

DETERMINISTIC WALKS IN RANDOM ENVIRONMENT

ROMAIN AIMINO AND CARLANGEOLO LIVERANI

ABSTRACT. Motivated by the random Lorentz gas, we study deterministic walks in random environment and show that (in simple, yet relevant, cases) they can be reduced to a class of random walks in random environment where the jump probability depends (weakly) on the past. In addition, we prove few basic results (hopefully the germ of a general theory) on the latter, purely probabilistic, model.

1. INTRODUCTION

The motion of a point particle among periodically distributed elastically reflecting convex bodies has been intensively studied for many years, both in the case of diluted obstacles and in the opposite case of high density. Diluted obstacles (the Boltzmann-Grad limit) can be treated by kinetic theory and lattice dynamics ideas (see [20] and related work). The high density case has required more specific dynamical systems tools, starting with the seminal work of Bunimovich, Sinai, Chernov [2] until the recent and much more precise results obtained in [7]. In particular, the latter result uses the new standard pairs and martingale problem techniques introduced in the field by Dolgopyat (see [6] for an elementary introduction to such ideas and references to the original works).

All the above deal with the periodic case, yet any material is expected to have defects. Hence, the study of obstacles distributed according to a random, translation invariant process is of paramount importance.

Unfortunately very few results are available in the non periodic case with the notable exception of the low density regime (Boltzmann-Grad limit), see [19]. For high obstacle density the only results are [10, 11], where recurrence is proven for special examples, and [8], where the Central Limit Theorem is proven when the periodicity is broken only in a finite region (hence translation invariance is violated).

2000 *Mathematics Subject Classification.* 60J15, 37A25, 37C30.

Key words and phrases. Random Lorentz gas, Random walk in random environment, decay of correlations.

L.C. thanks Dmitry Dolgopyat and Marco Lenci for several discussions on random walks through the years. Also we would like to thank Serge Troubetzkoy for pointing out the possibility that, in our setting, the ergodicity of the process as seen from the particle might hold under very weak extra assumptions. We also thank the anonymous referee for several helpful suggestions. This work was partially supported by the European Advanced Grant Macroscopic Laws and Dynamical Systems (MALADY) (ERC AdG 246953). L.C. acknowledges the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C18000100006. This work started while R.A. was affiliated to Università di Roma (Tor Vergata). R.A. was partially supported by FCT fellowship SFRH/BPD/123630/2016, by FCT projects PTDC/MAT-CAL/3884/2014 and PTDC/MAT-PUR/28177/2017, with national funds, and by CMUP (UID/MAT/00144/2019), which is funded by FCT with national (MCTES) and European structural funds through the programs FEDER, under the partnership agreement PT2020.

However some basic results hold in full generality: [3, 23] establish a criteria for recurrence and [10] shows that recurrent systems are ergodic. For example, by the criteria in [3, 23], the problems of recurrence and ergodicity are reduced, in the two dimensional case, to establishing a CLT, see [11] for details. Also, establishing recurrence in the case of a one dimensional array of obstacles (tubes) is substantially simpler and has been obtained in [5, 4, 25, 13]. Nevertheless, even in the simpler one dimensional situation, the study of rate of mixing and CLT is wide open.

Part of the difficulty in studying the above problems stems from the fact that, on the one hand, one needs non trivial results for the dynamical system and, on the other hand, one has to overcome the same obstacles that exist in analysing the problems of random walks in random environment (see [27] for a review on the subject). In particular, on the dynamical system side it seems necessary to establish some type of *memory loss*. In other words, one must show that the deterministic dynamics is akin to a random process.

In this article we aim at separating the above two difficulties, so that they can be (hopefully) solved independently. To this end we investigate more general models: a) deterministic random walks in random environment (see section 3); b) random walks (with memory) in random environment (see section 4). The former include the random Lorentz gases; the latter include *persistent random walks* but allows for infinite memory.

These two classes of models are connected by the following conjecture: relevant classes of deterministic walks in random environment are equivalent to the above purely probabilistic models. Hence, e.g., establishing the CLT for the probabilistic model implies the CLT for the deterministic walk.

1.1. Results and structure of the paper. The paper is organised as follows: in the next section we discuss briefly the Lorentz gases establishing, in the process, several notations needed in the following. In Section 3 we describe a general class of deterministic walks in random environment which encompass both the Lorentz gas and the example with a simpler dynamics that we will consider later.

In Section 4 we describe the class of random systems. We establish properties under which one can prove the ergodicity of the random process as seen from the particle. In particular, if one would succeed in reducing the Lorentz gas to such probabilistic model, then one would automatically recover all the known results. Of course, we believe that much more would follow from such a reduction. As a first step, in Section 5, we prove the reversibility of the associated probabilistic model. In the same section we show that if one restricts the dynamics to a Markov one dimensional expanding map, then the dynamical part of the problem can be completely obliterated and one is lead to known purely probabilistic models.

Next, in Section 6, we explore more realistic (but still one dimensional) models. We explain under which conditions the deterministic dynamics can be obliterated yielding a probabilistic model as in Section 4. In Section 7 we verify such conditions for a class of one dimensional non-Markov expanding maps. In the following sections we prove the statements of Section 6.



Notation. *In the following we will use $C_{\#}$ to designate a generic constant that depends only on the parameters or the assumptions of the considered model. The actual value of $C_{\#}$ is immaterial. In particular, the value of $C_{\#}$ can change from one occurrence to the next.*

2. THE RANDOM LORENTZ GAS

The random Lorentz gas consists of a distribution of convex, non overlapping, obstacles in \mathbb{R}^d , $d \geq 2$ and independent point particles that move of free motion and collides elastically with the obstacles. If one describes the particle density by a distribution, then the problem is reduced to studying the motion of one particle with an initial distribution given by a measure. If the obstacle distribution is described by some probability measure, then the goal is to study the dynamics of the particle for almost all obstacle distributions.

Of course, the problem depends heavily on the obstacle distribution. Two reasonable assumptions are that the distribution enjoys some type of stationarity and ergodicity with respect to some subgroup of the space translations. Also a key issue is the existence or not of trajectories that can spend an unboundedly long time without experiencing any collision. Given the many possibilities, let us restrict to small perturbations of a periodic array of discs on \mathbb{Z}^2 . Note however that similar examples can be considered for $d = 1$ (Lorentz tubes [5, 4, 25]) and $d > 2$.

Following [11] we start with an exactly periodic distribution of discs of radius R centred on the square lattice $\mathbb{R}^2/\mathbb{Z}^2$, such a periodic array divides naturally \mathbb{R}^2 in cells. In each cell we assume that there is another disc of radius r with center in a δ -neighborhood of the center of the cell. We assume that $0 < R < 1/2$ and $2/\sqrt{2} - R - \delta > r > 1 - 2R + \delta$ which insures that the obstacles do not overlap and that the trajectories cannot enter and leave a cell without experiencing a collision. The locations of the central obstacle in different cells is an i.i.d. random variable. See Figure 1.a for an illustration.

Fig 1.a *Random obstacle configuration*Fig 1.b *Poincaré section (in bold)*

The above is a reasonable model for a material with a periodic structure and random impurities. Let $\|\omega_z\| \leq \delta$ be the displacement of the center obstacle from the center of the cell $z \in \mathbb{Z}^2$. The obstacles are distributed according to a product measure \mathbb{P} on $\Omega = \{\omega \in \mathbb{R}^2 : \|\omega\| \leq \delta\}^{\mathbb{Z}^2}$. On Ω are defined the translations $\tau_z : \Omega \rightarrow \Omega$ by $z \in \mathbb{Z}^2$: for all $\bar{\omega} \in \Omega$ and $w \in \mathbb{Z}^2$

$$\tau_z(\bar{\omega})_w = (\bar{\omega})_{w+z}.$$

Next, consider a particle with position $q \in \mathbb{R}^2$ and velocity p , $\|p\| = 1$, moving among the obstacles with elastic collisions.

The dynamics is deterministic but we consider stochastic initial conditions: Assume w.l.o.g. that the particle starts from the zero cell with positions and velocity described by a smooth distribution h_0 .

Let $(q(t), p(t))$ be the position and velocity at time t . Natural questions are: does it exist an asymptotic velocity (Law of large numbers)

$$(2.1) \quad V = \lim_{t \rightarrow \infty} t^{-1}[q(t) - q(0)].$$

If so, does the Central Limit Theorem hold? That is, does

$$(2.2) \quad \frac{1}{\sqrt{t}}[q(t) - q(0) - Vt]$$

converges in law to a Gaussian Random Variable \mathbb{P} a.s. (quenched CLT).

2.1. Poincaré section. Consider a single cell (see Figure 1.b) and let $C = \cup_{i=1}^4 C_i$. C_i being one of the bold arcs centered at the corners of the box, and let $G_{(0,0)} = C \times [-\pi/2, \pi/2]$; $[-\pi/2, \pi/2]$ being the angle that the post-collisional velocity forms with the external normal. Next, let $\{B_i\}_{i=1}^4$ be the segments constituting the boundary of the box not contained in the obstacles (see Figure 1.b) and set $G_{(1,0)} = B_1 \times [-\pi/2, \pi/2]$, $G_{(-1,0)} = B_2 \times [-\pi/2, \pi/2]$, $G_{(0,1)} = B_3 \times [-\pi/2, \pi/2]$ and $G_{(0,-1)} = B_4 \times [-\pi/2, \pi/2]$. Next, let $\mathcal{W} = \{(0, 0), (\pm 1, 0), (0, \pm 1)\}$, $\mathcal{B} = \cup_{w \in \mathcal{W}} G_w$ and consider the phase space $\mathbb{B} = \mathcal{B} \times \mathbb{Z}^2$. This is a Poincaré section.

For each $\bar{\omega}$, given a particle at $(x, z) \in \mathbb{B}$ follow its motion till it hits \mathcal{B} again, this defines a map $f_{\bar{\omega}_z}(x)$. The Poincaré map $\mathbb{F}_{\bar{\omega}} : \mathbb{B} \rightarrow \mathbb{B}$ associated to the billiard flows is then defined by

$$(2.3) \quad \begin{aligned} \mathbb{F}_{\bar{\omega}}(x, z) &= (f_{\bar{\omega}_{z+e(\bar{\omega}_z, x)}}(x), z + e(\bar{\omega}_z, x)) \\ e(\bar{\omega}_z, x) &= \sum_{w \in \mathcal{W}} \mathbf{1}_{G_w}(x)w, \end{aligned}$$

where $\mathbf{1}_B$ is the characteristic function of the set B .

2.2. The random process. For each initial density h_0 and $\bar{\omega} \in \Omega$ we have thus defined the random variables¹

$$(x(\bar{\omega}, n), z(\bar{\omega}, n)) = \mathbb{F}_{\bar{\omega}}^n(x, z).$$

Let $\mathbb{M} = \{(z(n)) \in (\mathbb{Z}^2)^{\mathbb{N}} : z(0) = 0, z(n+1) - z(n) \in \mathcal{W}\}$ be the path space and consider the random process on $\Omega_{\star} = \Omega \times \mathbb{M}$ defined as follows. For each $\bar{\omega} \in \Omega$ define the conditional probability

$$(2.4) \quad \mathbb{P}_{\star}(\{z(1), \dots, z(n)\} | \bar{\omega}) = \int_{\mathcal{B}} \prod_{k=0}^{n-1} \mathbf{1}_{G_{w(k)}}(x(\bar{\omega}, k)) h_0(x) dx,$$

where $w(k) = z(k+1) - z(k)$. While $\bar{\omega}$ is distributed according to \mathbb{P} .

Next, we introduce the transfer operators²

$$\mathcal{L}_{\bar{\omega}, z, w} \varphi(x) = \sum_{\{y : f_{\bar{\omega}_{z+w}}(y) = x\}} |\det \partial_y f_{\bar{\omega}_{z+w}}(y)|^{-1} \mathbf{1}_{G_w}(y) \varphi(y).$$

Changing variable repeatedly, yields

$$(2.5) \quad \mathbb{P}_{\star}(\{z(1), \dots, z(n)\} | \bar{\omega}) = \int_{\mathcal{B}} \mathcal{L}_{\bar{\omega}, z(n-1), w(n-1)} \cdots \mathcal{L}_{\bar{\omega}, z(0), w(0)} h_0.$$

¹Using the probabilistic usage we will often suppress the $\bar{\omega}$ dependency, when this does not create confusion.

²In the present case the set on which we take the sum consists of only one element and the determinant of the Jacobian of the $f_{\bar{\omega}}$ is always one.

Thus the measure \mathbb{P}_\star is expressed as products of transfer operators. The process \mathbb{P}_\star is, in general, not Markov, however we conjecture:

Conjecture. *Under some appropriate technical conditions on \mathbb{P} and h_0 , there exist $\nu \in (0, 1)$ and $C_\# > 0$ such that for all $\bar{\omega} \in \Omega$*

$$|\mathbb{P}_\star(z(n) \mid z^n, \bar{\omega}) - \mathbb{P}_\star(\hat{z}_m(n-m) \mid \hat{z}^{n-m}, \tau_{z(m)}\bar{\omega})| \leq C_\# \nu^{n-m},$$

where $z^k = z(1), \dots, z(k-1)$, $\hat{z}^k = \hat{z}_m(1), \dots, \hat{z}_m(k-1)$ with $\hat{z}_m(k) = z(m+k) - z(m)$.

If the above were true, then one could reduce the study of the dynamical system to a purely probabilistic problem, albeit non an easy one. See however Section 4 for a beginning of a theory in some related cases.

Next, we introduce a large class of dynamical systems, generalising the Lorentz models. Then, we will prove the above conjecture for simple, and not so simple, dynamical systems, giving an idea of what *some appropriate technical conditions* might mean (see Theorem 6.1).

3. DETERMINISTIC WALKS IN RANDOM ENVIRONMENT

As already mentioned there is an extreme scarcity of results pertaining the high density random Lorentz Gas. It is then sensible to consider simpler models in which one can start to solve some of the outstanding difficulties. To this end, following Lenci [11], it is convenient to see the Lorentz gas as a special case of a more general class of models: *deterministic walks in random environment*. Even for such models very few results exist. Exceptions are [14] in which a zero-one law for systems with local dynamics which are markovian, but deterministic, is established and [26] which considers statistical properties for a related (simplified) model with local dynamics consisting of expanding linear maps of the circle or hyperbolic toral automorphisms.

The model can be stated in rather general terms, for simplicity let us restrict to the case of the \mathbb{Z}^d lattice with bounded jumps $\mathcal{W} \subset \mathbb{Z}^d$, $\#\mathcal{W} < \infty$,³ and local dynamics which live all on the same phase space \mathcal{M} .

Consider the set $\mathcal{A} = \{(f_\alpha, \mathcal{M}, \mathcal{G}_\alpha)\}_{\alpha \in A}$, where (A, \mathcal{S}) is a measurable space⁴, $f_\alpha : \mathcal{M} \rightarrow \mathcal{M}$ are maps such that $(\alpha, x) \in A \times \mathcal{M} \mapsto f_\alpha(x) \in \mathcal{M}$ is measurable, and $\mathcal{G}_\alpha = \{G_{\alpha, w}\}_{w \in \mathcal{W}}$ are partitions of \mathcal{M} such that, for each $w \in \mathcal{W}$, the map $(\alpha, x) \mapsto \mathbf{1}_{G_{\alpha, w}}(x)$ is measurable. The environment is described by the probability space $\Omega = A^{\mathbb{Z}^d}$ equipped with a translation invariant probability \mathbb{P} . Also we assume that all the maps $f_\alpha : \mathcal{M} \rightarrow \mathcal{M}$ are nonsingular with respect to some reference measure m on \mathcal{M} . Then, for each realisation $\bar{\omega} \in \Omega$ we can define the dynamics $\mathbb{F}_{\bar{\omega}}(\cdot, \cdot) : \mathcal{M} \times \mathbb{Z}^d \rightarrow \mathcal{M} \times \mathbb{Z}^d$:

$$\begin{aligned} \mathbb{F}_{\bar{\omega}}(x, z) &= (f_{\bar{\omega}_{z+e(\bar{\omega}_z, x)}}(x), z + e(\bar{\omega}_z, x)) \\ e(\alpha, x) &= \sum_{w \in \mathcal{W}} \mathbf{1}_{G_{\alpha, w}}(x)w, \end{aligned}$$

and $(x(n), z(n)) = \mathbb{F}_{\bar{\omega}}^n(x(0), z(0))$. The randomness at fixed environment rests in the initial condition: $z(0) = z_0$ while $x(0)$ is distributed according to some probability

³ That is only jumps $w \in \mathcal{W}$ are allowed.

⁴For most of our concrete applications, (A, \mathcal{S}) will simply be a finite set with the discrete σ -algebra.

measure μ absolutely continuous with respect to m . We will assume, w.l.o.g., $z_0 = 0$. Then the path $(z(n))_{n \in \mathbb{N}}$ belongs to $\mathbb{M} = \{(z_n) \in (\mathbb{Z}^d)^{\mathbb{N}} : z_0 = 0, z_{n+1} - z_n \in \mathcal{W}\}$, which we call the space of *admissible paths*, and \mathbb{P}_* is the law of the resulting process on $\Omega_* = \Omega \times \mathbb{M}$.

Each f_α admits a transfer operator $\mathcal{L}_{f_\alpha} : L^1(m) \rightarrow L^1(m)$ defined by

$$(3.1) \quad \int_{\mathcal{M}} (\mathcal{L}_{f_\alpha} \phi) \psi dm = \int_{\mathcal{M}} \phi \cdot \psi \circ f_\alpha dm$$

for all $\phi \in L^1(m)$ and $\psi \in L^\infty(m)$.

Let h_0 be the density of the initial condition ($d\mu = h_0 dm$). Then, setting $w(n) = z(n+1) - z(n)$, a repeated use of (3.1) in (2.4) yields

$$(3.2) \quad \mathbb{P}_*(z(1), \dots, z(n) \mid \bar{\omega}) = \int_{\mathcal{M}} \mathcal{L}_{\bar{\omega}, z(n-1), w(n-1)} \cdots \mathcal{L}_{\bar{\omega}, z(0), w(0)} h_0 dm$$

for each $\bar{\omega} \in \Omega$ and each admissible path $(z(n)) \in \mathbb{M}$, and with

$$(3.3) \quad \mathcal{L}_{\bar{\omega}, z, w}(\phi) = \mathcal{L}_{f_{\bar{\omega}_z + w}}(\mathbf{1}_{G_{\bar{\omega}_z, w}} \phi).$$

3.1. The point of view of the particle. An important process associated to our deterministic walk is the process of the environment as seen from the particle. This is the dynamical system defined on $\Omega \times \mathcal{M}$ by

$$\mathcal{F}(\bar{\omega}, x) = (\tau_{e(\bar{\omega}_0, x)} \bar{\omega}, f_{\bar{\omega}_e(\bar{\omega}_0, x)}(x)).$$

This is known to be a fruitful point of view, in particular if one knows the invariant measure. As noted by Lenci [10, 11], in an important subclass of deterministic random walks the invariant measure can be trivially computed.

Lemma 3.1. *If all the maps f_α have the same invariant measure λ and the set \mathcal{G}_α is deterministic (i.e., it does not depend on α), then the probability measure $\mathbb{P}_0 = \mathbb{P} \times \lambda$ is invariant for the map \mathcal{F} .*

Proof. Let \mathbb{E}_0 be the expectation with respect to \mathbb{P}_0 . Since the set \mathcal{G}_α is deterministic, we can write $e(\alpha, x) = e(x)$ for all $\alpha \in A$ and $x \in \mathcal{M}$. For each bounded measurable function φ we have

$$\begin{aligned} \mathbb{E}_0(\varphi \circ \mathcal{F}) &= \int \varphi(\tau_{e(x)} \bar{\omega}, f_{\bar{\omega}_{e(x)}}(x)) \mathbb{P}(d\bar{\omega}) \lambda(dx) \\ &= \int \varphi(\tau_{e(x)} \bar{\omega}, f_{(\tau_{e(x)} \bar{\omega})_0}(x)) \mathbb{P}(d\bar{\omega}) \lambda(dx) \\ &= \int \varphi(\bar{\omega}, f_{\bar{\omega}_0}(x)) \mathbb{P}(d\bar{\omega}) \lambda(dx) \\ &= \int \varphi(\bar{\omega}, x) \mathbb{P}(d\bar{\omega}) \lambda(dx) = \mathbb{E}_0(\varphi), \end{aligned}$$

where we have used first the invariance of \mathbb{P} with respect to the translations and then the invariance of λ with respect to the maps f_α . \square

Hence now we have a dynamical system with a finite measure.

Lemma 3.2. *In the hypotheses of Lemma 3.1 the limit*

$$V = \lim_{n \rightarrow \infty} \frac{1}{n} z(n)$$

exists \mathbb{P}_0 a.s.. Moreover, if $(\Omega \times \mathcal{M}, \mathcal{F}, \mathbb{P}_0)$ is ergodic, then $V = \lambda(e)$.

Proof. Setting $(\bar{\omega}(n), x(n)) = \mathcal{F}^n(\bar{\omega}, x)$ yields $\frac{1}{n}z(n) = \frac{1}{n} \sum_{k=0}^{n-1} e(x(k))$. Hence the existence of the limit follows from Birkhoff ergodic theorem for the dynamical system $(\Omega \times \mathcal{M}, \mathcal{F}, \mathbb{P}_0)$ applied to $\varphi(\bar{\omega}, x) = e(\bar{\omega}_0, x) = e(x)$. By ergodicity, the limit equals the average of φ with respect to \mathbb{P}_0 which is equal to $\mathbb{E}_0(\varphi) = \lambda(e)$. \square

As an application we have the following Lemma (which is implicit in [5]).

Lemma 3.3. *For the Lorentz gas described in section 2 we have $V = 0$.*

Proof. In [11, Theorem 5.4] is proven that $(\Omega \times \mathcal{M}, \mathcal{F}, \mathbb{P}_0)$ is ergodic. Then Lemma 3.2 implies that $V = \sum_{w \in \mathcal{W}} w \lambda(G_w)$ which average is zero due to the fact that λ is invariant for the change $p \rightarrow -p$. \square

4. GIBBS RANDOM WALKS IN RANDOM ENVIRONMENT

We do not expect the process described by \mathbb{P}_* to be Markov, yet we expect that the jump rates have a weak dependence of the past, provided the maps are strongly chaotic. To be more precise, we conjecture that the process is a *random walk in random environment with weak memory*. In the probabilistic literature random walks with a finite memory are called *persistent*, here we expect the memory to be infinite although depending weakly on the far past, like a potential of a Gibbs measure. Let us specify exactly what we mean by this.

Let (A, \mathcal{S}) be a measurable space and $\mathcal{W} \subset \mathbb{Z}^d$, $2 \leq \#\mathcal{W} < \infty$. Consider the measurable space $\Omega = A^{\mathbb{Z}^d}$ and a translation invariant, ergodic, probability distribution \mathbb{P} that describes the distribution of the environments $\bar{\omega} \in \Omega$. For each $n \in \mathbb{N}$ and $(w_0, \dots, w_{n-1}) \in \mathcal{W}^n$, assume that are given compatible probabilities $p(\bar{\omega}, n, \cdot)$ on \mathcal{W}^n , i.e.

$$\sum_{(w_0, \dots, w_{n-1}) \in \mathcal{W}^n} p(\bar{\omega}, n, w_0 \dots w_{n-1}) = 1,$$

and

$$p(\bar{\omega}, n, w_0 \dots w_{n-1}) = \sum_{w \in \mathcal{W}} p(\bar{\omega}, n+1, w_0 \dots w_{n-1}w)$$

for all $\bar{\omega} \in \Omega$, $n \geq 0$ and $(w_0, \dots, w_{n-1}) \in \mathcal{W}^n$. Assume also that all the maps $\bar{\omega} \mapsto p(\bar{\omega}, n, w_0 \dots w_{n-1})$ are measurable. We have then for each $\bar{\omega} \in \Omega$ a probability measure $\mathbb{P}^{\bar{\omega}}$ on the space $\mathcal{W}^{\mathbb{N}}$ by Kolmogorov extension theorem. By the monotone class theorem, the map $G \mapsto \mathbb{P}^{\bar{\omega}}(G)$ is measurable for any measurable set $G \subset \mathcal{W}^{\mathbb{N}}$, and we can thus define a probability measure \mathbb{P}_* on $\Omega \times \mathcal{W}^{\mathbb{N}}$ by

$$\mathbb{P}_*(d\bar{\omega}, d\bar{w}) = \mathbb{P}(d\bar{\omega}) \mathbb{P}^{\bar{\omega}}(d\bar{w}).$$

Remark 4.1. *The measure \mathbb{P}_* can be naturally identified with a measure on the space $\Omega_* = \Omega \times \mathbb{M}$, where $\mathbb{M} = \{(z_n) \in (\mathbb{Z}^d)^{\mathbb{N}} : z_0 = 0, z_{n+1} - z_n \in \mathcal{W}, \forall n\}$ is the space of admissible paths starting at 0, since there is a 1-to-1 correspondence between elements of \mathcal{W}^n and admissible paths of length n , via the relations $w_k = z_{k+1} - z_k$.*

4.1. The weak memory requirement. We find convenient, although not strictly necessary, to require the following assumption that ensures that all admissible paths have positive probability:

(Pos): for \mathbb{P} -a.e. $\bar{\omega} \in \Omega$, for all $n \geq 0$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$,

$$p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) > 0.$$

Let $\mathbb{P}_{\bar{\omega}}(\bar{w}_n \mid \bar{w}_0 \dots \bar{w}_{n-1})$ be the conditional probability $\frac{p(\bar{\omega}, n+1, \bar{w}_0 \dots \bar{w}_{n-1} \bar{w}_n)}{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}$.

The weak memory requirement is made precise by the following:

(Exp): there exist $C_* > 0$ and $\nu \in (0, 1)$ such that for \mathbb{P} -a.e. $\bar{\omega} \in \Omega$, all $n > m \geq 0$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$

$$(4.1) \quad \left| \mathbb{P}_{\bar{\omega}}(\bar{w}_n \mid \bar{w}_0 \dots \bar{w}_{n-1}) - \mathbb{P}_{\tau_{z_m} \bar{\omega}}(\bar{w}_n \mid \bar{w}_m \dots \bar{w}_{n-1}) \right| \leq C_* \nu^{n-m},$$

where $z_m = \sum_{k=0}^{m-1} \bar{w}_k$.

Remark 4.2. *This is the property we have conjectured to be true for the random Lorentz gas at the end of Section 2.*

Note that the above condition implies loss of memory:

Lemma 4.3. *If \mathbb{P}_* satisfies (4.1), then (using the notation of Remark 4.1),*

$$\left| \mathbb{P}_*(z_{n+1} \mid z_{n-m+1}, \dots, z_n, \bar{\omega}) - \mathbb{P}_*(z_{n+1} \mid z_1, \dots, z_n, \bar{\omega}) \right| \leq C_{\#} \nu^m.$$

Proof. Let $\mathcal{P}(z_{n-m+1})$ be the set of admissible paths (w_0, \dots, w_{n-m}) that arrive in z_{n-m+1} , i.e. $\sum_{i=0}^{n-m} w_i = z_{n-m+1}$, and set $w_i = z_{i+1} - z_i$ for $i = n-m+1, \dots, n$. Then we have, using **(Exp)**:

$$\begin{aligned} \mathbb{P}_*(z_{n+1} \mid z_{n-m+1}, \dots, z_n, \bar{\omega}) &= \sum_{w \in \mathcal{P}(z_{n-m+1})} \frac{p(\bar{\omega}, n+1, w_0 \dots w_n)}{\mathbb{P}_*(z_{n-m+1}, \dots, z_n \mid \bar{\omega})} \\ &= \sum_{w \in \mathcal{P}(z_{n-m+1})} \mathbb{P}_{\bar{\omega}}(w_n \mid w_0 \dots w_{n-1}) \frac{p(\bar{\omega}, n, w_0 \dots w_{n-1})}{\mathbb{P}_*(z_{n-m+1}, \dots, z_n \mid \bar{\omega})} \\ &= \mathbb{P}_{\tau_{z_{n-m+1}} \bar{\omega}}(w_n \mid w_{n-m+1} \dots w_{n-1}) + \mathcal{O}(\nu^m), \end{aligned}$$

and

$$\begin{aligned} \mathbb{P}_*(z_{n+1} \mid z_1, \dots, z_n, \bar{\omega}) &= \mathbb{P}_{\bar{\omega}}(w_n \mid w_0 \dots w_{n-1}) \\ &= \mathbb{P}_{\tau_{z_{n-m+1}} \bar{\omega}}(w_n \mid w_{n-m+1} \dots w_{n-1}) + \mathcal{O}(\nu^m). \end{aligned}$$

□

4.2. The point of view of the particle and three further assumptions. One can define the process of the environment as seen from the particle. It is given by the dynamical system on the space $\Omega_* = \Omega \times \mathcal{W}^{\mathbb{N}}$ defined by

$$\mathcal{F}_*(\bar{\omega}, \bar{w}) = (\tau_{\bar{w}_0} \bar{\omega}, \tau_* \bar{w}),$$

where $\tau_* : \mathcal{W}^{\mathbb{N}} \rightarrow \mathcal{W}^{\mathbb{N}}$ is the unilateral shift. Note that, in general, \mathbb{P}_* is not invariant for \mathcal{F}_* . Next, let us show that some easy properties of the Markov case persist in the present context, under reasonable extra conditions.

Remark 4.4. *Note that in the probabilistic literature, see for instance [27], it is more usual to consider the random process $\bar{\omega}(n) = \tau_{z(n)} \bar{\omega}$ on $\bar{\Omega} = \Omega^{\mathbb{N}}$, and its law $\bar{\mathbb{P}}$ when $\bar{\omega}(0)$ is distributed according to \mathbb{P} (as we did in Section 3.1). When the set of periodic environments has probability 0, these two points of view are equivalent, since the map which associates to each $(\bar{\omega}, \bar{w}) \in \Omega_*$ the corresponding sequence $(\bar{\omega}(n)) \in \bar{\Omega}$ is invertible almost everywhere and realizes a conjugacy between the two dynamical systems $(\Omega_*, \mathcal{F}_*, \mathbb{P}_*)$ and $(\bar{\Omega}, \bar{\tau}, \bar{\mathbb{P}})$, where $\bar{\tau}$ is the shift on $\bar{\Omega}$. The same comments hold also for the definition given in Section 3.1. See also Remark 4.24 for further comments.*

We are interested in the asymptotic properties for z_n . Note that if we define $\varphi(\bar{\omega}, \bar{w}) = \bar{w}_0$, then we have $z_n = \sum_{k=0}^{n-1} \varphi \circ \mathcal{F}_*^k$.

The next assumption will imply the existence of a \mathcal{F}_* -invariant measure:

(Abs): There exists $C_0 > 0$ such that for \mathbb{P} -a.e. $\bar{\omega} \in \Omega$, all $n \geq 0$, all $k \geq 1$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$,

$$C_0^{-1} \leq \sum_{(w^1, \dots, w^k) \in \mathcal{W}^k} \frac{p(\tau_{-(w^1 + \dots + w^k)} \bar{\omega}, n + k, w^1 \dots w^k \bar{w}_0 \dots \bar{w}_{n-1})}{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})} \leq C_0.$$

To legitimize this assumption, we prove two relevant facts.

Lemma 4.5. *If there exists a probability measure \mathbb{Q}_* equivalent to \mathbb{P}_* , invariant for \mathcal{F}_* and such that $c^{-1} \leq \frac{d\mathbb{Q}_*}{d\mathbb{P}_*} \leq c$ for some $c > 0$, then **(Abs)** holds with $C_0 = c^2$. In particular, **(Abs)** holds if \mathbb{P}_* is \mathcal{F}_* -invariant.*

Proof. For all measurable set $B \subset \Omega$ and all cylinder $[\bar{w}_0 \dots \bar{w}_{n-1}] \subset \mathcal{W}^{\mathbb{N}}$, the preimage $\mathcal{F}_*^{-k}(B \times [\bar{w}_0, \dots, \bar{w}_{n-1}])$ is equal to the disjoint union

$$\bigcup_{(w^1, \dots, w^k) \in \mathcal{W}^k} \tau_{-(w^1 + \dots + w^k)}(B) \times [w^1 \dots w^k \bar{w}_0 \dots \bar{w}_{n-1}].$$

Let $B' = B \times [\bar{w}_0 \dots \bar{w}_{n-1}]$. By definition of \mathbb{P}_* , we have

$$\mathbb{P}_*(B') = \int_B p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega})$$

and

$$\begin{aligned} \mathbb{P}_*(\mathcal{F}_*^{-k} B') &= \sum_{(w^1, \dots, w^k) \in \mathcal{W}^k} \int_{\tau_{-(w^1 + \dots + w^k)} B'} p(\bar{\omega}, n + k, w^1 \dots w^k \bar{w}_0 \dots \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}) \\ &= \int_B \sum_{(w^1, \dots, w^k) \in \mathcal{W}^k} p(\tau_{w^1 + \dots + w^k} \bar{\omega}, n + k, w^1 \dots w^k \bar{w}_0 \dots \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}) \end{aligned}$$

thanks to the translation invariance of \mathbb{P} .

Since $\mathbb{P}_*(B') \leq c \mathbb{Q}_*(B') = c \mathbb{Q}_*(\mathcal{F}_*^{-k} B') \leq c^2 \mathbb{P}_*(\mathcal{F}_*^{-k} B')$, we have

$$\begin{aligned} &\int_B p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}) \\ &\leq c^2 \int_B \sum_{(w^1, \dots, w^k) \in \mathcal{W}^k} p(\tau_{w^1 + \dots + w^k} \bar{\omega}, n + k, w^1 \dots w^k \bar{w}_0 \dots \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}), \end{aligned}$$

and similarly,

$$\begin{aligned} &\int_B \sum_{(w^1, \dots, w^k) \in \mathcal{W}^k} p(\tau_{w^1 + \dots + w^k} \bar{\omega}, n + k, w^1 \dots w^k \bar{w}_0 \dots \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}) \\ &\leq c^2 \int_B p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}). \end{aligned}$$

Since the set of cylinders is countable, this proves the lemma. \square

We say that the process is *reversible* if \mathcal{W} is symmetric (i.e. $-w \in \mathcal{W}$ for all $w \in \mathcal{W}$) and, for \mathbb{P} -a.e. $\bar{\omega} \in \Omega$, all $n \geq 0$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$:

$$p(\bar{\omega}, n, w_0 \dots w_{n-1}) = p(\tau_{(w_0 + \dots + w_{n-1})} \bar{\omega}, n, -w_{n-1}, \dots, -w_0).$$

Note that this definition of reversibility is more akin to the one used in Dynamical Systems than the one used in Markov processes (the self-adjointness of the generator). However it does have relevant implications.

Lemma 4.6. *If the process is reversible, then \mathbb{P}_\star is \mathcal{F}_\star -invariant. In particular, (Abs) is verified, by Lemma 4.5.*

Proof. It suffices to consider a measurable set B' of the form $B' = B \times [\bar{w}_0, \dots, \bar{w}_{n-1}]$. We have:

$$\begin{aligned}
\mathbb{P}_\star(\mathcal{F}_\star^{-1}B') &= \mathbb{P}_\star\left(\bigcup_{w \in \mathcal{W}} \tau_{-w}(B) \times [w, \bar{w}_0, \dots, \bar{w}_{n-1}]\right) \\
&= \sum_{w \in \mathcal{W}} \int_{\tau_{-w}(B)} p(\bar{\omega}, n+1, w, \bar{w}_0, \dots, \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}) \\
&= \sum_{w \in \mathcal{W}} \int_B p(\tau_{-w}\bar{\omega}, n+1, w, \bar{w}_0, \dots, \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}) \\
&= \sum_{w \in \mathcal{W}} \int_B p(\tau_{(\bar{w}_0 + \dots + \bar{w}_{n-1})}\bar{\omega}, n+1, -\bar{w}_{n-1}, \dots, -\bar{w}_0, -w) \mathbb{P}(d\bar{\omega}) \\
&= \int_B p(\tau_{(\bar{w}_0 + \dots + \bar{w}_{n-1})}\bar{\omega}, n, -\bar{w}_{n-1}, \dots, -\bar{w}_0) \mathbb{P}(d\bar{\omega}) \\
&= \int_B p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) \mathbb{P}(d\bar{\omega}) = \mathbb{P}_\star(B').
\end{aligned}$$

□

Remark 4.7. *Lemma 4.5 suggests that (Abs) may be too strong. Yet, there are simple models (e.g. Sinai walk, see Example 1 in Section 5.2) for which there does not exist an invariant probability measure absolutely continuous with respect to \mathbb{P}_\star . Hence some condition is necessary.*

As common for random walks, we require an ellipticity assumption:

(Ell): There exist $\gamma_0 > 0$ and $n_\star \geq 0$ such that for \mathbb{P}_\star -a.e. $\bar{\omega} \in \Omega$, all $n \geq n_\star$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$,

$$\mathbb{P}_{\bar{\omega}}(\bar{w}_n \mid \bar{w}_0 \dots \bar{w}_{n-1}) \geq \gamma_0.$$

We finally formulate an assumption on the probability measure \mathbb{P} :

(Pro): Let $\mathcal{V}(\mathcal{W}) = \{z \in \mathbb{Z}^d : z = w_0 + \dots + w_{n-1}, n \in \mathbb{N}, w_i \in \mathcal{W}\}$. We assume that $G = \mathcal{V}(\mathcal{W})$ is an additive group and that \mathbb{P} is ergodic with respect to the action of G .

Remark 4.8. *It might not be necessary to assume that $\mathcal{V}(\mathcal{W})$ is a group, but we will not pursue this direction, as the examples we have in mind satisfies the above assumption: e.g., $\mathcal{V}(\mathcal{W})$ is an additive group whenever \mathcal{W} is symmetric. We leave to the interested reader possible weakening of property (Pro). We nevertheless mention that a closer look at its proof reveals that Theorem 4.9 remains valid if \mathbb{P} is ergodic for each translation τ_z , $z \in \mathbb{Z}^d$, $z \neq 0$ (for instance if \mathbb{P} is mixing when $d = 1$, or if \mathbb{P} is i.i.d. when $d > 1$), without any extra assumption on \mathcal{W} .*

4.3. A few basic results.

The above assumptions are justified by the following Theorem. The rest of the section is devoted to its proof.

Theorem 4.9. *Suppose that the conditions **(Pos)**, **(Exp)**, **(Abs)**, **(Ell)** and **(Pro)** hold. Then there exists a unique \mathcal{F}_* -invariant probability measure \mathbb{Q}_* equivalent to \mathbb{P}_* and the dynamical system $(\Omega_*, \mathcal{F}_*, \mathbb{Q}_*)$ is ergodic. In particular, we have*

$$(4.2) \quad \lim_{n \rightarrow \infty} \frac{z_n}{n} = V,$$

\mathbb{P}_* -a.s., with $V = \int_{\Omega_*} \varphi d\mathbb{Q}_* \in \mathbb{R}^d$.⁵

Remark 4.10. *If \mathbb{P}_* is invariant, then*

$$V = \int_{\Omega_*} \varphi d\mathbb{P}_* = \sum_{w \in \mathcal{W}} w \int_{\Omega} p(\bar{\omega}, 1, w) \mathbb{P}(d\bar{\omega}).$$

As a simple but important consequence, we deduce the recurrence in 1-d:

Corollary 4.11. *Under the conditions of the above theorem, if $d = 1$ and $V = 0$, then the process (z_n) is recurrent: $z_n = 0$ infinitely often, \mathbb{P}_* -a.s..*

Proof of Corollary 4.11. We refer to [24] for a nice survey on the recurrence of cocycles. Since $(\Omega_*, \mathcal{F}_*, \mathbb{Q}_*)$ is ergodic by Theorem 4.9 when $d = 1$, the walk is recurrent if $V = 0$, see [1] or [24, Theorem 3]. Note that this result for recurrence of cocycles is stated for invertible dynamical systems, but it can be extended to non-invertible systems using the natural extension, see for instance [12, Appendix A.2]. \square

Remark 4.12. *When $d = 2$ and $V = 0$, if (z_n) satisfies an annealed central limit theorem, i.e. if $\frac{z_n}{\sqrt{n}}$ converges in law to a Gaussian distribution under the probability measure \mathbb{Q}_* , then the process (z_n) is recurrent by the results of Conze [3] and Schmidt [23].*

From now on and till the end of the section we will assume conditions **(Pos)**, **(Exp)**, **(Abs)**, **(Ell)** and **(Pro)** if not explicitly stated otherwise.

To prove Theorem 4.9, we will analyze the properties of the transfer operator \mathcal{L}_* associated to \mathcal{F}_* with respect to \mathbb{P}_* . More precisely, we will show that the operator \mathcal{L}_* enjoys some regularization properties on a space of Hölder functions. We first define the usual separation time on $\mathcal{W}^{\mathbb{N}}$ by

$$s(\bar{w}, \bar{w}') = \inf\{n \geq 0 : \bar{w}_n \neq \bar{w}'_n\},$$

and for $0 < \nu < 1$, the metric $d_\nu(\bar{w}, \bar{w}') = \nu^{s(\bar{w}, \bar{w}')}$ on $\mathcal{W}^{\mathbb{N}}$.

For a measurable function $f : \Omega \times \mathcal{W}^{\mathbb{N}} \rightarrow \mathbb{C}$, we set:

$$\begin{aligned} \|f\|_\infty &= \operatorname{ess\,sup}_{\bar{\omega} \in \Omega} \sup_{\bar{w} \in \mathcal{W}^{\mathbb{N}}} |f(\bar{\omega}, \bar{w})|, \\ |f|_\nu &= \operatorname{ess\,sup}_{\bar{\omega} \in \Omega} \sup_{\bar{w} \neq \bar{w}'} \frac{|f(\bar{\omega}, \bar{w}) - f(\bar{\omega}, \bar{w}')|}{d_\nu(\bar{w}, \bar{w}')}, \end{aligned}$$

and define

$$\mathcal{H}_\infty = \{f : \Omega \times \mathcal{W}^{\mathbb{N}} \rightarrow \mathbb{C} : \|f\|_\infty < \infty\},$$

⁵ Recall that $\varphi(\bar{\omega}, \bar{w}) = \bar{w}_0$.

$$\mathcal{H}_\nu = \{f : \Omega \times \mathcal{W}^{\mathbb{N}} \rightarrow \mathbb{C} : \|f\|_\nu := \|f\|_\infty + |f|_\nu < \infty\}.$$

The space \mathcal{H}_ν is a Banach algebra. The following result about density of \mathcal{H}_ν is based on very classical ideas, but we include it here for completeness:

Lemma 4.13. *For any function $\varphi \in L^1(\mathbb{P}_\star)$, there exists $(\varphi_\epsilon)_\epsilon \in \mathcal{H}_\nu$ such that $\varphi_\epsilon \rightarrow \varphi$ in $L^1(\mathbb{P}_\star)$. Moreover, if φ is bounded, $(\varphi_\epsilon)_\epsilon$ can be chosen such that $\sup_\epsilon \|\varphi_\epsilon\|_\infty < \infty$.*

Proof. We first consider the case where $\varphi = \mathbf{1}_B$ is the indicator function of a measurable set $B \subset \Omega_\star$. We endow Ω_\star with the metric $d_\star((\bar{\omega}, \bar{w}), (\bar{\omega}', \bar{w}')) = d_\Omega(\bar{\omega}, \bar{\omega}') + d_\nu(\bar{w}, \bar{w}')$, where d_Ω is any metric defining the product topology on Ω . The metric d_\star defines the product topology on Ω_\star . For any open set $O \subset \Omega_\star$, we define

$$\varphi_{k,O}(\bar{\omega}, \bar{w}) = \min\{k d_\star((\bar{\omega}, \bar{w}), \Omega_\star \setminus O), 1\}.$$

We clearly have $0 \leq \varphi_{k,O} \leq \varphi_{k+1,O} \leq \mathbf{1}_O \leq 1$, and $\lim_k \varphi_{k,O}(\bar{\omega}, \bar{w}) = \mathbf{1}_O(\bar{\omega}, \bar{w})$ for all $(\bar{\omega}, \bar{w}) \in \Omega_\star$. The function $\varphi_{k,O}$ is clearly lipschitzian with respect to the metric d_\star , which also implies that $\varphi_{k,O} \in \mathcal{H}_\nu$. Since \mathbb{P}_\star is a probability measure on the compact metric space Ω_\star , it is outer regular, see [22, Theorem 2.17]: for any Borel set $B \subset \Omega_\star$ and any $\epsilon > 0$, there exists an open set $O_\epsilon \subset \Omega_\star$ such that $B \subset O_\epsilon$ and $\mathbb{P}_\star(O_\epsilon \setminus B) \leq \epsilon$. By the dominated convergence theorem

$$\lim_{k \rightarrow \infty} \int_{\Omega_\star} |\varphi_{k,O_\epsilon} - \mathbf{1}_{O_\epsilon}| d\mathbb{P}_\star = 0.$$

We choose $k_\epsilon \geq 0$ such that $\int_{\Omega_\star} |\varphi_{k_\epsilon, O_\epsilon} - \mathbf{1}_{O_\epsilon}| d\mathbb{P}_\star \leq \epsilon$ and set $\varphi_\epsilon = \varphi_{k_\epsilon, O_\epsilon}$. By the above arguments, we have $\varphi_\epsilon \in \mathcal{H}_\nu$, $\|\varphi_\epsilon\|_\infty \leq 1$, and

$$\|\varphi_\epsilon - \mathbf{1}_B\|_{L^1(\mathbb{P}_\star)} \leq \|\varphi_\epsilon - \mathbf{1}_{O_\epsilon}\|_{L^1(\mathbb{P}_\star)} + \mathbb{P}_\star(O_\epsilon \setminus B) \leq 2\epsilon,$$

which proves the convergence in $L^1(\mathbb{P}_\star)$. Next, assume $\varphi \in L^\infty$. Without loss of generality we can assume $\varphi \geq 0$ and $\|\varphi\|_\infty = 2$. Let $B_1 = \{\xi \in \Omega_\star : \varphi(\xi) \geq 1\}$ and

$$B_k = \left\{ \xi \in \Omega_\star : \varphi(\xi) \geq 2^{-k} + \sum_{j=0}^{k-1} \mathbf{1}_{B_j} 2^{-j} \right\}.$$

By construction $\|\varphi - \sum_{j=0}^{k-1} \mathbf{1}_{B_j} 2^{-j}\|_\infty \leq 2^{-k}$, hence we can use the above approximations of the characteristic functions to approximate φ in L^1 with a sequence with norm bounded by 2. The case $\varphi \in L^1$ can be obtained by approximation by bounded functions. \square

Next, we state a useful technical lemma.

Lemma 4.14. *Under assumptions **(Pos)**, **(Exp)** and **(Ell)** there exists $\gamma_\star \in L^\infty(\Omega, \mathbb{P})$, $\gamma_\star > 0$ such that, for all $n > n_\star$ and $\bar{w} \in \mathcal{W}^{\mathbb{N}}$, we have*

$$\gamma_\star(\omega) \leq \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} \leq \gamma_\star(\omega)^{-1}$$

Proof. For all $n > n_*$, we have

$$\begin{aligned}
& \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} = \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\bar{\omega}, n-1, \bar{w}_0 \dots \bar{w}_{n-2})} \frac{p(\bar{\omega}, n-1, \bar{w}_0 \dots \bar{w}_{n-2})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} \\
& = \mathbb{P}_{\bar{\omega}}(\bar{w}_{n-1} \mid \bar{w}_0 \dots \bar{w}_{n-2}) \frac{p(\bar{\omega}, n-1, \bar{w}_0 \dots \bar{w}_{n-2})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} \\
& \geq \left[\mathbb{P}_{\tau_{\bar{w}_0} \bar{\omega}}(\bar{w}_{n-1} \mid \bar{w}_1 \dots \bar{w}_{n-2}) - C_{\#} \nu^n \right] \frac{p(\bar{\omega}, n-1, \bar{w}_0 \dots \bar{w}_{n-2})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} \\
& = \left[\frac{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-2, \bar{w}_1 \dots \bar{w}_{n-2})} - C_{\#} \nu^n \right] \frac{p(\bar{\omega}, n-1, \bar{w}_0 \dots \bar{w}_{n-2})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} \\
& = \left[1 - C_{\#} \nu^n \frac{p(\tau_{\bar{w}_0} \bar{\omega}, n-2, \bar{w}_1 \dots \bar{w}_{n-2})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} \right] \frac{p(\bar{\omega}, n-1, \bar{w}_0 \dots \bar{w}_{n-2})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-2, \bar{w}_1 \dots \bar{w}_{n-2})} \\
& \geq (1 - C_{\#} \gamma_0^{-1} \nu^n) \frac{p(\bar{\omega}, n-1, \bar{w}_0 \dots \bar{w}_{n-2})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-2, \bar{w}_1 \dots \bar{w}_{n-2})} \\
& \geq \prod_{j=n_*+1}^n (1 - C_{\#} \gamma_0^{-1} \nu^j) \frac{p(\bar{\omega}, n_*+1, \bar{w}_0 \dots \bar{w}_{n_*})}{p(\tau_{\bar{w}_0} \bar{\omega}, n_*, \bar{w}_1 \dots \bar{w}_{n_*})},
\end{aligned}$$

where we have used **(Exp)** at the third line and **(Ell)** at the last line. The lower bound follows then by **(Pos)**, provided that $C_{\#} \gamma_0^{-1} \nu^{n_*} < 1$, which we can always ensure by eventually redefining n_* . The upper bound can be established similarly.⁶ \square

Remark 4.15. Note that a slight strengthening of **(Pos)** would imply that γ_* can be chosen to be constant.⁷ Then Lemma 4.14 would imply: for each $k \in \mathbb{N}$, there exists $C_k > 0$ such that for \mathbb{P} -a.e. $\bar{\omega} \in \Omega$, all $n \geq 0$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$,

$$C_k^{-1} \leq \sum_{(w_1, \dots, w_k) \in \mathcal{W}^k} \frac{p(\tau_{-(w_1+\dots+w_k)} \bar{\omega}, n+k, w_1 \dots w_k \bar{w}_0 \dots \bar{w}_{n-1})}{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})} \leq C_k.$$

Hence the all point of **(Abs)** rests in the uniformity with respect to k .

We now define what will turn out to be the potential associated to \mathcal{L}_* :

Lemma 4.16. There exists a measurable function $J : \Omega \times \mathcal{W}^{\mathbb{N}} \rightarrow \mathbb{R}^+$ such that for all $n \geq 0$

$$(4.3) \quad \operatorname{ess\,sup}_{\bar{\omega} \in \Omega} \sup_{\bar{w} \in \mathcal{W}^{\mathbb{N}}} \left| \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} - J(\bar{\omega}, \bar{w}) \right| \leq C_{\#} \nu^n.$$

Moreover, J belongs to \mathcal{H}_ν and $J(\bar{\omega}, \bar{w}) > 0$ for \mathbb{P} -a.e. $\bar{\omega} \in \Omega$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$.

⁶ Note however that **(Ell)** is not needed to prove the upper bound.

⁷ That is, one could ask, for all $n > n_*$, $\inf_{\omega \in \Omega} p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) > 0$, which holds in all the example we have in mind.

Proof. Let $p_n(\bar{\omega}, \bar{w}) = \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})}$. Using **(Exp)**, we have

$$\begin{aligned} |p_n(\bar{\omega}, \bar{w}) - p_{n+1}(\bar{\omega}, \bar{w})| &= \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n, \bar{w}_1 \dots \bar{w}_n)} \\ &\quad \times \left| \frac{p(\tau_{\bar{w}_0} \bar{\omega}, n, \bar{w}_1 \dots \bar{w}_n)}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})} - \frac{p(\bar{\omega}, n+1, \bar{w}_0 \dots \bar{w}_n)}{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})} \right| \\ &= \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n, \bar{w}_1 \dots \bar{w}_n)} \left| \mathbb{P}_{\tau_{\bar{w}_0} \bar{\omega}}(\bar{w}_n \mid \bar{w}_1 \dots \bar{w}_{n-1}) - \mathbb{P}_{\bar{\omega}}(\bar{w}_n \mid \bar{w}_0 \dots \bar{w}_{n-1}) \right| \\ &\leq C_{\#} \nu^n \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n, \bar{w}_1 \dots \bar{w}_n)}. \end{aligned}$$

From **(Abs)**, substituting $\tau_{\bar{w}_0} \bar{\omega}$ to $\bar{\omega}$, we have

$$(4.4) \quad p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) \leq C_{\#} p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1}),$$

from which it follows, using **(Ell)**, for all $n \geq n_{\star}$,

$$\frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n, \bar{w}_1 \dots \bar{w}_n)} \leq C_{\#} \frac{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n, \bar{w}_1 \dots \bar{w}_n)} \leq C_{\#} \gamma_0^{-1}.$$

We thus get for all $n \geq n_{\star}$

$$|p_{n+1}(\bar{\omega}, \bar{w}) - p_n(\bar{\omega}, \bar{w})| \leq C_{\#} \nu^n,$$

and so for any $m \geq 0$,

$$(4.5) \quad |p_{n+m}(\bar{\omega}, \bar{w}) - p_n(\bar{\omega}, \bar{w})| \leq C_{\#} \sum_{k=0}^{m-1} \nu^{n+k} \leq C_{\#} \nu^n.$$

It follows that $(p_n(\bar{\omega}, \bar{w}))_n$ is a Cauchy sequence for \mathbb{P} -a.e. $\bar{\omega} \in \Omega$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$, and has thus a limit $J(\bar{\omega}, \bar{w})$. Taking the limit $m \rightarrow \infty$ in (4.5), we obtain (4.3) for all $n \geq n_{\star}$. From (4.4), it follows that $\|J\|_{\infty} < \infty$, which also allows to deduce (4.3) for all $n \geq 0$. The fact that $|J|_{\nu} < \infty$ is a direct consequence of (4.3).

The positivity of J follows then by Lemma 4.14. \square

Accordingly, $\log J$ is Hölder with respect to the usual metric on the shift. Hence it can be seen as a potential of a Gibbs measure. Of course, such a Gibbs measure is random, depending on $\bar{\omega}$, and non translation invariant, but it is a natural generalisation of the usual random walk in random environment situation in which one has a random Markov chain on $\mathcal{W}^{\mathbb{N}}$.

The transfer operator \mathcal{L}_{\star} has the following expression:

Lemma 4.17. *For any $f \in L^1(\mathbb{P}_{\star})$, we have*

$$(4.6) \quad \mathcal{L}_{\star} f(\bar{\omega}, \bar{w}) = \sum_{w \in \mathcal{W}} J(\tau_{-w} \bar{\omega}, w \bar{w}) f(\tau_{-w} \bar{\omega}, w \bar{w}).$$

Proof. We have to prove that, for all $f \in L^1(\mathbb{P}_{\star})$ and $g \in L^{\infty}(\mathbb{P}_{\star})$,

$$(4.7) \quad \int_{\Omega_{\star}} f g \circ \mathcal{F}_{\star} d\mathbb{P}_{\star} = \int_{\Omega_{\star}} \mathcal{L}_{\star} f g d\mathbb{P}_{\star},$$

where $\mathcal{L}_* f$ is given by (4.6). We first assume that both f and g are bounded, and depend only on $(\bar{\omega}, \bar{w}_0, \dots, \bar{w}_{k-1})$ for some $k \geq 1$. For any $n \geq k$, we have

$$\begin{aligned} \int_{\Omega_*} f g \circ \mathcal{F}_* d\mathbb{P}_* &= \int_{\Omega} \int_{\mathcal{W}^{\mathbb{N}}} f(\bar{\omega}, \bar{w}_0, \dots, \bar{w}_{k-1}) g(\tau_{\bar{w}_0} \bar{\omega}, \bar{w}_1, \dots, \bar{w}_k) \mathbb{P}_{\bar{\omega}}(d\bar{w}) \mathbb{P}(d\bar{\omega}) \\ &= \int_{\Omega} \sum_{\bar{w}_0, \dots, \bar{w}_n \in \mathcal{W}} p(\bar{\omega}, n+1, \bar{w}_0 \dots \bar{w}_n) f(\bar{\omega}, \bar{w}_0, \dots, \bar{w}_{k-1}) g(\tau_{\bar{w}_0} \bar{\omega}, \bar{w}_1, \dots, \bar{w}_k) \mathbb{P}(d\bar{\omega}) \\ &= \sum_{\bar{w}_0, \dots, \bar{w}_n \in \mathcal{W}} \int_{\Omega} p(\tau_{-\bar{w}_0} \bar{\omega}, n+1, \bar{w}_0 \dots \bar{w}_n) f(\tau_{-\bar{w}_0} \bar{\omega}, \bar{w}_0, \dots, \bar{w}_{k-1}) g(\bar{\omega}, \bar{w}_1, \dots, \bar{w}_k) \mathbb{P}(d\bar{\omega}) \\ &= \int_{\Omega} \left(\sum_{w \in \mathcal{W}} \frac{p(\tau_{-w} \bar{\omega}, n+1, w \bar{w}_0 \dots \bar{w}_{n-1})}{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})} f(\tau_{-w} \bar{\omega}, w, \bar{w}_0, \dots) \right) g(\bar{\omega}, \bar{w}) \mathbb{P}_*(d\bar{\omega}, d\bar{w}), \end{aligned}$$

where we have used the translation invariance of \mathbb{P} at the third line. Taking the limit as $n \rightarrow \infty$ and using Lemma 4.16, we obtain (4.7). The result for general f and g is obtained by approximation. \square

Define for each $k \geq 1$,

$$J_k(\bar{\omega}, \bar{w}) = \prod_{i=0}^{k-1} J(\mathcal{F}_*^i(\bar{\omega}, \bar{w})) = \lim_{n \rightarrow \infty} \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0 + \dots + \bar{w}_{k-1}} \bar{\omega}, n-k, \bar{w}_k \dots \bar{w}_{n-1})}.$$

It is immediate to verify that, for any $f \in L^1(\mathbb{P}_*)$,

$$\mathcal{L}_*^k f(\bar{\omega}, \bar{w}) = \sum_{w^k \in \mathcal{W}^k} J_k(\tau_{-w^k} \bar{\omega}, w^k \bar{w}) f(\tau_{-w^k} \bar{\omega}, w^k \bar{w})$$

where, for $w^k = (w_0, \dots, w_{k-1})$, $\tau_{-w^k} = \tau_{-(w_0 + \dots + w_{k-1})}$.

We introduce, for $w^k = (w_0, \dots, w_{k-1}) \in \mathcal{W}^k$, the map

$$\psi_{w^k}(\bar{\omega}, \bar{w}) = (\tau_{-w^k} \bar{\omega}, w^k \bar{w}),$$

so that

$$\mathcal{L}_*^k f = \sum_{w^k \in \mathcal{W}^k} J_k \circ \psi_{w^k} f \circ \psi_{w^k}.$$

Lemma 4.18. J_k belong to \mathcal{H}_ν for all $k \geq 1$.

Proof. Recall the notation $p_n(\bar{\omega}, \bar{w}) = \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0} \bar{\omega}, n-1, \bar{w}_1 \dots \bar{w}_{n-1})}$, and set

$$p_{n,k}(\bar{\omega}, \bar{w}) := \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0 + \dots + \bar{w}_{k-1}} \bar{\omega}, n-k, \bar{w}_k \dots \bar{w}_{n-1})} = \prod_{i=0}^{k-1} p_{n-i}(\mathcal{F}_*^i(\bar{\omega}, \bar{w})).$$

By Lemma 4.16, we have

$$|J(\mathcal{F}_*^i(\bar{\omega}, \bar{w})) - p_{n-i}(\mathcal{F}_*^i(\bar{\omega}, \bar{w}))| \leq C_{\#} \nu^{n-i},$$

and, consequently, using the inequality

$$\left| \prod_{i=0}^{k-1} a_i - \prod_{i=0}^{k-1} b_i \right| \leq \sum_{i=0}^{k-1} |a_i - b_i| \prod_{j \neq i} \max\{a_j, b_j\},$$

valid for all non-negative sequences $(a_i), (b_i)$, we obtain

$$\begin{aligned} |J_k(\bar{\omega}, \bar{w}) - p_{n,k}(\bar{\omega}, \bar{w})| &\leq C_{\#} \nu^n \sum_{i=0}^{k-1} \nu^{-i} \prod_{j \neq i} \max\{\|J\|_{\infty}, \|p_{n-i}\|_{\infty}\} \\ &= C_k \nu^n. \end{aligned}$$

where C_k depends only on k , since $\sup_n \|p_n\|_{\infty} < \infty$ by **(Abs)** and $\|J\|_{\infty} < \infty$ by Lemma 4.16. The lemma follows immediately. \square

Lemma 4.19. *There exists $C_{\#} > 0$ such that $C_{\#}^{-1} \leq \mathcal{L}_{\star}^k \mathbf{1}(\bar{\omega}, \bar{w}) \leq C_{\#}$ for all $k \geq 0$, \mathbb{P} -a.e. $\bar{\omega} \in \Omega$ and all $\bar{w} \in \mathcal{W}^{\mathbb{N}}$.*

Proof. This is a simple reformulation of **(Abs)**, as

$$\begin{aligned} \mathcal{L}_{\star}^k \mathbf{1}(\bar{\omega}, \bar{w}) &= \sum_{(w_0, \dots, w_{k-1}) \in \mathcal{W}^k} J_k(\tau_{-(w_0 + \dots + w_{k-1})} \bar{\omega}, w_0 \dots w_{k-1} \bar{w}) \\ &= \lim_{n \rightarrow \infty} \sum_{(w_0, \dots, w_{k-1}) \in \mathcal{W}^k} \frac{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})}{p(\tau_{\bar{w}_0 + \dots + \bar{w}_{k-1}} \bar{\omega}, n - k, \bar{w}_k \dots \bar{w}_{n-1})}. \end{aligned}$$

\square

Lemma 4.20. *There exist $C_{\#} > 0$ and $\xi \in (0, 1)$ such that for all $n \geq 0$ and all $f \in \mathcal{H}_{\nu}$,*

$$\begin{aligned} \|\mathcal{L}_{\star}^n f\|_{\infty} &\leq C_{\#} \|f\|_{\infty}, \\ \|\mathcal{L}_{\star}^n f\|_{\nu} &\leq C_{\#} \xi^n \|f\|_{\nu} + C_{\#} \|f\|_{\infty}. \end{aligned}$$

Proof. For $f \in \mathcal{H}_{\nu}$, we have

$$|\mathcal{L}_{\star}^n f| \leq \sum_{w^n \in \mathcal{W}^n} J_n \circ \psi_{w^n} |f| \circ \psi_{w^n} \leq \|f\|_{\infty} \mathcal{L}_{\star}^n \mathbf{1} \leq C_{\#} \|f\|_{\infty},$$

by Lemma 4.19. This proves that $\|\mathcal{L}_{\star}^n f\|_{\infty} \leq C \|f\|_{\infty}$. We also have, setting $\eta = \psi_{w^n}(\bar{\omega}, \bar{w})$ and $\eta' = \psi_{w^n}(\bar{\omega}, \bar{w}')$

$$\begin{aligned} |\mathcal{L}_{\star}^n f(\bar{\omega}, \bar{w}) - \mathcal{L}_{\star}^n f(\bar{\omega}, \bar{w}')| &\leq \sum_{w^n \in \mathcal{W}^n} J_n(\eta) |f(\eta) - f(\eta')| + |J_n(\eta) - J_n(\eta')| |f(\eta')| \\ &\leq \left(\sum_{w^n \in \mathcal{W}^n} J_n(\eta) |f|_{\nu} + \sum_{w^n \in \mathcal{W}^n} |J_n|_{\nu} \|f\|_{\infty} \right) d_{\nu}(w^n \bar{w}, w^n \bar{w}') \\ &\leq (\mathcal{L}_{\star}^n \mathbf{1}(\bar{\omega}, \bar{w}) |f|_{\nu} + (\#\mathcal{W})^n |J_n|_{\nu} \|f\|_{\infty}) \nu^n d_{\nu}(\bar{w}, \bar{w}'). \end{aligned}$$

By Lemma 4.19, this shows that, for all $n \geq 0$ and $f \in \mathcal{H}_{\nu}$,

$$\begin{aligned} \|\mathcal{L}_{\star}^n f\|_{\nu} &\leq C_{\#} \nu^n \|f\|_{\nu} + (1 + (\#\mathcal{W})^n \nu^n |J_n|_{\nu}) \|f\|_{\infty} \\ &\leq C_{\#} \nu^n \|f\|_{\nu} + C_n \|f\|_{\infty}. \end{aligned}$$

In particular, $\mathcal{L}_{\star} : \mathcal{H}_{\nu} \rightarrow \mathcal{H}_{\nu}$ is a continuous operator.

Take $k \geq 0$ such that the term $\tilde{\nu} := C_{\#} \nu^k$ in front of $\|f\|_{\nu}$ is strictly less than 1 and set $\xi = \tilde{\nu}^{\frac{1}{k}}$. Writing $n = qk + r$, with $0 \leq r < k$, we have, by iterating the previous inequality,

$$\begin{aligned} \|\mathcal{L}_{\star}^n f\|_{\nu} &\leq \tilde{\nu}^q \|\mathcal{L}_{\star}^r f\|_{\nu} + C_{\#} C_k (1 - \tilde{\nu}^{-1}) \|f\|_{\infty} \\ &\leq \tilde{\nu}^q \sup_{r < k} \|\mathcal{L}_{\star}^r\|_{\mathcal{H}_{\nu} \rightarrow \mathcal{H}_{\nu}} \|f\|_{\nu} + C_{\#} \|f\|_{\infty} \\ &\leq C_{\#} \xi^n \|f\|_{\nu} + C_{\#} \|f\|_{\infty}. \end{aligned}$$

□

Lemma 4.21. *There exists a continuous projection $\Pi : L^1(\mathbb{P}_\star) \rightarrow L^1(\mathbb{P}_\star)$ with $\Pi(L^1(\mathbb{P}_\star)) = \ker(\text{id} - \mathcal{L}_\star)$ such that*

$$\frac{1}{n} \sum_{k=0}^{n-1} \mathcal{L}_\star^k \rightarrow \Pi$$

in the strong operator topology.

Proof. For $h \in \mathcal{H}_\nu$, Lemma 4.20 implies that $\left\{ \frac{1}{n} \sum_{k=0}^{n-1} \mathcal{L}_\star^k h \right\}_{n \geq 1}$ is bounded in $L^\infty(\mathbb{P}_\star)$. By the Banach-Alaoglu theorem, since $L^\infty(\mathbb{P}_\star)$ is the dual of $L^1(\mathbb{P}_\star)$, the set $\left\{ \frac{1}{n} \sum_{k=0}^{n-1} \mathcal{L}_\star^k h \right\}_{n \geq 1}$ is weakly relatively compact in $L^1(\mathbb{P}_\star)$. This holds for all $h \in \mathcal{H}_\nu$, which is dense in $L^1(\mathbb{P}_\star)$ by Lemma 4.13, and so by the Kakutani-Yosida theorem [9, VIII.5.2, 5.3], the operators $\frac{1}{n} \sum_{k=0}^{n-1} \mathcal{L}_\star^k$ converge in the strong operator topology to the projection Π with range the set of fixed points of \mathcal{L}_\star in $L^1(\mathbb{P}_\star)$ and kernel the closure of $(\text{id} - \mathcal{L}_\star)(L^1(\mathbb{P}_\star))$. □

Define

$$h_\star = \Pi \mathbf{1} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathcal{L}_\star^k \mathbf{1},$$

in $L^1(\mathbb{P}_\star)$. By Lemma 4.21, we have $\mathcal{L}_\star h_\star = h_\star$. We clearly have $\int_{\Omega_\star} h_\star d\mathbb{P}_\star = 1$, and the fact that $C_\#^{-1} \leq h_\star \leq C_\#$, \mathbb{P}_\star -a.e., is an immediate consequence of Lemma 4.19. Consequently, the probability measure \mathbb{Q}_\star defined by

$$(4.8) \quad d\mathbb{Q}_\star = h_\star d\mathbb{P}_\star$$

is \mathcal{F}_\star -invariant and equivalent to \mathbb{P}_\star .

Next, we show that $\Pi(L^1(\mathbb{P}_\star))$ is the one-dimensional subspace generated by h_\star . Firstly, we prove a useful inclusion in \mathcal{H}_ν .

Lemma 4.22. *If $f \in L^\infty(\mathbb{P}_\star)$ and $\mathcal{L}_\star f = f$, then $f \in \mathcal{H}_\nu$.⁸*

Proof. Let $(\varphi_\epsilon)_\epsilon \subset \mathcal{H}_\nu$ be such that $\|f - \varphi_\epsilon\|_{L^1(\mathbb{P}_\star)} = \mathcal{O}(\epsilon)$ and $\|\varphi_\epsilon\|_\infty = \mathcal{O}(1)$, such a sequence exists by Lemma 4.13. We have

$$\begin{aligned} f &= \mathcal{L}_\star^n f = \mathcal{L}_\star^n \varphi_\epsilon + \mathcal{L}_\star^n (f - \varphi_\epsilon) \\ &=: \widehat{\varphi}_\epsilon^{(n)} + \gamma_\epsilon^{(n)}. \end{aligned}$$

This decomposition satisfies

$$\|\gamma_\epsilon^{(n)}\|_{L^1(\mathbb{P}_\star)} = \|\mathcal{L}_\star^n (f - \varphi_\epsilon)\|_{L^1(\mathbb{P}_\star)} \leq \|f - \varphi_\epsilon\|_{L^1(\mathbb{P}_\star)} = \mathcal{O}(\epsilon),$$

and

$$\|\widehat{\varphi}_\epsilon^{(n)}\|_\nu = \|\mathcal{L}_\star^n \varphi_\epsilon\|_\nu \leq C_\# \xi^n \|\varphi_\epsilon\|_\nu + C_\# \|\varphi_\epsilon\|_\infty = \mathcal{O}(\xi^n \|\varphi_\epsilon\|_\nu + 1),$$

using Lemma 4.20. If we choose n_ϵ such that $\xi^{n_\epsilon} \|\varphi_\epsilon\|_\nu = \mathcal{O}(1)$ and set $\widehat{\varphi}_\epsilon = \widehat{\varphi}_\epsilon^{(n_\epsilon)}$ and $\gamma_\epsilon = \gamma_\epsilon^{(n_\epsilon)}$, we then have $f = \widehat{\varphi}_\epsilon + \gamma_\epsilon$ with $\|\gamma_\epsilon\|_{L^1(\mathbb{P}_\star)} = \mathcal{O}(\epsilon)$ and $\|\widehat{\varphi}_\epsilon\|_\nu = \mathcal{O}(1)$.

For $\delta > 0$, we define

$$B_{\epsilon, \delta} = \{|f - \widehat{\varphi}_\epsilon| > \delta\} = \{|\gamma_\epsilon| > \delta\},$$

which satisfies $\mathbb{P}_\star(B_{\epsilon, \delta}) \leq \delta^{-1} \|\gamma_\epsilon\|_{L^1(\mathbb{P}_\star)}$ by Markov's inequality.

⁸ That is, there exists an element in the equivalence class of f that belongs to \mathcal{H}_ν .

For \mathbb{P} -a.e. $\bar{\omega} \in \Omega$ and all $\bar{w}, \bar{w}' \in \mathcal{W}^{\mathbb{N}}$ such that both $(\bar{\omega}, \bar{w})$ and $(\bar{\omega}, \bar{w}')$ do not belong to $B_{\epsilon, \delta}$, we have

$$\begin{aligned} |f(\bar{\omega}, \bar{w}) - f(\bar{\omega}, \bar{w}')| &\leq |\widehat{\varphi}_\epsilon(\bar{\omega}, \bar{w}) - \widehat{\varphi}_\epsilon(\bar{\omega}, \bar{w}')| + |\gamma_\epsilon(\bar{\omega}, \bar{w}) - \gamma_\epsilon(\bar{\omega}, \bar{w}')| \\ &\leq |\widehat{\varphi}_\epsilon|_\nu d_\nu(\bar{w}, \bar{w}') + |\gamma_\epsilon(\bar{\omega}, \bar{w})| + |\gamma_\epsilon(\bar{\omega}, \bar{w}')| \\ &\leq C d_\nu(\bar{w}, \bar{w}') + 2\delta. \end{aligned}$$

We set $B_\delta = \bigcap_{k \geq 0} \bigcup_{j \geq k} B_{2^{-j}, \delta}$, which satisfies $\mathbb{P}_*(B_\delta) = 0$, since

$$\mathbb{P}_* \left(\bigcup_{j \geq k} B_{2^{-j}, \delta} \right) = \mathcal{O} \left(\sum_{j \geq k} \|\gamma_{2^{-j}}\|_{L^1(\mathbb{P}_*)} \right) = \mathcal{O} \left(\sum_{j \geq k} 2^{-j} \right) = o(1).$$

Thus, $B = \bigcup_{n \in \mathbb{N}} B_{1/n}$ is also of zero measure and, eventually changing f on the zero measure set B , we have $f \in \mathcal{H}_\nu$. \square

We can now prove the main theorem:

Proof of Theorem 4.9. The probability measure \mathbb{Q}_* defined by (4.8) is \mathcal{F}_* -invariant and equivalent to \mathbb{P}_* . If $B \subset \Omega_*$ is a \mathcal{F}_* -invariant set, we have

$$\mathcal{L}_*(\mathbf{1}_B h_*) = \mathcal{L}_*((\mathbf{1}_B \circ \mathcal{F}_*) h_*) = \mathbf{1}_B \mathcal{L}_*(h_*) = \mathbf{1}_B h_*,$$

and so $\mathbf{1}_B h_*$ is a fixed point of \mathcal{L}_* in $L^\infty(\mathbb{P}_*)$. By Lemma 4.22, we have $\mathbf{1}_B h_* \in \mathcal{H}_\nu$, and so $\mathbf{1}_B = h_*^{-1}(h_* \mathbf{1}_B) \in \mathcal{H}_\nu$.⁹ This implies that there exists $N_B > 0$ such that $\mathbf{1}_B(\bar{\omega}, \bar{w}) = \mathbf{1}_B(\bar{\omega}, w_0, \dots, w_{N_B-1})$.

By the invariance of B it follows, for each $m \geq N_B$,

$$(4.9) \quad \begin{aligned} \mathbf{1}_B(\bar{\omega}, w_0, \dots, w_{N_B-1}) &= \mathbf{1}_B \circ \mathcal{F}_*^m(\bar{\omega}, w_0, \dots, w_{N_B-1}) \\ &= \mathbf{1}_B(\tau_{w_0 + \dots + w_{m-1}} \bar{\omega}, w_m, \dots, w_{N_B+m-1}). \end{aligned}$$

By **(Pro)** we can choose m and w_{N_B}, \dots, w_{m-1} such that $w_0 + \dots + w_{m-1} = 0$. It follows that $\mathbf{1}_B(\bar{\omega}, \bar{w}) = \mathbf{1}_B(\bar{\omega})$. Then (4.9) implies $\tau_{w_0 + \dots + w_{m-1}} B \subset B$ for all $(w_0, \dots, w_{m-1}) \in \mathcal{W}^m$. Accordingly, B is invariant for the group generated by \mathcal{W} and, by **(Pro)** again, it is either of zero or full measure due to ergodicity of \mathbb{P} , which concludes the proof. \square

Lemma 4.23. For all $f \in L^1(\mathbb{P}_*)$, we have

$$\Pi f = \left(\int_{\Omega_*} f d\mathbb{P}_* \right) h_*.$$

Proof. For each $\varphi \in L^\infty$ we have

$$\begin{aligned} \int_{\Omega_*} \varphi \Pi f d\Pi_* &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \int_{\Omega_*} \varphi \mathcal{L}_*^k f d\mathbb{P}_* = \int_{\Omega_*} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \varphi \circ \mathcal{F}_*^k \cdot f d\mathbb{P}_* \\ &= \left[\int_{\Omega_*} \varphi h_* d\mathbb{P}_* \right] \left[\int_{\Omega_*} f d\mathbb{P}_* \right] \end{aligned}$$

where, in the second equality, we have used Lebesgue dominated convergence Theorem and, in the second line, we have used the Birkhoff theorem and the ergodicity of \mathbb{Q}_* (and hence of $f\mathbb{P}_*$) established in Theorem 4.9. \square

⁹ Since h_* belongs to \mathcal{H}_ν by Lemma 4.22 and so does h_*^{-1} since $\inf h_* > 0$.

4.4. Application to deterministic walks in random environment. Deterministic walks in random environment, as presented in Section 3, naturally define random processes as described in the previous subsections. Indeed, if $\mathcal{A} = \{(f_\alpha, \mathcal{M}, \mathcal{G}_\alpha)\}_{\alpha \in A}$ is a deterministic walk in random environment, where all maps f_α are non-singular with respect to some reference measure m on \mathcal{M} , and the initial condition is given by an absolutely continuous probability measure $d\mu = h_0 dm$, then the probabilities $p(\bar{\omega}, n, w_0 \dots w_{n-1})$ are given by

$$(4.10) \quad p(\bar{\omega}, n, w_0 \dots w_{n-1}) = \int_{\mathcal{M}} \mathcal{L}_{\bar{\omega}, z_{n-1}, w_{n-1}} \dots \mathcal{L}_{\bar{\omega}, z_0, w_0} h_0 dm,$$

as we have seen in Section 3. Recall that $w_n = e(\bar{\omega}_{z_n}, x_n)$, where $(x_n, z_n) = \mathbb{F}_{\bar{\omega}}^n(x_0, z_0)$.

Remark 4.24. Note that we have a priori defined two different notions of environment as seen from the particle, in subsections 3.1 and 4.2, but the map $\Phi : \Omega \times \mathcal{M} \rightarrow \Omega \times \mathcal{W}^{\mathbb{N}}$ defined by $\Phi(\bar{\omega}, x) = (\bar{\omega}, \bar{w})$ with $\bar{w} = (w_n)_n$, is a semi-conjugacy between $(\Omega \times \mathcal{M}, \mathcal{F})$ and $(\Omega \times \mathcal{W}^{\mathbb{N}}, \mathcal{F}_\star)$, and if the maps f_α are expansive, it is invertible a.e.

If we are able to check the assumptions **(Pos)**, **(Exp)**, **(Abs)**, **(Ell)** and **(Pro)**, then Theorem 4.9 applies, and we deduce the existence of a deterministic drift V .

A particular situation, which we have already encountered in Lemma 3.1, occurs when all maps f_α preserve the same invariant measure $d\lambda = h_0 dm$, and the set \mathcal{G}_α is deterministic, i.e. $G_{\alpha, w} = G_w$ does not depend on $\alpha \in A$. In this case, the measure \mathbb{P}_\star on $\Omega \times \mathcal{W}^{\mathbb{N}}$ is invariant under \mathcal{F}_\star , since it is the push-forward of $\mathbb{P}_0 = \mathbb{P} \times \lambda$, which is \mathcal{F} -invariant and the condition **(Abs)** is automatically satisfied by Lemma 4.5.

Remark 4.25. Lemma 3.2 and Remark 4.10 agree, since

$$\begin{aligned} \sum_{w \in \mathcal{W}} w \int_{\Omega} p(\bar{\omega}, 1, w) \mathbb{P}(d\bar{\omega}) &= \sum_{w \in \mathcal{W}} w \int_{\Omega} \int_{\mathcal{M}} \mathcal{L}_{\bar{\omega}, 0, w} h_0 dm \mathbb{P}(d\bar{\omega}) \\ &= \sum_{w \in \mathcal{W}} w \int_{\Omega} \int_{\mathcal{M}} \mathcal{L}_{f_{\bar{\omega} w}} \mathbf{1}_{G_w} h_0 dm \mathbb{P}(d\bar{\omega}) \\ &= \sum_{w \in \mathcal{W}} w \int_{G_w} h_0 dm \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}_0(e \circ \pi) &= \int_{\Omega} \int_{\mathcal{M}} e(\pi(\bar{\omega}, x)) h_0(x) m(dx) \mathbb{P}(d\bar{\omega}) \\ &= \int_{\Omega} \int_{\mathcal{M}} e(\bar{\omega}_0, x) h_0(x) m(dx) \mathbb{P}(d\bar{\omega}) \\ &= \sum_{w \in \mathcal{W}} w \int_{G_w} h_0 dm. \end{aligned}$$

5. EXAMPLES

In this section we discuss several concrete example of the previous abstract models.

5.1. Lorentz gas as a deterministic walk in random environment. It is easy to verify that the Random Lorentz gas presented in Section 2 is a special example of a deterministic walk in random environment. However the Random Lorentz gas has several special features:

- (1) the maps f_α have all the same invariant measure λ , the Lebesgue measure;
- (2) the set \mathcal{G}_α is non random, i.e. it does not depend from α ;
- (3) the associated random process is reversible, in particular $\lambda(e) = 0$.

The first two facts are obvious. Let us discuss the third. The dynamics is reversible under the involution $(q, p) \rightarrow (q, -p)$, which, at the level of the Poincaré map, writing $x \in \cup_{w \in \mathcal{W}} G_w$ as $x = (w, s, \theta)$,¹⁰ reads $i(w, s, \theta, z) = (-w, s, \theta, z + w)$.¹¹ Also, let $\pi(w, s, \theta, z) = (w, s, \theta)$ and $i_1(w, s, \theta) = (-w, s, \theta)$ so that $i_1 \circ \pi = \pi \circ i$. Thus, choosing as initial measure the common invariant measure of the Poincaré maps h_0 (hence $h_0 \circ i_1 = h_0$), we have

$$\begin{aligned} \mathbb{P}_\star(\{z(1), \dots, z(n)\} \mid \bar{\omega}) &= \int_{\mathcal{B}} \prod_{k=0}^{n-1} \mathbf{1}_{G_{w^{(k)}}} \circ \pi \circ \mathbb{F}_{\bar{\omega}}^k(x, 0) h_0(x) dx \\ &= \int_{\mathcal{B}} \prod_{k=0}^{n-1} \mathbf{1}_{G_{w^{(k)}}} \circ \pi \circ \mathbb{F}_{\bar{\omega}}^k \circ i(i_1(x), e(x)) h_0(x) dx \\ &= \int_{\mathcal{B}} \prod_{k=0}^{n-1} \mathbf{1}_{G_{-w^{(k)}}} \circ \pi \circ \mathbb{F}_{\bar{\omega}}^{-k}(x, -e(x)) h_0(x) dx \\ &= \int_{\mathcal{B}} \prod_{k=0}^{n-1} \mathbf{1}_{G_{-w^{(k)}}} (f_{\bar{\omega}_{z^{(k)}}}^{-1} \circ \dots \circ f_{\bar{\omega}_{z^{(1)}}}^{-1}(x)) h_0(x) dx, \end{aligned}$$

where, in the third line, we have used the invariance of the measure with respect to i_1 and the relation $e(i_1(x)) = -e(x)$; while, in the last line, we have used the formula $\mathbb{F}_{\bar{\omega}}(x, z) = (f_{\bar{\omega}_z}^{-1}(x), z - e(f_{\bar{\omega}_z}^{-1}(x)))$. Next, using the invariance of the measure with respect to the maps $f_{\bar{\omega}_z}$,

$$\mathbb{P}_\star(\{z(1), \dots, z(n)\} \mid \bar{\omega}) = \int_{\mathcal{B}} \prod_{k=0}^{n-1} \mathbf{1}_{G_{-w^{(k)}}} (f_{\bar{\omega}_{z^{(k+1)}}} \circ \dots \circ f_{\bar{\omega}_{z^{(n-1)}}}(x)) h_0(x) dx.$$

Then, setting $\tilde{w}(k) = -w(n-1-k)$, $\tilde{z}(k) = \sum_{j=0}^{k-1} \tilde{w}(j)$ and $\tilde{\omega} = \tau_{z(n)} \bar{\omega}$,

$$\begin{aligned} (5.1) \quad \mathbb{P}_\star(\{z(1), \dots, z(n)\} \mid \bar{\omega}) &= \int_{\mathcal{B}} \prod_{k=0}^{n-1} \mathbf{1}_{G_{\tilde{w}^{(k)}}} (f_{\tilde{\omega}_{\tilde{z}^{(k)}}} \circ \dots \circ f_{\tilde{\omega}_{\tilde{z}^{(1)}}}(x)) h_0(x) dx \\ &= \mathbb{P}_\star(\{\tilde{z}(n-1), \dots, \tilde{z}(0)\} \mid \tau_{z(n)} \bar{\omega}). \end{aligned}$$

Note that $\tilde{z}(k) = z(n-k) - z(n)$. Which is the reversibility of the random process. In particular Lemma 4.6 applies.

The Lorentz gas has a dynamics that is hard to study. To make further progresses let us consider simpler local dynamics.

¹⁰ The curvilinear coordinate s is chosen such that the segments $s \rightarrow (w, s)$ and $s \rightarrow (-w, s)$, $w \neq 0$ go in the same direction.

¹¹ Recall that we are considering the case of deterministic gates, although the following consideration easily extend to the general case. In particular, e is a function of x only, and we have $e(w, s, \theta) = w$.

5.2. Markovian models. To try to get a better feeling for the difficulties involved in studying the above questions, let us try to invent a model stripped of all the technical difficulties present in the Lorentz gas dynamics. For simplicity let us discuss the case $d = 1$, although similar considerations hold in any higher dimensional lattice. To simplify the dynamics f in (2.3) let us suppose that it is a map from $[0, 1]$ to itself. Hence the map $\mathbb{F}_{\bar{\omega}}$ acts on $[0, 1] \times \mathbb{Z}$. Also, we assume that the environment is a random variable distributed according to a Bernoulli product measure over the space $\Omega = A^{\mathbb{Z}} = \{-1, 1\}^{\mathbb{Z}}$.

Example 1. The dynamics is defined by the map $f_{\alpha}(x) = 4x \bmod 1$ for $\alpha \in A$, with $G_{-1,-1} = [0, 1/4]$, $G_{-1,+1} = [1/4, 1]$ and $G_{+1,-1} = [0, 3/4]$, $G_{+1,+1} = [3/4, 1]$.

Remark 5.1. Here we are considering a more general situation than the one described for the Lorentz gas insofar also the gates are random. This is indeed the general case also for the Lorentz gas. We considered the case of deterministic gates only to simplify the exposition.

Also, we consider the initial distribution $h_0 = 1$. Then an elementary computation shows that

$$\mathbb{P}_{\star}(z(n+1) - z(n) = \pm 1 \mid \bar{\omega}, z(n), \dots, z(0)) = |G_{\bar{\omega}_{z(n)}, \pm 1}| = \frac{1}{2} \mp \frac{\bar{\omega}_{z(n)}}{4}.$$

This is an example of Sinai's walk, hence we do not have the classical CLT.

Example 2. Assume that $G_{\alpha,-1} = G_{-1} = [0, 1/2]$ and $G_{\alpha,+1} = G_{+1} = (1/2, 1]$ for any α and the maps are defined by

$$f_{-1}(x) = \begin{cases} 2x & x \in [0, 1/4] \\ 4x \bmod 1 & x > 1/4 \end{cases}$$

$$f_{+1}(x) = \begin{cases} 4x \bmod 1 & x \in [0, 3/4] \\ 2x - 1 & x > 3/4. \end{cases}$$

Again let us consider the initial distribution $h_0 = 1$. Denote by $\mathcal{L}_{\alpha,w}$ the operator $\mathcal{L}_{\alpha,w}(\phi) = \mathcal{L}_{f_{\alpha}}(\mathbf{1}_{G_w}\phi)$. The two dimensional vector space $\mathbb{V} = \{a_{-1}\mathbf{1}_{G_{-1}} + a_{+1}\mathbf{1}_{G_{+1}} : a_{-1}, a_{+1} \in \mathbb{R}\}$ is left invariant by the operators $\{\mathcal{L}_{\alpha,w}\}_{\alpha,w}$. Since $h_0 \in \mathbb{V}$, this allows to compute the transition probabilities by using formula (3.2).

If $\phi = a_{-1}\mathbf{1}_{G_{-1}} + a_{+1}\mathbf{1}_{G_{+1}}$, a direct computation shows that $\mathcal{L}_{\alpha,w}(\phi) = a_w\mathcal{L}_{\alpha,w}(\mathbf{1})$, and thus $\mathcal{L}_{\alpha',w'}\mathcal{L}_{\alpha,w}(\phi) = a_w\mathcal{L}_{\alpha',w'}\mathcal{L}_{\alpha,w}\mathbf{1}$. For any $\bar{\omega}$ and $z(1), \dots, z(n), z(n+1)$, denote by $\alpha_k = \bar{\omega}_{z(k)+w(k)}$ and $w_k = w(k) = z(k+1) - z(k)$. We have

$$\mathbb{P}_{\star}(z(1), \dots, z(n) \mid \bar{\omega}) = \int \mathcal{L}_{\alpha_{n-1}, w_{n-1}} \dots \mathcal{L}_{\alpha_0, w_0} \mathbf{1}.$$

Set $\phi = \mathcal{L}_{\alpha_{n-2}, w_{n-2}} \dots \mathcal{L}_{\alpha_0, w_0} \mathbf{1} = a_{-1}\mathbf{1}_{G_{-1}} + a_{+1}\mathbf{1}_{G_{+1}} \in \mathbb{V}$. We have

$$\mathbb{P}_{\star}(z(1), \dots, z(n) \mid \omega) = \int \mathcal{L}_{\alpha_{n-1}, w_{n-1}} \phi = a_{w_{n-1}} \int \mathcal{L}_{\alpha_{n-1}, w_{n-1}} \mathbf{1}$$

and

$$\mathbb{P}_{\star}(z(1), \dots, z(n), z(n+1) \mid \bar{\omega}) = a_{w_{n-1}} \int \mathcal{L}_{\alpha_n, w_n} \mathcal{L}_{\alpha_{n-1}, w_{n-1}} \mathbf{1}.$$

It follows that

$$\mathbb{P}_{\star}(z(n+1) \mid z(1), \dots, z(n), \bar{\omega}) = \frac{\int \mathcal{L}_{\alpha_n, w_n} \mathcal{L}_{\alpha_{n-1}, w_{n-1}} \mathbf{1}}{\int \mathcal{L}_{\alpha_{n-1}, w_{n-1}} \mathbf{1}}$$

which is a function of $z(n-1)$, $z(n)$ and $z(n+1)$ only. We have obtained a *persistent random walk*, that is a walk where the transition probability depends not only on the current position of the particle but also on its previous position.

Initial conditions. A natural question that arises at this point is what happens if one starts by a different initial measure. A moment thought shows that this is a non trivial issue. For instance, in the first example, there exists a Cantor set C (of zero Lebesgue measure) that corresponds to the coordinates $x(n)$ never belonging to $(1/4, 3/4)$. For such points $x \in C$, the set $\{x(n) \in G_{\alpha,w}\}$ does not depend on α , and so the process $(z(n))$ is completely unaffected by the environment. If we identify naturally the Cantor set C with $\{-1, +1\}^{\mathbb{N}}$ (in such a way that $x \in C$ is identified with the sequence (i_n) such that $x(n) \in I_{i_n}$ for all $n \geq 0$, where $I_{-1} = [0, 1/4]$ and $I_{+1} = [3/4, 1]$), then the initial distribution of x can be identified with a probability measure on $\{-1, +1\}^{\mathbb{N}}$ and this measure will be the distribution law of the random process $(w(n))$. In particular, if we consider the Bernoulli measure with equal probabilities on such a Cantor set as the initial distribution of x , then we obtain a standard random walk which has a very different behaviour than the Sinai's walk.

Without going to such extremes, one can (perhaps more naturally) start from a measure absolutely continuous with respect to Lebesgue and wonder which kind of process this will yield. We do not discuss this issue at present because it is part of the more general discussion that we will start in the next section.

Remark 5.2. *The above examples (among other obvious limitations) are unreasonable in one key aspect: their Markov structure. It is inevitable to ask what happens when the Markov structure is absent (as for billiards). The next section is devoted to investigating such a situation.*

6. NON-MARKOVIAN EXAMPLES: GENERAL DISCUSSION

We now consider a model of d -dimensional deterministic random walk in random environment $\mathcal{A} = \{(f_\alpha, \mathcal{M}, \mathcal{G}_\alpha)\}_{\alpha \in A}$ for a finite set A , where $\mathcal{M} = [0, 1]$, all maps $f_\alpha : [0, 1] \rightarrow [0, 1]$ are piecewise C^2 and uniformly expanding (i.e. $|f'_\alpha| \geq \lambda > 1$), and the partitions $\mathcal{G}_\alpha = \{G_{\alpha,w}\}_{w \in \mathcal{W}}$ are made of subintervals of $[0, 1]$, for a given bounded subset $\mathcal{W} \subset \mathbb{Z}^d$.

Let \mathbb{P} be a translation invariant probability on the set $\Omega = A^{\mathbb{Z}^d}$. For a given environment $\bar{\omega} \in \Omega$, we have the dynamics $\mathbb{F}_{\bar{\omega}}(\cdot, \cdot) : \mathcal{M} \times \mathbb{Z}^d \rightarrow \mathcal{M} \times \mathbb{Z}^d$ given by $\mathbb{F}_{\bar{\omega}}(x, z) = (f_{\bar{\omega}_{z+e(\bar{\omega}_z, x)}}(x), z + e(\bar{\omega}_z, x))$, where $e(\alpha, x) = \sum_{w \in \mathcal{W}} \mathbf{1}_{G_{\alpha,w}}(x)w$.

We are interested in the quenched evolution, $(x_n, z_n) = \mathbb{F}_{\bar{\omega}}^n(x_0, z_0)$, of such a system when the initial condition x_0 is distributed according to the measure

$$\mu(\varphi) = \int_0^1 \varphi(x, 0) h_0(x) dx$$

for some $h_0 \in \text{BV}$, with $\inf h_0 > 0$.

Recall the definition of the probability measure \mathbb{P}_\star and the dynamical system $(\Omega_\star, \mathcal{F}_\star)$ of the point of view of the particle, from Section 4.

Our goal is to reduce the study of this model to a probabilistic one. To this end we need some technical conditions.

6.1. Conditions (C1), (C2), (C3).

Let $\mathcal{T} = \{f_\alpha\}_{\alpha \in A}$ be a finite set of maps on $[0, 1]$, and \mathcal{H} be the (finite) set of all the possible intervals of the partitions, i.e. $\mathcal{H} = \{G_{\alpha, w}\}_{w \in \mathcal{W}, \alpha \in A}$.

Notation. The set $\mathcal{T} \times \mathcal{H}$ is canonically isomorphic to $\Sigma = A \times (A \times \mathcal{W})$, with correspondence $\rho : \Sigma \rightarrow \mathcal{T} \times \mathcal{H}$ given by $\rho((\alpha, \beta)) = (f_\alpha, G_\beta)$: $(T_\sigma, H_\sigma) = (f_{\pi_1 \circ \rho^{-1}(\sigma)}, G_{\pi_2 \circ \rho^{-1}(\sigma)})$, $\pi_1(\alpha, \beta) = \alpha$ and $\pi_2(\alpha, \beta) = \beta$. Hence we will often write $\Sigma = \mathcal{T} \times \mathcal{H}$.

Let $\tau : \Sigma^{\mathbb{N}} \rightarrow \Sigma^{\mathbb{N}}$ be the unilateral shift. We denote $T_\sigma^n = T_{\sigma_n} \circ \dots \circ T_{\sigma_1}$ and $H_\sigma^n = \bigcap_{j=0}^{n-1} (T_\sigma^j)^{-1}(H_{\sigma_{j+1}})$.

Let \mathcal{L}_{σ_k} be the transfer operator of the map T_{σ_k} with respect to the Lebesgue measure, i.e.

$$\mathcal{L}_{\sigma_k} f(x) = \sum_{T_{\sigma_k} y = x} \frac{f(y)}{|T'_{\sigma_k}(y)|}.$$

We set $\widehat{\mathcal{L}}_{\sigma_k} f = \mathcal{L}_{\sigma_k}(f \mathbf{1}_{H_{\sigma_k}})$ and $\widehat{\mathcal{L}}_\sigma^n = \widehat{\mathcal{L}}_{\sigma_n} \circ \dots \circ \widehat{\mathcal{L}}_{\sigma_1}$. We can write $\widehat{\mathcal{L}}_\sigma^n f(x) = \sum_{T_\sigma^n y = x} g_\sigma^n(y) f(y)$, where $g_\sigma^n = g_{\sigma_1} \times \dots \times g_{\sigma_n} \circ T_\sigma^{n-1}$, with $g_{\sigma_k} = \mathbf{1}_{H_{\sigma_k}} \frac{1}{|T'_{\sigma_k}|}$. Let $\Theta^{-1} = \inf_{T \in \mathcal{T}} \inf_x |T'(x)|$, then $\|g_\sigma^n\|_\infty \leq \Theta^n$.

We will only consider systems that satisfy

(C1): There exists $\delta_\star > 0$ such that for all $\sigma_1 \in \mathcal{T} \times \mathcal{H}$, $\inf \widehat{\mathcal{L}}_{\sigma_1} \mathbf{1} \geq \delta_\star$.

Observe that this condition is satisfied if, for any choice of T and H , T admits at least one full branch inside H . By iteration, we also have $\inf \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \geq \delta_\star^n$ for any $\sigma \in \Sigma^{\mathbb{N}}$ and $n \geq 1$. Next, we define the functionals

$$\Lambda_\sigma(f) = \liminf_{n \rightarrow \infty} \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}}.$$

This limit exists since the sequence is increasing and bounded. Indeed,

$$\begin{aligned} \inf \frac{\widehat{\mathcal{L}}_\sigma^{n+1} f}{\widehat{\mathcal{L}}_\sigma^{n+1} \mathbf{1}} &\geq \inf \frac{\widehat{\mathcal{L}}_{\sigma_{n+1}}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1} \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}})}{\widehat{\mathcal{L}}_\sigma^{n+1} \mathbf{1}} \\ &\geq \inf \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} \inf \frac{\widehat{\mathcal{L}}_{\sigma_{n+1}}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}{\widehat{\mathcal{L}}_\sigma^{n+1} \mathbf{1}} = \inf \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}}; \end{aligned}$$

and $-\|f\|_\infty \leq \inf \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} \leq \|f\|_\infty$. In particular, for all $n \geq 0$, we have

$$(6.1) \quad \Lambda_\sigma(f) \geq \inf \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}}.$$

The Λ_σ satisfy the following properties:

- $\Lambda_\sigma(\mathbf{1}) = 1$;
- $|\Lambda_\sigma(f)| \leq \|f\|_\infty$;
- $f \geq g$ implies $\Lambda_\sigma(f) \geq \Lambda_\sigma(g)$ (monotonicity);
- $\Lambda_\sigma(\lambda f) = \lambda \Lambda_\sigma(f)$, for $\lambda > 0$ (positive homogeneity);
- $\Lambda_\sigma(f + g) \geq \Lambda_\sigma(f) + \Lambda_\sigma(g)$ (super-additivity);
- $\Lambda_\sigma(f + b) = \Lambda_\sigma(f) + b$ for all $b \in \mathbb{R}$.

All the above follows immediately from the definition. Note that it is not clear at the moment if Λ_σ is linear or not.

Next, we define

$$(6.2) \quad \rho_\sigma = \Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_{\sigma_1} \mathbf{1}); \quad \rho = \inf_{\sigma \in \Sigma^{\mathbb{N}}} \rho_\sigma.$$

Let \mathcal{Z}_σ^n be the partition of smoothness intervals of T_σ^n , and $\widehat{\mathcal{Z}}_\sigma^n$ be the coarsest partition which is finer than \mathcal{Z}_σ^n and enjoying the property that the elements of the partition are either disjoint from $\overline{H_\sigma^n}$ or contained in H_σ^n .

Let us define the collections of intervals

$$(6.3) \quad \begin{aligned} \mathcal{Z}_{\sigma,\star}^n &= \{Z \in \widehat{\mathcal{Z}}_\sigma^n \mid Z \subset H_\sigma^n\}, \\ \mathcal{Z}_{\sigma,b}^n &= \{Z \in \mathcal{Z}_{\sigma,\star}^n \mid \Lambda_\sigma(\mathbf{1}_Z) = 0\} \\ \mathcal{Z}_{\sigma,g}^n &= \{Z \in \mathcal{Z}_{\sigma,\star}^n \mid \Lambda_\sigma(\mathbf{1}_Z) > 0\}. \end{aligned}$$

Definition 1. We will call contiguous two elements of $\mathcal{Z}_{\sigma,\star}^n$ that are either contiguous in the usual sense, or separated by a connected component of $(H_\sigma^n)^c = \bigcup_{j=0}^{n-1} (T_\sigma^j)^{-1}(H_{\sigma_{j+1}}^c)$.

We can now introduce the second condition needed to state our results.

(C2): there exist constants $K \geq 0$ and $\xi \geq 1$ such that for any n and $\sigma \in \Sigma^{\mathbb{N}}$, at most $K\xi^n$ elements of $\mathcal{Z}_{\sigma,b}^n$ are contiguous. In addition, $\theta := \xi\Theta < \min\{\rho, 1\}$. In particular, $\rho > 0$.

Let us turn to the third and last condition.

$$(C3(N, N')): \hat{\epsilon}(N, N') := \min_{k \leq N} \inf_{\sigma \in \Sigma^{\mathbb{N}}} \min_{Z \in \mathcal{Z}_{\sigma,g}^k} \inf_{x \in [0,1]} \frac{(\widehat{\mathcal{L}}_\sigma^{N'} \mathbf{1}_Z)(x)}{(\widehat{\mathcal{L}}_\sigma^{N'} \mathbf{1})(x)} > 0.$$

Note that, by (6.1), condition $C3(N, N')$ implies $\Lambda_\sigma(\mathbf{1}_Z) \geq \epsilon(N, N') > 0$ for all $Z \in \mathcal{Z}_{\sigma,g}^n$, $n \leq N$, and $\sigma \in \Sigma^{\mathbb{N}}$. This is morally what we need in the following, however in Lemma 9.3 the above more precise condition is used.

We remark also that $m' \geq m$ implies $\hat{\epsilon}(n, m') \geq \hat{\epsilon}(n, m)$ and $n \geq n'$ implies $\hat{\epsilon}(n, m) \leq \hat{\epsilon}(n', m)$. Thus, setting

$$\epsilon_\star(n) = \min\{\hat{\epsilon}(n, m) : m \geq n, \hat{\epsilon}(n, m) > 0\},$$

we have $\epsilon_\star(N) > 0$ if and only if $(C3(N, N'))$ holds for some $N' \geq N$. Also if $\epsilon_\star(n) > 0$, then $\epsilon_\star(n') > 0$ for all $n' \leq n$, however ϵ_\star is not necessarily a decreasing function.

6.2. The results.

We are finally able to state our main result.

Theorem 6.1. *There exists an integers $n_2 \geq 1$, depending only on the classes \mathcal{T} and \mathcal{H} , explicitly computable (see Remark 6.2), such that if (C1), (C2) and $(C3(n_2, n_3))$ hold for some $n_3 \geq n_2$, then the condition **(Exp)** holds. In particular, the property of loss memory from Lemma 4.3 is verified.*

Remark 6.2. *The definition of n_2 is a bit cumbersome but explicit: given $(\mathcal{T}, \mathcal{H})$ define the constants (see Lemma 9.3 for their meaning)¹²*

$$C_\star = 3(C+1) + 2K(3C+2); \quad C_n = \frac{(3C+2)(2K\xi+1)\Theta}{\epsilon_\star(n)}$$

where C is such that $\bigvee_Z g_\sigma^n \leq C \|g_\sigma^n\|_\infty$ for all n, σ and $Z \in \mathcal{Z}_{\sigma,\star}^n$. Recall the definitions of ρ and θ from (C2) and define (as in Lemma 9.5) $n_0 = \lceil \frac{\ln 4C_\star^2}{\ln \rho \theta^{-1}} \rceil$. If

¹² Note that $C_n = \infty$ if $\epsilon_\star(n) = 0$.

$\epsilon_*(n_0) = 0$, then define $n_2 = n_0$.¹³ If $\epsilon_*(n_0) > 0$, then $C_n < \infty$ for all $n \leq n_0$, and we can set (as in Remark 9.9) $a = \max\{1, \frac{15}{11} \max_{i \leq n_0} \frac{C_i}{C_* \theta^i}\}$ and $B = 1 + 2aC_*$ (used in Lemma 9.6). Finally we define (as introduced in Lemmata 9.7, 9.8)

$$n_2 = \left\lceil \frac{\ln 4aB(1 + 2C_{n_0}\rho^{-n_0})}{\ln \theta^{-1}\rho} \right\rceil.$$

According to Remark 6.2, to check the hypothesis of Theorem 6.1 one has first to check (C1), (C2); compute n_0 and find N' for which $(C3(n_0, N'))$ holds; use it to compute n_2 and look for an n_3 for which $(C3(n_2, n_3))$ holds. Given $(\mathcal{T}, \mathcal{H})$, this can be rather laborious as we will see in Section 7.

As we already pointed out in Section 5, the choice of the initial condition might play an important role. The following result states that if we restrict ourselves to initial conditions absolutely continuous to Lebesgue, with density in BV bounded uniformly away from 0, this difference is not so important in the sense that for large times, the transition probabilities are exponentially close. For two different initial densities $h_0, h'_0 \in \text{BV}$ with $\inf h_0 > 0$ and $\inf h'_0 > 0$, we denote by \mathbb{P}_* and \mathbb{P}'_* the probability measures corresponding to h_0 and h'_0 respectively.

Theorem 6.3. *Under the assumptions of Theorem 6.1, we have for all realisation of the environment $\bar{\omega} \in \Omega$, $n \geq 0$ and all densities h_0, h'_0 as above:*

$$|\mathbb{P}_*(z(n) \mid z(1), \dots, z(n-1), \bar{\omega}) - \mathbb{P}'_*(z(n) \mid z(1), \dots, z(n-1), \bar{\omega})| \leq C_{h_0, h'_0} \nu^n,$$

where $C_{h_0, h'_0} > 0$ depends only on the densities h_0 and h'_0 .

Next, we consider the situation where all maps f_α preserve a common density $h_0 \in \text{BV}$ such that $\inf h_0 > 0$, and when the partitions \mathcal{G}_α are deterministic, i.e. $G_{\alpha, w} = G_w$ does not depend on $\alpha \in A$. In this situation, the dynamical system $(\Omega_*, \mathcal{F}_*, \mathbb{P}_*)$ is measure-preserving, and so condition **(Abs)** holds by Lemma 4.5.

Theorem 6.4. *Under the assumptions of Theorem 6.1, if the maps f_α preserve a common density, the partitions \mathcal{G}_α are deterministic, and if furthermore condition **(Pro)** of Section 4.2 holds, then the dynamical system $(\Omega_*, \mathcal{F}_*, \mathbb{P}_*)$ is ergodic. In particular, \mathbb{P}_* -a.e.,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} z(n) = \sum_{w \in \mathcal{W}} w \int_{G_w} h_0 dm.$$

Remark 6.5. *The assumption that the maps all preserve a common measure and that the partitions are deterministic is only used to check the validity of condition **(Abs)** thanks to Lemma 4.5, and to have an explicit formula for the drift. If for a concrete example, one is able to check **(Abs)** by any other mean, then Theorem 4.9 applies and there exists $V \in \mathbb{R}^d$ such that $\lim_{n \rightarrow \infty} \frac{1}{n} z(n) = V$, \mathbb{P}_* -a.s. .*

The proofs of Theorems 6.1, 6.3 and 6.4 will be provided in Section 8.

7. EXISTENCE OF NON-MARKOVIAN EXAMPLES: β -MAPS

To ensure that the hypotheses of Theorem 6.1 are non empty it suffices to verify them in some limiting regime, e.g. when the dynamics has a lot of expansion. This is the aim of the present section. To further simplify things we will limit ourselves to β maps, a popular class of dynamical systems.

¹³ In this case $(C3(n_2, n_3))$ fails for all $n_3 \geq n_2$ and hence Theorem 6.1 does not apply.

7.1. General β -maps. More precisely, we consider the situation where the class of maps is $\mathcal{T} = \{T_{\beta_1}, T_{\beta_2}\}$ for $\beta_2 > \beta_1 > 1$, with $T_\beta(x) = \beta x \bmod 1$ and the partitions \mathcal{G}_α are such that $\mathcal{H} = \{[0, \frac{1}{2}], (\frac{1}{2}, 1]\}$. Note that \mathcal{W} is not specified, and \mathcal{G}_α can be random.¹⁴ For $\sigma \in \Sigma^{\mathbb{N}} = (\mathcal{T} \times \mathcal{H})^{\mathbb{N}}$, we denote by β_{σ_i} the value of β such that $T_{\sigma_i} = T_{\beta_{\sigma_i}}$. For simplicity, we will fix $\varpi > 1$ and assume that $\beta_1 = \beta$ and $\beta_2 = \varpi\beta$. We will show that assumptions (C1), (C2) and (C3) are verified for a large set of β . When needed, we will denote by $\mathcal{Z}_\sigma^n(\beta)$ the partition of smoothness intervals of T_σ^n , to emphasise the dependence on β , and similarly for the objects defined in equation (6.3). We will do the same with the subsets of the partition defined in Section 6.

Proposition 7.1. *There exists a set $\mathcal{B} \subset (1, \infty)$ and $C > 0$, with $\text{Leb}(\mathcal{B} \cap (1, t)) \leq C \log t$, for $t > 1$, such that the model described above satisfies the assumptions of Theorem 6.1 when $\beta \notin \mathcal{B}$.*

The rest of the section is devoted to the proof of Proposition 7.1. It suffices to check conditions (C1), (C2) and (C3) as explained in Remark 6.2.

7.1.1. Condition (C1). This condition is satisfied if every map $T \in \mathcal{T}$ admits at least one full branch inside any interval $H \in \mathcal{H}$. This is the case whenever $\beta \geq 3$.

7.1.2. Condition (C2). We first give a general criterion to check this condition. Let $\mathcal{Z}_{\sigma, f}^n$ be the collection of elements Z in $\mathcal{Z}_{\sigma, \star}^n$ such that $T_\sigma^n Z = [0, 1]$, and let $\mathcal{Z}_{\sigma, u}^n = \mathcal{Z}_{\sigma, \star}^n \setminus \mathcal{Z}_{\sigma, f}^n$.

We call a system ξ -full branched, $\xi > 0$, if there exists $K > 0$ such that for all $\sigma \in \Sigma$ and n , the number of contiguous elements in $\mathcal{Z}_{\sigma, u}^n$ is less than $K\xi^n$. A system ξ -full branched satisfies (C2) with the same K, ξ .

Lemma 7.2. Calling C_σ^n the maximal number of contiguous elements in $\mathcal{Z}_{\sigma, u}^n$, holds

$$C_\sigma^n \leq 2 \sum_{i=0}^{n-1} (C^{(1)} + 2)^i C^{(1)},$$

where $C^{(1)}$ is the supremum over all σ of C_σ^1 .

Proof. The proof is by induction on n . Clearly it is true for $n = 1$. Let us suppose it true for n . The elements of the partition $\mathcal{Z}_{\sigma, \star}^{n+1}$ are formed by $\{T_{\sigma_1}^{-1} Z \cap Z_1\}$ where $Z \in \mathcal{Z}_{\tau\sigma, \star}^n$ and $Z_1 \in \mathcal{Z}_{\sigma, \star}^1$. Now, if $Z_1 \in \mathcal{Z}_{\sigma, f}^1$, the elements maintain the same nature, i.e. if $Z \in \mathcal{Z}_{\tau\sigma, f}^n$ (resp. $\mathcal{Z}_{\tau\sigma, u}^n$) then $T_{\sigma_1}^{-1} Z \cap Z_1 \in \mathcal{Z}_{\sigma, f}^{n+1}$ (resp. $\mathcal{Z}_{\sigma, u}^{n+1}$). So we have in Z_1 at most $C_{\tau\sigma}^n$ contiguous elements of $\mathcal{Z}_{\sigma, u}^{n+1}$. The only problem arises when a block of contiguous elements ends at the boundary of Z_1 since in such a case it can still be contiguous to others elements of $\mathcal{Z}_{\sigma, u}^{n+1}$. Yet, if the contiguous elements of Z_1 are in $\mathcal{Z}_{\sigma, f}^1$, then there can be at most a block of length $2C_{\tau\sigma}^n$. One must then analyze what can happen if $Z_1 \in \mathcal{Z}_{\sigma, u}^1$. In this case, a set of contiguous elements can either have only partial preimage in Z_1 , hence we get a shorter group of contiguous elements, or all the group can have preimage. In this last case, the worst case scenario is when the elements contiguous to the groups (that must belong to $\mathcal{Z}_{\tau\sigma, f}^n$) are cut while taking preimages. This means that at most two new contiguous elements can be generated, but in this case the group must end at the

¹⁴For instance, $\mathcal{W} = \{-1, +1\}$ and whether $[0, \frac{1}{2}]$ and $(\frac{1}{2}, 1]$ correspond to $-1, +1$ or $+1, -1$ respectively is random.

boundary of Z_1 . Since there are at most C_σ^1 contiguous elements in $\mathcal{Z}_{\sigma,u}^1$ in this way we can generate at most $C_\sigma^1(C_{\tau\sigma}^n + 2)$ contiguous elements that, again in the worst case scenario, can be contiguous to two blocks belonging to the neighboring elements in $\mathcal{Z}_{\sigma,f}^1$. Accordingly,

$$C_\sigma^{n+1} \leq C_\sigma^1(C_{\tau\sigma}^n + 2) + 2C_{\tau\sigma}^n = (C_\sigma^1 + 2)C_{\tau\sigma}^n + 2C_\sigma^1 \leq 2 \sum_{i=0}^n (C^{(1)} + 2)^i C^{(1)},$$

where we have used the induction hypothesis. \square

Hence any system is ξ -full branched with $\xi = C^{(1)} + 2$ and $K = \frac{2C^{(1)}}{C^{(1)}+1}$.

Next, we estimate ρ . Remark that $\rho_\sigma \geq \inf \widehat{\mathcal{L}}_{\sigma_1} \mathbf{1}$. Let N to be the minimal number of full branches of T inside H for any $T \in \mathcal{T}$ and $H \in \mathcal{H}$,

$$\widehat{\mathcal{L}}_{\sigma_1} \mathbf{1}(x) = \sum_{T_{\sigma_1} y=x} \mathbf{1}_{H_{\sigma_1}}(y) \frac{1}{|T'_{\sigma_1}(y)|} \geq \frac{N}{M},$$

with $M = \sup_T |T'(x)|$. Thus $\rho \geq \frac{N}{M}$, and condition (C2) is satisfied provided $(C^{(1)} + 2)\Theta < \frac{N}{M}$.

For β transformations $C^{(1)} = 2$, $\Theta = \beta^{-1}$, $M = \varpi\beta$ and $N \geq \lfloor \frac{\beta}{2} \rfloor - 1$. Thus, remembering (6.2), $\rho \geq \frac{\lfloor \frac{\beta}{2} \rfloor - 1}{\varpi\beta}$; $\xi = 4$; $\theta = 4\beta^{-1}$; $K = \frac{4}{3}$ and (C2) is satisfied if $4\varpi\beta < \beta(\lfloor \frac{\beta}{2} \rfloor - 1)$, which happens if $\beta \geq 8\varpi + 4$.

7.1.3. *Condition (C3)*. We start with a parameter selection.

Lemma 7.3. *For each $m \geq 1$, there exists a set $\mathcal{B}_m \subset (1, \infty)$ and $C_m > 0$ such that $\text{Leb}(\mathcal{B}_m \cap (1, t)) \leq C_m \log t$ for all $t > 1$ and if $1 < \beta \notin \mathcal{B}_m$, then*

$$(7.1) \quad \begin{aligned} & \forall n \leq m, \forall \sigma \in \Sigma^{\mathbb{N}}, \forall Z \in \mathcal{Z}_{\sigma,*}^n(\beta), \\ & T_\sigma^{n+1}(Z \cap H_\sigma^{n+1}) = [0, 1] \text{ or } T_\sigma^n Z \subset H_{\sigma_{n+1}}^c. \end{aligned}$$

This lemma, proven shortly, establishes a dichotomy between good and bad elements of $\mathcal{Z}_{\sigma,*}^n(\beta)$ when $\beta \notin \mathcal{B}_m$: either $T_\sigma^n Z \cap H_{\sigma_{n+1}}$ is large enough to cover $[0, 1]$ after one more iteration if Z is good, or it is empty if Z is bad.

Corollary 7.4. *If $\beta \notin \bigcup_{k \leq n} \mathcal{B}_k$ then $\hat{\epsilon}(n, n+1) \geq \frac{1}{(2\varpi\beta)^{n+1}}$.*

Proof. By Lemma 7.3, for $Z \in \mathcal{Z}_{\sigma,g}^k(\beta)$, $k \leq n$, we have $T_\sigma^{n+1}(Z \cap H_\sigma^{n+1}) = T_{\tau^{k+1}\sigma}^{n-k}(T_\sigma^{k+1}(Z \cap H_\sigma^{k+1}) \cap H_{\tau^{k+1}\sigma}^{n-k}) = [0, 1]$ and then

$$\widehat{\mathcal{L}}_\sigma^{n+1} \mathbf{1}_Z(x) = \sum_{T_\sigma^{n+1} y=x} \frac{\mathbf{1}_Z(y) \mathbf{1}_{H_\sigma^{n+1}}(y)}{|(T_\sigma^{n+1})'(y)|} \geq \frac{1}{\sup |(T_\sigma^{n+1})'|} \geq \frac{1}{(\varpi\beta)^{n+1}},$$

and $\widehat{\mathcal{L}}_\sigma^{n+1} \mathbf{1}(x) \leq \mathcal{L}_\sigma^{n+1} \mathbf{1}(x) \leq 2^{n+1}$, since $\mathcal{L}_{T_{\beta_i}} \mathbf{1}(x) \leq 2$. \square

Proof of Proposition 7.1. We choose $\mathcal{B} = (1, \beta_0) \cup \bigcup_{k=1}^6 \mathcal{B}_k$ for some $\beta_0 > 2 \cdot 25^2 \varpi + 1$. Since $K = \frac{4}{3}$, we have $C_* = \frac{25}{3}$. Note that if $\beta \geq \beta_0$, then $\lceil \frac{\ln 4C_*^2}{\ln \rho \theta^{-1}} \rceil < 1$ and, by Remark 6.2, $n_0 = \inf\{n \geq 1 : \epsilon_*(n) > 0\}$. But $\epsilon_*(1) \geq \hat{\epsilon}(1, 2) \geq (2\varpi\beta)^{-2}$, hence $n_0 = 1$. Accordingly, $C_1 \leq \frac{280}{3} \varpi^2 \beta$; $a \leq \frac{42}{11} \varpi^2 \beta^2$ and $B \leq 1 + \frac{700}{11} \varpi^2 \beta^2$ which yields, for all β_0 large enough,

$$n_2 = \left\lceil \frac{\ln 4aB(1 + 2C_1 \rho^{-1})}{\ln \theta^{-1} \rho} \right\rceil \leq \frac{5 \ln \beta + D_1}{\ln \beta - D_2}$$

for some constants $D_i \geq 0$ depending only on ϖ . We can thus choose β_0 large enough so that $n_2 \leq 6$. \square

The rest of the section is devoted to the proof of Lemma 7.3. The basic idea is to discard the β for which the n -th iterates of elements of $\partial\widehat{\mathcal{Z}}_\sigma^n(\beta)$ come too close to $\{0, \frac{1}{2}, 1\}$.

Lemma 7.5. *For all $\sigma \in \Sigma^{\mathbb{N}}$ and $n \geq 1$, one has $T_\sigma^n(\partial\widehat{\mathcal{Z}}_\sigma^n(\beta)) \subset \mathcal{Q}_n(\beta) := \{0, 1\} \cup \{T_{\sigma'}^i(1), T_{\sigma'}^i(\frac{1}{2}) / i = 1, \dots, n, \sigma' \in \Sigma^{\mathbb{N}}\}$.*

Proof. We proceed by induction, the result being clearly true for $n = 1$. Note that $\mathcal{Q}_n(\beta) \cup T_{\tau^n \sigma}^1(\mathcal{Q}_n(\beta)) \subset \mathcal{Q}_{n+1}(\beta)$. Since every $Z \in \widehat{\mathcal{Z}}_\sigma^{n+1}(\beta)$ is of the form $Z = Z' \cap (T_\sigma^n)^{-1}(Z'')$ for $Z' \in \widehat{\mathcal{Z}}_\sigma^n(\beta)$ and $Z'' \in \widehat{\mathcal{Z}}_{\tau^n \sigma}^1(\beta)$, if $a \in \partial\widehat{\mathcal{Z}}_\sigma^{n+1}(\beta)$, then either $a \in \partial\widehat{\mathcal{Z}}_\sigma^n(\beta)$ or $T_\sigma^n a \in \partial\widehat{\mathcal{Z}}_{\tau^n \sigma}^1(\beta)$. In the first case, $T_\sigma^{n+1} a \in T_{\tau^n \sigma}^1(T_\sigma^n(\widehat{\mathcal{Z}}_\sigma^n(\beta))) \subset \mathcal{Q}_{n+1}(\beta)$, and in the second case, $T_\sigma^{n+1} a \in T_{\tau^n \sigma}^1(\partial\widehat{\mathcal{Z}}_{\tau^n \sigma}^1(\beta)) \subset \mathcal{Q}_1(\beta) \subset \mathcal{Q}_{n+1}(\beta)$. \square

We thus see that we need to control all the orbits of 1 and $\frac{1}{2}$. To this end, for $x \in (0, 1]$ and $\sigma \in \Sigma^{\mathbb{N}}$,¹⁵ we introduce the map $\phi_n^{x, \sigma} : (1, \infty) \rightarrow [0, 1]$ defined by $\phi_n^{x, \sigma}(\beta) = T_\sigma^n(x)$.¹⁶ For non negative integers i_1, \dots, i_n , we define

$$I_{i_1, \dots, i_n}^{x, \sigma} = \{\beta \in (1, \infty) \mid \lfloor \beta_{\sigma_k} \phi_{k-1}^{x, \sigma}(\beta) \rfloor = i_k, k = 1, \dots, n\}.$$

Note that $I_{i_1}^{x, \sigma} = [\frac{i_1}{x}, \frac{i_1+1}{x})$ if $\beta_{\sigma_1} = \beta_1$, $I_{i_1}^{x, \sigma} = [\frac{i_1}{\varpi x}, \frac{i_1+1}{\varpi x})$ if $\beta_{\sigma_1} = \beta_2$, and that in both cases, $I_{i_1}^{x, \sigma} \subset [\frac{i_1}{\varpi x}, \frac{i_1+1}{x})$. The family $\{I_{i_1, \dots, i_n, i_{n+1}}^{x, \sigma}\}_{i_{n+1} \geq 0}$ forms a partition into finitely many (at most $\lfloor \varpi \frac{i_1+1}{x} \rfloor + 1$) intervals of $I_{i_1, \dots, i_n}^{x, \sigma}$. From the relation $\phi_{n+1}^{x, \sigma}(\beta) = \beta_{\sigma_{n+1}} \phi_n^{x, \sigma}(\beta) \pmod{1}$, we deduce easily by induction:

Lemma 7.6. *The map $\phi_n^{x, \sigma}$ is \mathcal{C}^1 and strictly increasing on each interval $I_{i_1, \dots, i_n}^{x, \sigma}$, and verifies $(\phi_n^{x, \sigma})'(\beta) \geq \beta^{n-1} x$.*

Lemma 7.7. *For each $x \in (0, 1]$ and $n \geq 1$, there exists $C_{x, n} > 0$ and a set $\mathcal{B}_{n, x} \subset (1, \infty)$, with $\text{Leb}(\mathcal{B}_{n, x} \cap (1, t)) \leq C_{x, n} \log t$, such that, if $\beta \notin \mathcal{B}_{n, x}$,*

$$\forall \sigma \in \Sigma^{\mathbb{N}}, d(\phi_n^{x, \sigma}(\beta), \{0, 1/2, 1\}) > 3\beta^{-1}.$$

Proof. Fix $n \geq 1$ and $\sigma \in \Sigma$, and consider $\beta \in I_{i_1, \dots, i_n}^{x, \sigma}$. We have $\beta^{-1} \leq \varpi x i_1^{-1}$ and thus $d(\phi_n^{x, \sigma}(\beta), \{0, 1/2, 1\}) > 3\beta^{-1}$ whenever

$$(7.2) \quad d(\phi_n^{x, \sigma}(\beta), \{0, 1/2, 1\}) > 3\varpi x i_1^{-1}.$$

Define $\mathcal{B}_{n, i_1}^{x, \sigma}$ to be the set of β in $I_{i_1}^{x, \sigma}$ which do not satisfy (7.2). By Lemma 7.6, the Lebesgue measure of $\mathcal{B}_{n, i_1}^{x, \sigma} \cap I_{i_1, \dots, i_n}^{x, \sigma}$ is less than¹⁷

$$3 \left(\inf_{I_{i_1, \dots, i_n}^{x, \sigma}} |(\phi_n^{x, \sigma})'| \right)^{-1} 3\varpi x i_1^{-1} \leq C_{\varpi, x, n} i_1^{-n}.$$

As $\{I_{i_1, \dots, i_n}^{x, \sigma}\}_{i_2, \dots, i_n}$ forms a partition of $I_{i_1}^{x, \sigma}$ into at most $C_{\varpi, x, n} i_1^{n-1}$ elements, the set $\mathcal{B}_{n, i_1}^{x, \sigma}$ has a measure less than $C_{\varpi, x, n} i_1^{-1}$. For $j = 1, 2$, set $\mathcal{B}_{n, i_1}^{x, j} = \bigcup_{\{\sigma \mid \beta_{\sigma_1} = \beta_j\}} \mathcal{B}_{n, i_1}^{x, \sigma}$.

¹⁵Note that we will only consider $x = \frac{1}{2}$ and $x = 1$ in the following.

¹⁶Recall that $T_{\sigma_i} = T_{\beta_{\sigma_i}}$ and $\beta_1 = \beta$, $\beta_2 = \varpi\beta$.

¹⁷In the following, $C_{\varpi, x, n}$ will denote a constant, the value of which may change from one line to another, depending on ϖ , x and n , but not on i_1 .

For n fixed, the condition (7.2) depends on σ only through its n first terms, and thus $\mathcal{B}_{n,i_1}^{x,j}$ is of measure less than $C_{\omega,x,n}i_1^{-1}$. The set $\mathcal{B}_{n,x} = \bigcup_{j=1,2} \bigcup_{i_1 \geq 1} \mathcal{B}_{n,i_1}^{x,j}$ then satisfies the conclusion of the lemma. \square

We can now conclude the proof:

Proof of Lemma 7.3. We set $\mathcal{B}_m = \bigcup_{1 \leq n \leq m} \mathcal{B}_{n,1/2} \cup \mathcal{B}_{n,1}$ and we proceed by induction over $n \leq m$. Note that if $Z \in \mathcal{Z}_{\sigma,*}^n(\beta)$, then $Z \cap H_\sigma^k = Z$ for all $k \leq n$.

If $Z \in \mathcal{Z}_{\sigma,*}^1(\beta)$, then either Z is a full interval of $\mathcal{Z}_{\sigma,*}^1(\beta)$ and so $T_\sigma^2(Z \cap H_\sigma^2) = T_{\sigma_2}([0,1] \cap H_{\sigma_2}) = [0,1]$, or one of the endpoints of Z is $\frac{1}{2}$ or 1. In the latter case, the other endpoint of Z must be sent after one iteration to 0 or 1 and so if $T_\sigma^1 Z \cap H_{\tau\sigma}^1$ is not empty, then $|T_\sigma^1 Z \cap H_{\tau\sigma}^1| > 3\beta^{-1}$ by Lemma 7.7. The interval $T_\sigma^1 Z \cap H_{\tau\sigma}^1$ therefore contains at least one full interval of $\mathcal{Z}_{\tau\sigma,*}^1(\beta)$, which implies $T_\sigma^2(Z \cap H_\sigma^2) = [0,1]$.

Now, we suppose that (7.1) holds for n and we prove it holds for $n+1 < m$.

Any $Z \in \mathcal{Z}_{\sigma,*}^{n+1}(\beta)$ is of the form $Z = Z' \cap (T_\sigma^n)^{-1}(Z'')$ with $Z' \in \mathcal{Z}_{\sigma,*}^n(\beta)$ and $Z'' \in \mathcal{Z}_{\tau^n\sigma,*}^1(\beta)$. If both endpoints of Z belong to the interior of Z' , then $T_\sigma^n Z = Z'' \in \mathcal{Z}_{\tau^n\sigma,*}^1(\beta)$ and we have

$$T_\sigma^{n+2}(Z \cap H_\sigma^{n+2}) = T_{\tau^n\sigma}^2(T_\sigma^n Z \cap H_{\tau^n\sigma}^2) = T_{\tau^n\sigma}^2(Z'' \cap H_{\tau^n\sigma}^2) = [0,1],$$

or

$$T_\sigma^{n+1} Z = T_{\tau^n\sigma}^1(T_\sigma^n Z) = T_{\tau^n\sigma}^1 Z'' \subset H_{\sigma_{n+1}}^c,$$

according to whether Z'' is a good a or bad element of $\mathcal{Z}_{\tau^n\sigma,*}^1(\beta)$ respectively.

If one endpoint of Z is also an endpoint of Z' , then $T_\sigma^n Z = [T_\sigma^n a, b]$ ¹⁸ with $a \in \partial\mathcal{Z}_{\sigma,*}^n(\beta)$ and $b \in \partial\mathcal{Z}_{\tau^n\sigma,*}^1(\beta)$. By Lemma 7.5, $T_\sigma^n a = T_\sigma^i x$, with $\sigma' \in \Sigma$, $0 \leq i \leq n$ and $x \in \{\frac{1}{2}, 1\}$.¹⁹ We consider two subcases: either $b \notin \{\frac{1}{2}, 1\}$ or $b \in \{\frac{1}{2}, 1\}$.

In the first subcase, we have $y := T_{\sigma_{n+1}} b \in \{0, 1\}$. Therefore, $T_\sigma^{n+1} Z = [T_{\sigma_{n+1}} T_\sigma^i x, y]$. Since $i+1 \leq n+1 \leq m$, $\beta \notin \mathcal{B}_{n+1,y}$ and one of the endpoints of $T_\sigma^{n+1} Z$ belongs to $\{0, 1\}$, it follows that $T_\sigma^{n+1} Z \cap H_{\sigma_{n+2}}$, if it is non empty, is an interval of length strictly larger than $3\beta^{-1}$ by Lemma 7.7 and thus contains at least one full interval of $\mathcal{Z}_{\tau^{n+1}\sigma,*}^1(\beta)$. Consequently, $T_\sigma^{n+2}(Z \cap H_\sigma^{n+2}) = [0,1]$.

In the second subcase, $T_\sigma^n Z = [T_\sigma^i x, b]$ has one endpoint belonging to $\{0, \frac{1}{2}, 1\}$, and so, as above, we obtain that $T_\sigma^n Z \cap H_{\sigma_n}$, if it is non empty, is of length strictly larger than $3\beta^{-1}$ by Lemma 7.7. We deduce that $T_\sigma^n Z \cap H_{\sigma_n}$ contains at least one full interval of $\mathcal{Z}_{\tau^n\sigma,*}^1(\beta)$, which implies that $T_\sigma^{n+2}(Z \cap H_\sigma^{n+2}) = [0,1]$.

Finally, if both endpoints of Z are also endpoints of Z' , then $Z = Z' \in \mathcal{Z}_{\sigma,*}^n(\beta)$. Since $Z \in \mathcal{Z}_{\sigma,*}^{n+1}(\beta)$, we have $T_\sigma^n Z \subset H_{\sigma_{n+1}}$ and so by our induction hypothesis, we can only have $T_\sigma^{n+1} Z = T_\sigma^{n+1}(Z' \cap H_\sigma^{n+1}) = [0,1]$. This implies $T_\sigma^{n+2}(Z \cap H_\sigma^{n+2}) = [0,1]$ and concludes the induction. \square

7.2. Markov maps with non-Markov gates. The reader might wonder if it is possible to produce an example more similar to the Lorentz gas. In particular, one in which the invariant measure of the maps is always the same and the gates are deterministic, so that one knows explicitly the invariant measure of the process of the environment as seen from the particle.

¹⁸We write $[x,y]$ to denote the interval joining x and y , disregarding whether $x \leq y$ or $x \geq y$.

¹⁹Note that if $T_\sigma^n a = 0$, then we are reduced to the previous situation.

This is indeed possible: let $\mathcal{T} = \{T_{\beta_1}, T_{\beta_2}\}$, with $\beta_2 > \beta_1$ both integers, and partitions $\mathcal{G}_\alpha = \mathcal{G}$ deterministic with $\mathcal{W} = \{-1, 0, +1\}$ and $G_{-1} = [0, y]$, $G_0 = (y, 1 - y]$ and $G_{+1} = (1 - y, 1]$, for $0 < y < \frac{1}{2}$. Proceeding similarly to the previous section, but using y as a parameter, instead of β , we have:

Proposition 7.8. *Let $\varpi > 1$. Then for each $1 < \beta_1 < \beta_2 < \varpi\beta_1$ integers, there exists a measurable set $\mathcal{B}_{\beta_1, \beta_2} \subset (0, \frac{1}{2})$, with $\text{Leb}(\mathcal{B}_{\beta_1, \beta_2}) = \mathcal{O}(\beta_1^{-1})$ as $\beta_1 \rightarrow \infty$, such that the model described above, with $y \notin \mathcal{B}_{\beta_1, \beta_2}$, satisfies the assumptions of Theorems 6.1.*

In particular, this class of models is non-empty when β_1 is large enough. Since all maps in \mathcal{T} preserve the Lebesgue measure and all gates are deterministic, Theorem 6.4 also applies²⁰ and we therefore have $\frac{1}{n}z(n) \rightarrow 0$ a.e., since the drift is equal to

$$V = \sum_{w \in \mathcal{W}} w \int_{G_w} h_0 dm = \sum_{w \in \{-1, 0, +1\}} w |G_w| = 0.$$

By corollary 4.11, the walk (z_n) is then recurrent.

8. EQUIVALENCE WITH A GIBBS RANDOM WALK

In this section, we prove Theorem 6.1. The proof will rely on a property of exponential loss of memory for compositions of the operators $\widehat{\mathcal{L}}_\sigma$. More precisely, we will investigate the properties of compositions of the form $\widehat{\mathcal{L}}_\sigma^n f$, in order to understand better the asymptotics of the probabilities $p(\bar{\omega}, n, w_0 \dots w_{n-1})$. For convenience, we will consider bi-infinite sequences $\sigma \in \Sigma^{\mathbb{Z}} = (\mathcal{T} \times \mathcal{H})^{\mathbb{Z}}$, with $\tau : \Sigma^{\mathbb{Z}} \rightarrow \Sigma^{\mathbb{Z}}$ the bilateral shift. We can extend the definitions of $\widehat{\mathcal{L}}_\sigma^n$, ρ_σ , C_σ^a and Λ_σ to the case $\sigma \in \Sigma^{\mathbb{Z}}$ in a straightforward way. What we mean by loss of memory is made precise by the following result proven in Section 9.

Proposition 8.1. *Under the hypothesis of Theorem 6.1 there exist $\nu \in (0, 1)$, a family of positive numbers $\{\rho_\sigma\}_{\sigma \in \Sigma^{\mathbb{Z}}}$ and a family of positive functions $\{h_\sigma\}_{\sigma \in \Sigma^{\mathbb{Z}}}$ in BV such that $\Lambda_\sigma(h_\sigma) = 1$ and for all $\sigma \in \Sigma^{\mathbb{Z}}$, $f \in \text{BV}$ and $n \geq 0$:*

$$(8.1) \quad \left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\rho_\sigma \cdots \rho_{\tau^{n-1}\sigma}} - \Lambda_\sigma(f) h_{\tau^n \sigma} \right\|_\infty \leq C_{\#} \nu^n \|f\|_{\text{BV}}.$$

Remark 8.2. *The above statement is similar to the one in [21]. Note however that here the setting is very different insofar in [21] only small holes and nearby maps are considered. The upgrade of the result to large holes and arbitrary maps (as we inescapably need) turns out to be highly non trivial.*

In the following we will need the following information on Λ_σ and h_σ , which are proven at the end of Section 9.

Lemma 8.3. *Under the hypothesis of Theorem 6.1 there exist $\varrho > 0$ such that, for all $n \geq 0$ and $\sigma \in \Sigma^{\mathbb{Z}}$, $\inf h_\sigma \geq \varrho$ and $\|h_\sigma\|_\infty \leq 1 + a$.²¹ In addition, $\sup_\sigma \rho_\sigma < \infty$ and for all $f \in \text{BV}$, $\Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n f) = \rho_\sigma \cdots \rho_{\tau^{n-1}\sigma} \Lambda_\sigma(f)$; $\widehat{\mathcal{L}}_\sigma^n h_\sigma = \rho_\sigma \cdots \rho_{\tau^{n-1}\sigma} h_{\tau^n \sigma}$.*

²⁰Condition **(Pro)** is satisfied here, since \mathcal{W} is symmetric.

²¹The constant a is defined in Remark 6.2

To prove Proposition 8.1, we will adapt the strategy of [17]. More precisely, we will show that the family of cones

$$C_\sigma^a = \{h \in \text{BV} \mid h \neq 0, h \geq 0, \bigvee h \leq a\Lambda_\sigma(h)\}$$

is strictly invariant under compositions of large enough length of transfer operators (i.e. $\widehat{\mathcal{L}}_\sigma^n C_\sigma^a \subset C_{\tau^n \sigma}^{a/2}$ for all $n \geq n_0$) for a suitable $a > 0$, see Lemma 9.5. From this, we will deduce that $\widehat{\mathcal{L}}_\sigma^n C_\sigma^a$ has uniform finite diameter in $C_{\tau^n \sigma}^a$ for the corresponding Hilbert metric (Lemma 9.14), which will imply that $\widehat{\mathcal{L}}_\sigma^n$ is a strict contraction for the Hilbert metric, and then enjoys exponential loss of memory.

To deduce Theorem 6.1 from Proposition 8.1, we will also need the following technical lemma, which will be proven at the end of Section 9:

Lemma 8.4. *There exists $\varrho' > 0$ such that for all $n \geq 0$ and $\sigma \in \Sigma^{\mathbb{N}}$, we have $\inf \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \geq \varrho' \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})$ and $\|\widehat{\mathcal{L}}_\sigma^n \mathbf{1}\|_\infty \leq B \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})$.²²*

Recall from Section 4 that the transition probabilities are given by (4.10). For $(\bar{\omega}, \bar{w}) \in \Omega_*$, we define $\sigma = \sigma(\bar{\omega}, \bar{w}) \in \Sigma^{\mathbb{N}}$ such that $\widehat{\mathcal{L}}_{\sigma_n} = \mathcal{L}_{\bar{\omega}, z_{n-1}, \bar{w}_{n-1}}$, so that

$$p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) = \int \widehat{\mathcal{L}}_\sigma^n h_0 dm.$$

We will still denote by $\sigma \in \Sigma^{\mathbb{Z}}$ an arbitrary element of $\Sigma^{\mathbb{Z}}$ which coincides with σ for future components (for instance, given an arbitrary $\sigma_* \in \Sigma^{\mathbb{Z}}$, we identify $\sigma \in \Sigma^{\mathbb{N}}$ with the element $\tilde{\sigma} \in \Sigma^{\mathbb{Z}}$ defined by $\tilde{\sigma}_i = \sigma_i$ if $i \geq 1$ and $\tilde{\sigma}_i = (\sigma_*)_i$ if $i \leq 0$).

We are now ready to prove the announced results.

Proof of Theorem 6.1. We have

$$(8.2) \quad \mathbb{P}_{\bar{\omega}}(\bar{w}_n \mid \bar{w}_0 \dots \bar{w}_{n-1}) = \frac{p(\bar{\omega}, n+1, \bar{w}_0 \dots \bar{w}_n)}{p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1})} = \frac{\int \widehat{\mathcal{L}}_\sigma^{n+1} h_0 dm}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm}$$

and

$$(8.3) \quad \begin{aligned} \mathbb{P}_{\tau_{z_m} \bar{\omega}}(\bar{w}_n \mid \bar{w}_m \dots \bar{w}_{n-1}) &= \frac{p(\tau_{z_m} \bar{\omega}, n-m+1, \bar{w}_m \dots \bar{w}_n)}{p(\tau_{z_m} \bar{\omega}, n-m, \bar{w}_m \dots \bar{w}_{n-1})} \\ &= \frac{\int \widehat{\mathcal{L}}_{\tau_{z_m} \sigma}^{n-m+1} h_0 dm}{\int \widehat{\mathcal{L}}_{\tau_{z_m} \sigma}^{n-m} h_0 dm}. \end{aligned}$$

We can then write

$$\frac{\int \widehat{\mathcal{L}}_\sigma^{n+1} h_0 dm}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm} = \rho_{\tau^n \sigma} \frac{\int \widehat{\mathcal{L}}_\sigma^{n+1} h_0 dm}{\rho_\sigma \dots \rho_{\tau^{n-1} \sigma}} \frac{\rho_\sigma \dots \rho_{\tau^{n-1} \sigma}}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm}.$$

Using Proposition 8.1, we have

$$(8.4) \quad \left| \frac{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm}{\rho_\sigma \dots \rho_{\tau^{n-1} \sigma}} - \Lambda_\sigma(h_0) \int h_{\tau^n \sigma} dm \right| \leq C_{\#} \nu^n \|h_0\|_{\text{BV}},$$

which also implies

$$(8.5) \quad \left| \frac{\rho_\sigma \dots \rho_{\tau^{n-1} \sigma}}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm} - \left(\Lambda_\sigma(h_0) \int h_{\tau^n \sigma} dm \right)^{-1} \right| \leq C_{\#} \nu^n \frac{\|h_0\|_{\text{BV}}}{(\inf h_0)^2},$$

²² The constant B is defined in Remark 6.2.

since, by Lemmata 8.3 and 8.4,

$$\frac{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm}{\rho_\sigma \cdots \rho_{\tau^{n-1}\sigma}} \geq (\inf h_0) \frac{\inf \widehat{\mathcal{L}}_\sigma^n \mathbf{1}}{\rho_\sigma \cdots \rho_{\tau^{n-1}\sigma}} \geq (\inf h_0) \frac{\varrho' \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}{\rho_\sigma \cdots \rho_{\tau^{n-1}\sigma}} = \varrho'(\inf h_0),$$

and $\Lambda_\sigma(h_0) \int h_{\tau^n \sigma} dm \geq \varrho \inf h_0$.

Note that we also have, using again Lemmata 8.4 and 8.3,

$$\frac{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm}{\rho_\sigma \cdots \rho_{\tau^{n-1}\sigma}} \leq \|h_0\|_\infty \frac{\|\widehat{\mathcal{L}}_\sigma^n \mathbf{1}\|_\infty}{\rho_\sigma \cdots \rho_{\tau^{n-1}\sigma}} \leq B \|h_0\|_\infty.$$

Consequently, using (8.4) with n replaced by $n+1$, (8.5), Lemma 8.3 and the above inequalities:

$$\begin{aligned} & \left| \frac{\int \widehat{\mathcal{L}}_\sigma^{n+1} h_0 dm}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm} - \rho_{\tau^n \sigma} \frac{\int h_{\tau^{n+1}\sigma} dm}{\int h_{\tau^n \sigma} dm} \right| \leq \left[\left| \frac{\int \widehat{\mathcal{L}}_\sigma^{n+1} h_0 dm}{\rho_\sigma \cdots \rho_{\tau^n \sigma}} - \Lambda_\sigma(h_0) \int h_{\tau^{n+1}\sigma} dm \right| \right. \\ & \times \left. \frac{\rho_\sigma \cdots \rho_{\tau^n \sigma}}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm} + \rho_{\tau^n \sigma} \Lambda_\sigma(h_0) \int h_{\tau^{n+1}\sigma} dm \left| \frac{\rho_\sigma \cdots \rho_{\tau^{n-1}\sigma}}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm} - \frac{1}{\Lambda_\sigma(h_0) \int h_{\tau^n \sigma} dm} \right| \right] \\ & \leq C_\# \nu^n \left(\frac{\|h_0\|_{\text{BV}}}{\inf h_0} + \frac{\|h_0\|_{\text{BV}}^2}{(\inf h_0)^2} \right). \end{aligned}$$

Hence, there exists C_{h_0} depending only on h_0 such that, for all $n \geq 0$,

$$(8.6) \quad \sup_{\sigma \in \Sigma} \left| \frac{\int \widehat{\mathcal{L}}_\sigma^{n+1} h_0 dm}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm} - \rho_{\tau^n \sigma} \frac{\int h_{\tau^{n+1}\sigma} dm}{\int h_{\tau^n \sigma} dm} \right| \leq C_{h_0} \nu^n.$$

Equation (8.6), by (8.2) and (8.3), implies **(Exp)**. \square

Proof of Theorem 6.3. By (8.6), we have

$$\sup_{\sigma \in \Sigma} \left| \frac{\int \widehat{\mathcal{L}}_\sigma^{n+1} h_0 dm}{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm} - \frac{\int \widehat{\mathcal{L}}_\sigma^{n+1} h'_0 dm}{\int \widehat{\mathcal{L}}_\sigma^n h'_0 dm} \right| \leq (C_{h_0} + C_{h'_0}) \nu^n,$$

for all $n \geq 0$, and the theorem follows with $C_{h_0, h'_0} = C_{h_0} + C_{h'_0}$. \square

Proof of Theorem 6.4. By Theorem 4.9, the discussion in Section 4.4, and since **(Exp)** already holds by Theorem 6.1, it is enough to show that assumptions **(Pos)** and **(Ell)** are satisfied for the probabilities defined by (4.10).

Verification of (Pos). By Proposition 8.1, Lemma 8.3 and equation (6.2)

$$\begin{aligned} p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) &= \int \widehat{\mathcal{L}}_\sigma^n h_0 dm = \rho_\sigma \cdots \rho_{\tau^{n-1}\sigma} \frac{\int \widehat{\mathcal{L}}_\sigma^n h_0 dm}{\rho_\sigma \cdots \rho_{\tau^{n-1}\sigma}} \\ &\geq \rho^n \left(\Lambda_\sigma(h_0) \int h_{\tau^n \sigma} - C_\# \nu^n \|h_0\|_{\text{BV}} \right) \geq \rho^n (\varrho \inf h_0 - C_\# \nu^n \|h_0\|_{\text{BV}}). \end{aligned}$$

Consequently, $p(\bar{\omega}, n, \bar{w}_0 \dots \bar{w}_{n-1}) > 0$ for all n large enough, and since this quantity is non-increasing, this proves the positivity for all $n \geq 0$.

Verification of **(EII)**. By (8.6) and Lemma 8.3, we have for all $n \geq 0$,

$$\begin{aligned} \mathbb{P}_{\bar{\omega}}(\bar{w}_n \mid \bar{w}_{n-1} \dots \bar{w}_0) &= \frac{\int \widehat{\mathcal{L}}_{\sigma}^{n+1} h_0 dm}{\int \widehat{\mathcal{L}}_{\sigma}^n h_0 dm} \geq \rho_{\tau^n \sigma} \frac{\int h_{\tau^{n+1} \sigma} dm}{\int h_{\tau^n \sigma} dm} - C_{h_0} \nu^n \\ &\geq \rho \frac{\varrho}{1+a} - C_{h_0} \nu^n, \end{aligned}$$

which proves **(EII)**. \square

9. LOSS OF MEMORY

To prove Proposition 8.1, we will adapt the strategy of [17] to our non-stationary case, and employ the theory of Hilbert metrics, that we recall below.

Definition 2. Let \mathcal{V} be a vector space. We will call convex cone a subset $\mathcal{C} \subset \mathcal{V}$ which enjoys the following properties:

- (i) $\mathcal{C} \cap -\mathcal{C} = \emptyset$.
- (ii) $\forall \lambda > 0, \lambda \mathcal{C} = \mathcal{C}$.
- (iii) \mathcal{C} is a convex set.
- (iv) $\forall f, g \in \mathcal{C}, \forall \alpha_n \in \mathbb{R} \alpha_n \rightarrow \alpha, g - \alpha_n f \in \mathcal{C} \Rightarrow g - \alpha f \in \mathcal{C} \cup \{0\}$.

We now define the Hilbert metric on \mathcal{C} :

Definition 3. The distance $d_{\mathcal{C}}(f, g)$ between two points f, g in \mathcal{C} is given by

$$\begin{aligned} \alpha(f, g) &= \sup\{\lambda > 0 \mid g - \lambda f \in \mathcal{C}\}, \\ \beta(f, g) &= \inf\{\mu > 0 \mid \mu f - g \in \mathcal{C}\}, \\ d_{\mathcal{C}}(f, g) &= \log \frac{\beta(f, g)}{\alpha(f, g)}, \end{aligned}$$

where we take $\alpha = 0$ or $\beta = \infty$ when the corresponding sets are empty.

The next theorem shows that every positive linear operator is a contraction, provided that the diameter of the image is finite.

Theorem 9.1 ([15, Theorem 1.1]). Let \mathcal{V}_1 and \mathcal{V}_2 be two vector spaces, $\mathcal{C}_1 \subset \mathcal{V}_1$ and $\mathcal{C}_2 \subset \mathcal{V}_2$ two convex cones and $L : \mathcal{V}_1 \rightarrow \mathcal{V}_2$ a positive linear operator (which implies $L(\mathcal{C}_1) \subset \mathcal{C}_2$). If we denote

$$\Delta = \sup_{f, g \in L(\mathcal{C}_1)} d_{\mathcal{C}_2}(f, g),$$

then

$$d_{\mathcal{C}_2}(Lf, Lg) \leq \tanh\left(\frac{\Delta}{4}\right) d_{\mathcal{C}_1}(f, g) \quad \forall f, g \in \mathcal{C}_1.$$

The following lemma links the Hilbert metric to suitable norms on \mathcal{V} :

Lemma 9.2 ([18, Lemma 2.2]). Let $\|\cdot\|$ be a norm on \mathcal{V} such that

$$\forall f, g \in \mathcal{V} \quad g - f, g + f \in \mathcal{C} \Rightarrow \|f\| \leq \|g\|$$

and let $\ell : \mathcal{C} \rightarrow \mathbb{R}^+$ be a homogeneous and order preserving function, i.e.

$$\begin{aligned} \forall f \in \mathcal{C}, \forall \lambda \in \mathbb{R}^+ \quad \ell(\lambda f) &= \lambda \ell(f), \\ \forall f, g \in \mathcal{C} \quad g - f \in \mathcal{C} \quad \ell(f) &\leq \ell(g), \end{aligned}$$

then

$$\forall f, g \in \mathcal{C} \quad \ell(f) = \ell(g) > 0 \Rightarrow \|f - g\| \leq (e^{d_{\mathcal{C}}(f, g)} - 1) \min(\|f\|, \|g\|).$$

From now on, we will always assume that conditions (C1) and (C2) hold. Our main tool will be the following Lasota-Yorke type inequality:

Lemma 9.3. *If condition (C3(N, N')) holds, then for any $n \leq N$, for any $\sigma \in \Sigma^{\mathbb{N}}$ and $h \in \text{BV}$, we have*

$$\bigvee \widehat{\mathcal{L}}_{\sigma}^n h \leq C_{\star} (\xi \Theta)^n \bigvee h + C_N \Lambda_{\sigma}(|h|),$$

where $C_{\star} = 3(C + 1) + 2K(3C + 2)$, $C_N = \frac{(3C+2)(2K\xi+1)\Theta}{\epsilon_{\star}(N)}$, and C is such that $\bigvee_Z g_{\sigma}^n \leq C \|g_{\sigma}^n\|_{\infty}$ for all n, σ and $Z \in \mathcal{Z}_{\sigma, \star}^n$.²³

Proof. Remark that the case $n = 0$ is immediate, since $C_{\star} \geq 1$ and $C_N \geq 0$. We thus only consider $n \geq 1$.

First notice that $\widehat{\mathcal{L}}_{\sigma}^n(h \mathbf{1}_Z) = 0$ if $Z \in \widehat{\mathcal{Z}}_{\sigma}^n \setminus \mathcal{Z}_{\sigma, \star}^n$. We can then write

$$\widehat{\mathcal{L}}_{\sigma}^n h = \sum_{Z \in \mathcal{Z}_{\sigma, \star}^n} \widehat{\mathcal{L}}_{\sigma}^n(\mathbf{1}_Z h) = \sum_{Z \in \mathcal{Z}_{\sigma, \star}^n} (\mathbf{1}_Z g_{\sigma}^n h) \circ (T_{\sigma, Z}^n)^{-1},$$

where $(T_{\sigma, Z}^n)^{-1}$ is the inverse branch of T_{σ}^n restricted to Z . Accordingly,

$$\bigvee \widehat{\mathcal{L}}_{\sigma}^n h \leq \sum_{Z \in \mathcal{Z}_{\sigma, \star}^n} \bigvee \mathbf{1}_{T_{\sigma}^n Z}(g_{\sigma}^n h) \circ (T_{\sigma, Z}^n)^{-1}.$$

We estimate each term of the sum separately.

$$\begin{aligned} \bigvee \mathbf{1}_{T_{\sigma}^n Z}(g_{\sigma}^n h) \circ (T_{\sigma, Z}^n)^{-1} &\leq \bigvee_Z h g_{\sigma}^n + 2 \sup_Z |h g_{\sigma}^n| \\ &\leq 3 \bigvee_Z h g_{\sigma}^n + 2 \inf_Z |h g_{\sigma}^n| \\ &\leq 3 \|g_{\sigma}^n\|_{\infty} \bigvee_Z h + 3 \sup_Z |h| \bigvee_Z g_{\sigma}^n + 2 \inf_Z |h g_{\sigma}^n| \\ &\leq 3 \|g_{\sigma}^n\|_{\infty} \bigvee_Z h + 3C \sup_Z |h| \|g_{\sigma}^n\|_{\infty} + 2 \|g_{\sigma}^n\|_{\infty} \inf_Z |h| \\ &\leq 3(C + 1) \|g_{\sigma}^n\|_{\infty} \bigvee_Z h + (3C + 2) \|g_{\sigma}^n\|_{\infty} \inf_Z |h|. \end{aligned}$$

By assumption (C3(N, N')), we have for each $x \in [0, 1]$,

$$\inf_{Z \in \mathcal{Z}_{\sigma, g}^n} \frac{\widehat{\mathcal{L}}_{\sigma}^{N'} \mathbf{1}_Z(x)}{\widehat{\mathcal{L}}_{\sigma}^{N'} \mathbf{1}(x)} \geq \hat{\epsilon}(N, N') \geq \epsilon_{\star}(N) > 0.$$

Accordingly, for each $x \in [0, 1]$, $h \in \text{BV}$ and $Z \in \mathcal{Z}_{\sigma, g}^n$ holds

$$\widehat{\mathcal{L}}_{\sigma}^{N'}(|h| \mathbf{1}_Z)(x) \geq \inf_Z |h| \widehat{\mathcal{L}}_{\sigma}^{N'} \mathbf{1}_Z(x) \geq \inf_Z |h| \epsilon_{\star}(N) \widehat{\mathcal{L}}_{\sigma}^{N'} \mathbf{1}(x).$$

To deal with elements in $\mathcal{Z}_{\sigma, b}^n$, we use condition (C2) which insures that elements of $\mathcal{Z}_{\sigma, g}^n$ can be separated by at most $K\xi^n$ elements of $\mathcal{Z}_{\sigma, b}^n$. For each $Z \in \mathcal{Z}_{\sigma, b}^n$, let $I_{\pm}(Z)$ be the union of the contiguous elements of $\mathcal{Z}_{\sigma, b}^n$ on the left and on the right of Z respectively. Clearly, for each $Z' \subset I_{\pm}(Z)$, holds

$$\inf_{Z'} |h| \leq \inf_Z |h| + \bigvee_{I_{\pm}(Z)} h.$$

²³The constant C exists by the usual bounded distortion estimates. In particular, note that if all the maps in \mathcal{T} are piecewise linear, then $C = 0$.

Accordingly,

$$\sum_{Z \in \mathcal{Z}_{\sigma, b}^n} \inf_Z |h| \leq 2K\xi^n \left[\sum_{Z \in \mathcal{Z}_{\sigma, g}^n} \inf_Z |h| + \bigvee h \right].$$

For all x , we thus have

$$\begin{aligned} \sum_{Z \in \mathcal{Z}_{\sigma, \star}^n} \inf_Z |h| &\leq (2K\xi^n + 1) \sum_{Z \in \mathcal{Z}_{\sigma, g}^n} \inf_Z |h| + 2K\xi^n \bigvee h \\ &\leq (2K\xi^n + 1) \frac{1}{\epsilon_{\star}(N)} \sum_{Z \in \mathcal{Z}_{\sigma, g}^n} \frac{\widehat{\mathcal{L}}_{\sigma}^{N'}(|h|\mathbf{1}_Z)(x)}{\widehat{\mathcal{L}}_{\sigma}^{N'}\mathbf{1}(x)} + 2K\xi^n \bigvee h \\ &\leq (2K\xi^n + 1) \frac{1}{\epsilon_{\star}(N)} \frac{\widehat{\mathcal{L}}_{\sigma}^{N'}(|h|)(x)}{\widehat{\mathcal{L}}_{\sigma}^{N'}\mathbf{1}(x)} + 2K\xi^n \bigvee h. \end{aligned}$$

We can then conclude

$$\begin{aligned} \bigvee \widehat{\mathcal{L}}_{\sigma}^n h &\leq [3(C+1) + (3C+2)2K\xi^n] \|g_{\sigma}^n\|_{\infty} \bigvee h \\ &\quad + (3C+2)(2K\xi^n + 1) \|g_{\sigma}^n\|_{\infty} \frac{1}{\epsilon_{\star}(N)} \frac{\widehat{\mathcal{L}}_{\sigma}^{N'}(|h|)(x)}{\widehat{\mathcal{L}}_{\sigma}^{N'}\mathbf{1}(x)}. \end{aligned}$$

Taking the inf over x and recalling that $\|g_{\sigma}^n\|_{\infty} \leq \Theta^n$, $\inf_x \frac{\widehat{\mathcal{L}}_{\sigma}^{N'}(|h|)(x)}{\widehat{\mathcal{L}}_{\sigma}^{N'}\mathbf{1}(x)} \leq \Lambda_{\sigma}(|h|)$, $\xi \geq 1$ and $\xi\Theta < 1$ by (C2), we obtain the result. \square

We will show that the family of cones

$$\mathcal{C}_{\sigma}^a = \{h \in \text{BV} \mid h \neq 0, h \geq 0, \bigvee h \leq a\Lambda_{\sigma}(h)\}$$

is strictly invariant under the transfer operators defined above.

Recall that, under assumption (C2), $\theta = \xi\Theta < \rho$.

Lemma 9.4. *For all σ and all $g \in \text{BV}$, $g \geq 0$, we have $\Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_{\sigma_1} g) \geq \rho_{\sigma}\Lambda_{\sigma}(g)$. In particular, $\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_{\sigma}^n g) \geq \rho^n\Lambda_{\sigma}(g)$ and $\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_{\sigma}^n \mathbf{1}) \geq \rho^n$.*

Proof. If we prove the first part of the statement, the second part follows by iteration, since $\rho_{\sigma} \geq \rho$ for all σ . For each $g \in \text{BV}$, $g \geq 0$ and $x \in [0, 1]$,

$$\frac{\widehat{\mathcal{L}}_{\tau\sigma}^n \widehat{\mathcal{L}}_{\sigma_1} g(x)}{\widehat{\mathcal{L}}_{\tau\sigma}^n \mathbf{1}(x)} \geq \frac{\widehat{\mathcal{L}}_{\sigma_{n+1}} \left[\frac{\widehat{\mathcal{L}}_{\sigma}^n g \widehat{\mathcal{L}}_{\sigma}^n \mathbf{1}}{\widehat{\mathcal{L}}_{\sigma}^n \mathbf{1}} \right] (x)}{\widehat{\mathcal{L}}_{\tau\sigma}^n \mathbf{1}(x)} \geq \frac{\widehat{\mathcal{L}}_{\tau\sigma}^n (\widehat{\mathcal{L}}_{\sigma_1} \mathbf{1})(x)}{\widehat{\mathcal{L}}_{\tau\sigma}^n \mathbf{1}(x)} \inf \frac{\widehat{\mathcal{L}}_{\sigma}^n g}{\widehat{\mathcal{L}}_{\sigma}^n \mathbf{1}}$$

and taking the inf on x and the limit $n \rightarrow \infty$, we get the result. \square

Lemma 9.5. *Let $n_0 \geq \lceil \frac{\ln 4C_{\star}^2}{\ln \rho\theta^{-1}} \rceil$ such that C3(n_0, N') holds for some $N' \geq n_0$, then for all $a \geq a_0 = \frac{15}{11} \max_{i \leq n_0} \frac{C_i}{C_{\star}^{\theta^i}}$ and σ , we have*

$$\widehat{\mathcal{L}}_{\sigma}^n \mathcal{C}_{\sigma}^a \subset C_{\tau^n\sigma}^{2aC_{\star}} \quad \forall n \geq 0 \quad \text{and} \quad \widehat{\mathcal{L}}_{\sigma}^n \mathcal{C}_{\sigma}^a \subset C_{\tau^n\sigma}^{a/2} \quad \forall n \geq n_0.$$

Proof. Let $n_0 \in \mathbb{N}$ which will be chosen later. Let $h \in \mathcal{C}_\sigma^a$, then we can write each n as $n = kn_0 + m$, $m < n_0$, and by Lemma 9.3, we have

$$\begin{aligned}
\bigvee \widehat{\mathcal{L}}_\sigma^n h &\leq C_\star \theta^{n_0} \bigvee \widehat{\mathcal{L}}_\sigma^{(k-1)n_0+m} h + C_{n_0} \Lambda_{\tau^{(k-1)n_0+m}\sigma}(\widehat{\mathcal{L}}_\sigma^{(k-1)n_0+m} h) \\
&\leq C_\star^k \theta^{kn_0} \bigvee \widehat{\mathcal{L}}_\sigma^m h + \sum_{i=0}^{k-1} C_{n_0} (C_\star \theta^{n_0})^i \Lambda_{\tau^{(k-i-1)n_0+m}\sigma}(\widehat{\mathcal{L}}_\sigma^{(k-i-1)n_0+m} h) \\
(9.1) \quad &\leq C_\star^{k+1} \theta^n \bigvee h + \sum_{i=0}^{k-1} C_{n_0} (C_\star \theta^{n_0})^i \Lambda_{\tau^{(k-i-1)n_0+m}\sigma}(\widehat{\mathcal{L}}_\sigma^{(k-i-1)n_0+m} h) \\
&\quad + C_m (C_\star \theta^{n_0})^k \Lambda_\sigma(h).
\end{aligned}$$

Using Lemma 9.4, we obtain

$$\bigvee \widehat{\mathcal{L}}_\sigma^n h \leq \left[\left(a + \frac{C_m}{C_\star \theta^m} \right) \frac{C_\star^{k+1} \theta^n}{\rho^n} + \frac{C_{n_0}}{\rho^{n_0}} \sum_{i=0}^{k-1} \left(\frac{C_\star \theta^{n_0}}{\rho^{n_0}} \right)^i \right] \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n h).$$

If $k = 0$ we have

$$\bigvee \widehat{\mathcal{L}}_\sigma^n h \leq 2a C_\star \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n h).$$

For $k > 0$ let us set $\tau = \theta \rho^{-1}$, by (C2) $\tau < 1$, and let n_0 such that $\alpha = C_\star^2 \tau^{n_0} \leq \frac{1}{4}$. Since $C_\star \geq 3$ then $C_\star \tau^{n_0} \leq \frac{1}{12}$. Hence

$$\bigvee \widehat{\mathcal{L}}_\sigma^n h \leq \left\{ \frac{1}{4} a + a_0 \frac{11}{15} \left[\frac{1}{4} + \frac{C_\star^2 \tau^{n_0}}{C_\star - \alpha} \right] \right\} \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n h) \leq \frac{1}{2} a \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n h).$$

□

Lemma 9.6. *Let $B = 1 + 2a_0 C_\star$. If (C3(n_0, N')) holds, then for each $h \in \text{BV}$, $h \geq 0$, $n \in \mathbb{N}$ and $\sigma \in \Sigma^{\mathbb{N}}$,*

$$\Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1}) \Lambda_\sigma(h) \leq \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n h) \leq B \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1}) \Lambda_\sigma(h).$$

Proof. For $x \in [0, 1]$, we have

$$\frac{\widehat{\mathcal{L}}_{\tau^n \sigma}^m(\widehat{\mathcal{L}}_\sigma^n h)(x)}{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \mathbf{1}(x)} \geq \frac{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \left[\frac{\widehat{\mathcal{L}}_\sigma^m h}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} \widehat{\mathcal{L}}_\sigma^m \mathbf{1} \right](x)}{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \mathbf{1}(x)} \geq \frac{\widehat{\mathcal{L}}_{\tau^n \sigma}^m(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})(x)}{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \mathbf{1}(x)} \inf \frac{\widehat{\mathcal{L}}_\sigma^m h}{\widehat{\mathcal{L}}_\sigma^m \mathbf{1}}$$

where we have used twice the fact that $\widehat{\mathcal{L}}_{\tau^n \sigma}^m \widehat{\mathcal{L}}_\sigma^n = \widehat{\mathcal{L}}_{\sigma^n \sigma}^m \widehat{\mathcal{L}}_\sigma^m$. Taking the inf on x and the limit $m \rightarrow \infty$, we get the first inequality. For the second, for $x \in [0, 1]$, we have

$$\frac{\widehat{\mathcal{L}}_{\tau^n \sigma}^m(\widehat{\mathcal{L}}_\sigma^n h)(x)}{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \mathbf{1}(x)} = \frac{\widehat{\mathcal{L}}_{\tau^n \sigma}^m(\widehat{\mathcal{L}}_\sigma^n h)(x)}{\widehat{\mathcal{L}}_{\tau^n \sigma}^m(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})(x)} \frac{\widehat{\mathcal{L}}_{\tau^n \sigma}^m(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})(x)}{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \mathbf{1}(x)} \leq \frac{\widehat{\mathcal{L}}_\sigma^{n+m} h(x)}{\widehat{\mathcal{L}}_\sigma^{n+m} \mathbf{1}(x)} \|\widehat{\mathcal{L}}_\sigma^n \mathbf{1}\|_\infty,$$

which, by taking the inf on x and the limit $m \rightarrow \infty$, yields

$$\Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n h) \leq \|\widehat{\mathcal{L}}_\sigma^n \mathbf{1}\|_\infty \Lambda_\sigma(h).$$

By applying Lemma 9.5 to $\mathbf{1} \in \mathcal{C}_\sigma^{a_0}$, we obtain $\bigvee \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \leq 2a_0 C_\star \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})$. Thus

$$\|\widehat{\mathcal{L}}_\sigma^n \mathbf{1}\|_\infty \leq \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1}) + \bigvee \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \leq (1 + 2a_0 C_\star) \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1}) = B \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1}).$$

□

Lemma 9.7. *Let $n_1(\delta) = \lceil \frac{\ln \delta^{-1}}{\ln \theta^{-1} \rho} \rceil$. Then, for each $\delta \in (0, 1)$ and $n \geq n_1(\delta)$, the partition $\widehat{\mathcal{Z}}_\sigma^n$ has the property*

$$\sup_{Z \in \widehat{\mathcal{Z}}_\sigma^n} \Lambda_\sigma(\mathbf{1}_Z) \leq \delta.$$

Proof. Remark that $n_1(\delta)$ is well defined due to condition (C2) and is such that for all $n \geq n_1(\delta)$, $\theta^n \rho^{-n} \leq \delta$. Then, for $Z \in \widehat{\mathcal{Z}}_\sigma^n$,

$$\widehat{\mathcal{L}}_\sigma^n \mathbf{1}_Z(x) = \sum_{T_\sigma^n y=x} g_\sigma^n(y) \mathbf{1}_Z(y) \leq \|g_\sigma^n\|_\infty \leq \theta^n.$$

Accordingly, for each $x \in [0, 1]$,

$$\frac{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \widehat{\mathcal{L}}_\sigma^n \mathbf{1}_Z(x)}{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \widehat{\mathcal{L}}_\sigma^n \mathbf{1}(x)} \leq \frac{\theta^n}{\inf \frac{\widehat{\mathcal{L}}_{\tau^n \sigma}^m(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}{\widehat{\mathcal{L}}_{\tau^n \sigma}^m \mathbf{1}}}.$$

Taking the inf on x and the limit $m \rightarrow \infty$, this yields

$$\Lambda_\sigma(\mathbf{1}_Z) \leq \frac{\theta^n}{\Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})} \leq \theta^n \rho^{-n} \leq \delta$$

where we have used Lemma 9.4. \square

Lemma 9.8. *Let $n_2(a) = n_1([4aB(1 + 2C_{n_0} \rho^{-n_0})]^{-1})$. If (C3(n_0, N')) holds, then for each $a \geq a_0$, $n \geq n_2$ and $h \in \mathcal{C}_\sigma^a$ there exists $Z \in \mathcal{Z}_{\sigma, g}^n$ with*

$$\inf_Z h \geq \frac{1}{4} \Lambda_\sigma(h).$$

Proof. For each n, m with $n < m$, we can write

$$\widehat{\mathcal{L}}_\sigma^m h(x) = \sum_{Z \in \widehat{\mathcal{Z}}_\sigma^n} \widehat{\mathcal{L}}_\sigma^m(h \mathbf{1}_Z)(x) = \sum_{Z \in \mathcal{Z}_{\sigma, *}} \widehat{\mathcal{L}}_\sigma^m(h \mathbf{1}_Z)(x).$$

Suppose the lemma is not true. Then, we have

$$\begin{aligned} \widehat{\mathcal{L}}_\sigma^m h(x) &= \sum_{Z \in \mathcal{Z}_{\sigma, g}^n} \widehat{\mathcal{L}}_\sigma^m(h \mathbf{1}_Z)(x) + \sum_{Z \in \mathcal{Z}_{\sigma, b}^n} \widehat{\mathcal{L}}_\sigma^m(h \mathbf{1}_Z)(x) \\ &\leq \sum_{Z \in \mathcal{Z}_{\sigma, g}^n} \widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z(x) \frac{\Lambda_\sigma(h)}{4} + \sum_{Z \in \mathcal{Z}_{\sigma, g}^n} \widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z(x) \bigvee_Z h + \|h\|_\infty \sum_{Z \in \mathcal{Z}_{\sigma, b}^n} \widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z(x) \\ &\leq \widehat{\mathcal{L}}_\sigma^m \mathbf{1}(x) \frac{\Lambda_\sigma(h)}{4} + \sum_{Z \in \mathcal{Z}_{\sigma, g}^n} \left[\Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z) + \bigvee_Z \widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z \right] \bigvee_Z h \\ &\quad + \|h\|_\infty \sum_{Z \in \mathcal{Z}_{\sigma, b}^n} \widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z(x). \end{aligned}$$

If $Z \in \mathcal{Z}_{\sigma, b}^n$, by Lemma 9.6, we have $\Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z) \leq B \Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_\sigma^m \mathbf{1}) \Lambda_\sigma(\mathbf{1}_Z) = 0$, which implies

$$\widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z(x) \leq \bigvee_Z \widehat{\mathcal{L}}_\sigma^m \mathbf{1}_Z \leq 2C_*^{m/n_0+1} \theta^m \leq 2C_*(C_*^{1/n_0} \theta \rho^{-1})^m \Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_\sigma^m \mathbf{1}),$$

by inequality (9.1) and Lemma 9.4.

If $Z \in \mathcal{Z}_{\sigma,g}^n$, the same argument gives

$$\begin{aligned} \bigvee \widehat{\mathcal{L}}_{\sigma}^m \mathbf{1}_Z &\leq 2C_{\star}^{m/n_0+1} \theta^m + 2C_{n_0} \rho^{-n_0} \Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_{\sigma}^m \mathbf{1}_Z) \\ &\leq \left[2C_{\star} (C_{\star}^{1/n_0} \theta \rho^{-1})^m + 2C_{n_0} \rho^{-n_0} B \Lambda_{\sigma}(\mathbf{1}_Z) \right] \Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_{\sigma}^m \mathbf{1}). \end{aligned}$$

Setting $\kappa = C_{\star}^{1/n_0} \theta \rho^{-1} \leq 4^{-1/n_0}$ and using Lemma 9.6 again, we have

$$\begin{aligned} \Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_{\sigma}^m h) &\leq \frac{\Lambda_{\sigma}(h)}{4} \Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_{\sigma}^m \mathbf{1}) \\ &\quad + \sum_{Z \in \mathcal{Z}_{\sigma,g}^n} \Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_{\sigma}^m \mathbf{1}) \bigvee_Z h \left[B(1 + 2C_{n_0} \rho^{-n_0}) \Lambda_{\sigma}(\mathbf{1}_Z) + 2C_{\star} \kappa^m \right] \\ &\quad + \|h\|_{\infty} \sum_{Z \in \mathcal{Z}_{\sigma,b}^n} 2C_{\star} \kappa^m \Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_{\sigma}^m \mathbf{1}). \end{aligned}$$

Dividing this inequality by $\Lambda_{\tau^m \sigma}(\widehat{\mathcal{L}}_{\sigma}^m \mathbf{1})$ and taking the limit $m \rightarrow \infty$ yields

$$\begin{aligned} \Lambda_{\sigma}(h) &\leq \frac{\Lambda_{\sigma}(h)}{4} + B(1 + 2C_{n_0} \rho^{-n_0}) \bigvee h \sup_{Z \in \mathcal{Z}_{\sigma,g}^n} \Lambda_{\sigma}(\mathbf{1}_Z) \\ &\leq \left[\frac{1}{4} + aB(1 + 2C_{n_0} \rho^{-n_0}) \sup_{Z \in \mathcal{Z}_{\sigma,g}^n} \Lambda_{\sigma}(\mathbf{1}_Z) \right] \Lambda_{\sigma}(h) \leq \frac{1}{2} \Lambda_{\sigma}(h) \end{aligned}$$

where we applied Lemma 9.7. This yields the announced contradiction. \square

Remark 9.9. From now on we set $a = \max\{a_0, 1\}$.

Remark 9.10. Since $\mathcal{T} \times \mathcal{H}$ is finite, there exists $M > 1$ such that $\|\widehat{\mathcal{L}}_{\sigma}^1 \mathbf{1}\|_{\infty} \leq M$. Hence, by condition (C1), $\delta_{\star}^n \leq \inf \widehat{\mathcal{L}}_{\sigma}^n \mathbf{1} \leq \|\widehat{\mathcal{L}}_{\sigma}^n \mathbf{1}\|_{\infty} \leq M^n$ for all $n \geq 0$ and $\sigma \in \Sigma^{\mathbb{N}}$.

Lemma 9.11. If (C3(n_2, n_3)) holds for some $n_3 \geq n_2$, then there exists $\varrho > 0$ such that

$$\inf \widehat{\mathcal{L}}_{\sigma}^n f \geq \varrho \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_{\sigma}^n f),$$

for all $n \geq n_3$, all $\sigma \in \Sigma^{\mathbb{N}}$ and all $f \in C_{\sigma}^a$.

Proof. We first remark that, by Lemma 9.8, for each $f \in C_{\sigma}^a$, there exists $Z \in \mathcal{Z}_{\sigma,g}^{n_2}$ such that

$$\inf_Z f \geq \frac{1}{4} \Lambda_{\sigma}(f).$$

Consequently, for any $\ell \geq 0$, we have

$$\inf \widehat{\mathcal{L}}_{\sigma}^{\ell} f \geq \frac{1}{4} \Lambda_{\sigma}(f) \inf \frac{\widehat{\mathcal{L}}_{\sigma}^{\ell} \mathbf{1}_Z}{\widehat{\mathcal{L}}_{\sigma}^{\ell} \mathbf{1}} \inf \widehat{\mathcal{L}}_{\sigma}^{\ell} \mathbf{1}.$$

Since condition (C3(n_2, n_3)) holds, one has

$$\inf \frac{\widehat{\mathcal{L}}_{\sigma}^{\ell} \mathbf{1}_Z}{\widehat{\mathcal{L}}_{\sigma}^{\ell} \mathbf{1}} \geq \epsilon_{\star}(n_2)$$

for all $\sigma \in \Sigma^{\mathbb{N}}$, $Z \in \mathcal{Z}_{\sigma,g}^{n_2}$ and $\ell \geq n_3$, and we thus get

$$(9.2) \quad \inf \widehat{\mathcal{L}}_{\sigma}^{\ell} f \geq \frac{\epsilon_{\star}(n_2)}{4} \Lambda_{\sigma}(f) \inf \widehat{\mathcal{L}}_{\sigma}^{\ell} \mathbf{1}.$$

We first prove Lemma 9.11 when $n = n_3$ and then extend it to all $n \geq n_3$. By Lemma 9.6,

$$\Lambda_\sigma(f) \geq B^{-1} \frac{\Lambda_{\tau^{n_3}\sigma}(\widehat{\mathcal{L}}_\sigma^{n_3} f)}{\Lambda_{\tau^{n_3}\sigma}(\widehat{\mathcal{L}}_\sigma^{n_3} \mathbf{1})} \geq B^{-1} \frac{\Lambda_{\tau^{n_3}\sigma}(\widehat{\mathcal{L}}_\sigma^{n_3} f)}{\sup \widehat{\mathcal{L}}_\sigma^{n_3} \mathbf{1}}.$$

By Remark 9.10 and (9.2) with $\ell = n_3$, we obtain

$$\inf \widehat{\mathcal{L}}_\sigma^{n_3} f \geq \hat{\varrho} \Lambda_{\tau^{n_3}\sigma}(\widehat{\mathcal{L}}_\sigma^{n_3} f),$$

with $\hat{\varrho} = B^{-1} \frac{\epsilon_*(n_2)}{4} \left(\frac{\delta_*}{M}\right)^{n_3}$.

By Lemma 9.5, $\widehat{\mathcal{L}}_\sigma^{n_0} C_\sigma^a \subset C_{\tau^{n_0}\sigma}^a$ for all σ . We thus have

$$\inf \widehat{\mathcal{L}}_\sigma^{n_3+kn_0} f \geq \hat{\varrho} \Lambda_{\tau^{n_3+kn_0}\sigma}(\widehat{\mathcal{L}}_\sigma^{n_3+kn_0} f),$$

for all $k \geq 0$, $\sigma \in \Sigma^{\mathbb{N}}$ and $f \in C_\sigma^a$.

Let now $n \geq n_3$. We write $n = kn_0 + n_3 + r = n' + r$ with $r < n_0$. We have

$$\begin{aligned} \inf \widehat{\mathcal{L}}_\sigma^n f &= \widehat{\mathcal{L}}_{\tau^{n'}\sigma}^r \widehat{\mathcal{L}}_\sigma^{n'} f \geq \hat{\varrho} \Lambda_{\tau^{n'}\sigma}(\widehat{\mathcal{L}}_\sigma^{n'} f) \inf \widehat{\mathcal{L}}_{\tau^{n'}\sigma}^r \mathbf{1} \\ &\geq \hat{\varrho} \delta_*^r \Lambda_{\tau^{n'}\sigma}(\widehat{\mathcal{L}}_\sigma^{n'} f) \\ &\geq \hat{\varrho} \delta_*^{n_0} \Lambda_{\tau^{n'}\sigma}(\widehat{\mathcal{L}}_\sigma^{n'} f). \end{aligned}$$

But,

$$\begin{aligned} \Lambda_{\tau^{n'}\sigma}(\widehat{\mathcal{L}}_\sigma^{n'} f) &= \liminf_{k \rightarrow \infty} \frac{\widehat{\mathcal{L}}_{\tau^{n'+r}\sigma}^k \widehat{\mathcal{L}}_{\tau^{n'}\sigma}^r \widehat{\mathcal{L}}_\sigma^{n'} f}{\widehat{\mathcal{L}}_{\tau^{n'+r}\sigma}^k \widehat{\mathcal{L}}_{\tau^{n'}\sigma}^r \mathbf{1}} \\ &\geq M^{-r} \liminf_{k \rightarrow \infty} \frac{\widehat{\mathcal{L}}_{\tau^{n'+r}\sigma}^k \widehat{\mathcal{L}}_\sigma^{n'+r} f}{\widehat{\mathcal{L}}_{\tau^{n'+r}\sigma}^k \mathbf{1}} \\ &= M^{-r} \Lambda_{\tau^{n'+r}\sigma}(\widehat{\mathcal{L}}_\sigma^{n'+r} f) \\ &\geq M^{-n_0} \Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f). \end{aligned}$$

We have thus proved the result with $\varrho = \hat{\varrho} \left(\frac{\delta_*}{M}\right)^{n_0}$. \square

Corollary 9.12. *It holds true $\rho \leq 1$, where ρ is defined in (6.2).²⁴*

Proof. By Lemma 9.4 and Lemma 9.11, with $f = 1$, we see that, for all $n \geq n_3$, $\inf \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \geq \varrho \rho^n$ but $\inf \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \leq \int \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \leq 1$, so $\rho \leq 1$. \square

Remark 9.13. *Since by Remark 9.10, $\|\widehat{\mathcal{L}}_\sigma^n \mathbf{1}\|_\infty \leq M^n$ and $\inf \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \geq \delta_*^n$ for all $n \geq 0$ and $\sigma \in \Sigma^{\mathbb{N}}$, we have*

$$\inf \widehat{\mathcal{L}}_\sigma^n \mathbf{1} \geq \varrho' \Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1}),$$

with $\varrho' = \min\{\varrho, 1, \frac{\delta_*}{M}, \dots, \left(\frac{\delta_*}{M}\right)^{n_3-1}\}$.

Lemma 9.14. *If (C3(n_2, n_3)) holds for some $n_3 \geq n_2$, then for all $\sigma \in \Sigma^{\mathbb{N}}$ and $n \geq n_3$, one has*

$$\widehat{\mathcal{L}}_\sigma^n C_\sigma^a \subset C_{\tau^n\sigma}^a$$

with finite diameter less than Δ_n , uniformly in σ .

²⁴ Hence the min in condition (C2) was, a posteriori, superfluous.

Proof. By (9.2), we have

$$\inf \widehat{\mathcal{L}}_\sigma^n h \geq \frac{\epsilon_*(n_2)}{4} \Lambda_\sigma(h) \inf \widehat{\mathcal{L}}_\sigma^n \mathbf{1},$$

for any $\sigma \in \Sigma^\mathbb{N}$, $h \in C_\sigma^a$ and $n \geq n_3$ and, using Lemmata 9.5 and 9.6,

$$\sup \widehat{\mathcal{L}}_\sigma^n h \leq \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n h) + \bigvee \widehat{\mathcal{L}}_\sigma^n h \leq \left(1 + \frac{a}{2}\right) B \Lambda_\sigma(h) \Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1}).$$

A simple adaptation of the proof of [16, Lemma 3.1] yields

$$d_{C_\sigma^a}(g, \mathbf{1}) \leq \log \left[\frac{\max \{(1 + \nu) \Lambda_\sigma(g), \sup g\}}{\min \{(1 - \nu) \Lambda_\sigma(g), \inf g\}} \right]$$

for any $g \in C_\sigma^{\nu a}$ with $0 < \nu < 1$. By Lemma 9.5 $\widehat{\mathcal{L}}_\sigma^n(C_\sigma^a) \subset C_{\tau^n \sigma}^{a/2}$, so we can choose $\nu = \frac{1}{2}$ and, recalling Remark 9.10, we obtain:

$$\text{diam}_{C_{\tau^n \sigma}^a} \widehat{\mathcal{L}}_\sigma^n(C_\sigma^a) \leq 2 \log \left[\frac{\max \left\{ \frac{3}{2}, BM^n \left(1 + \frac{a}{2}\right) \right\}}{\min \left\{ \frac{1}{2}, \frac{\epsilon_*(n_2) \delta_*^n}{4} \right\}} \right] =: \Delta_n < \infty$$

for any $\sigma \in \Sigma^\mathbb{N}$. □

Since we are interested in functions of the form $\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^n f$, we will need to consider functions f that belong to the intersections of all the cones C_σ^a , $\sigma \in \Sigma^\mathbb{Z}$. For this purpose, we introduce the family of cones

$$C_{\text{inf}}^a = \{f \in \text{BV} : f \neq 0, f \geq 0, \bigvee f \leq a \inf f\}.$$

We have $C_{\text{inf}}^a \subset C_\sigma^a$ for any $\sigma \in \Sigma^\mathbb{Z}$, and thus $d_{C_\sigma^a} \leq d_{C_{\text{inf}}^a}$ by Theorem 9.1.

Lemma 9.15. *There exist $\nu \in (0, 1)$ and a family of positive functions $\{h_\sigma\}_{\sigma \in \Sigma^\mathbb{Z}}$ in BV such that for all $f \in C_{\text{inf}}^a$, $\sigma \in \Sigma^\mathbb{Z}$ and $n \geq 0$:*

$$\left\| \frac{\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^n f)} - h_\sigma \right\|_\infty \leq C_\# \nu^n,$$

and

$$\left\| \frac{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^n f)}{\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^n f} - h_\sigma^{-1} \right\|_\infty \leq C_\# \nu^n.$$

Furthermore, $h_\sigma \in C_\sigma^{a/2}$, $\|h_\sigma\|_\infty \leq 1 + a$ and $\inf h_\sigma \geq \varrho > 0$ for all $\sigma \in \Sigma^\mathbb{Z}$, where ϱ is defined in Lemma 9.11.

Proof. Writing $n \geq 2n_3$ as $n = kn_3 + r$, with $k \geq 2$ and $r < n_3$, for all $f \in C_{\text{inf}}^a$, $\sigma \in \Sigma^\mathbb{Z}$ and $m \geq 0$, we have by Theorem 9.1 and Lemma 9.14:

$$d_{C_\sigma^a}(\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^n f, \widehat{\mathcal{L}}_{\tau^{-(n+m)} \sigma}^{n+m} f) \leq \gamma^{k-2} d_{C_{\tau^{-(k-2)n_3} \sigma}^a}(\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^{2n_3+r} f, \widehat{\mathcal{L}}_{\tau^{-(n+m)} \sigma}^{2n_3+r+m} f),$$

with $\gamma = \tanh\left(\frac{\Delta_{n_3}}{4}\right) < 1$.

Since both $\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^{n_3+r} f$ and $\widehat{\mathcal{L}}_{\tau^{-(n+m)} \sigma}^{n_3+r+m} f$ belong to $C_{\tau^{-(k-1)n_3} \sigma}^a$ by Lemma 9.14 again, as $f \in C_{\text{inf}}^a \subset C_{\tau^{-n} \sigma}^a \cap C_{\tau^{-(n+m)} \sigma}^a$, we have, using Lemma 9.14 one more time:

$$d_{C_{\tau^{-(k-2)n_3} \sigma}^a}(\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^{2n_3+r} f, \widehat{\mathcal{L}}_{\tau^{-(n+m)} \sigma}^{2n_3+r+m} f) \leq \Delta_{n_3}.$$

Consequently, for all $n \geq 2n_3$, $m \geq 0$, $\sigma \in \Sigma^\mathbb{Z}$ and $f \in C_{\text{inf}}^a$,

$$d_{C_\sigma^a}(\widehat{\mathcal{L}}_{\tau^{-n} \sigma}^n f, \widehat{\mathcal{L}}_{\tau^{-(n+m)} \sigma}^{n+m} f) \leq C_\# \nu^n,$$

with $\nu = \gamma^{\frac{1}{n_3}}$.

Using Lemma 9.2 with $\|\cdot\| = \|\cdot\|_\infty$ and $\ell(\cdot) = \Lambda_\sigma(\cdot)$, we get

$$\left\| \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)} - \frac{\widehat{\mathcal{L}}_{\tau-(n+m)\sigma}^{n+m} f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-(n+m)\sigma}^{n+m} f)} \right\|_\infty \leq (e^{d_{C_\sigma^a}(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f, \widehat{\mathcal{L}}_{\tau-(n+m)\sigma}^{n+m} f)} - 1) \left\| \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)} \right\|_\infty.$$

Since $\widehat{\mathcal{L}}_{\tau-n\sigma}^n f \in C_\sigma^a$, we have

$$(9.3) \quad \|\widehat{\mathcal{L}}_{\tau-n\sigma}^n f\|_\infty \leq \Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f) + \bigvee \widehat{\mathcal{L}}_{\tau-n\sigma}^n f \leq (1+a)\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f),$$

and we deduce that

$$\left\| \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)} - \frac{\widehat{\mathcal{L}}_{\tau-(n+m)\sigma}^{n+m} f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-(n+m)\sigma}^{n+m} f)} \right\|_\infty \leq C_\# \nu^n.$$

This implies that $\frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)}$ is a Cauchy sequence in L^∞ , and thus converges to a function $h_\sigma \in L^\infty$. Since $C_\sigma^{a/2}$ is closed in L^∞ and $\widehat{\mathcal{L}}_{\tau-n\sigma}^n f \in C_\sigma^{a/2}$ for $n \geq n_0$ by Lemma 9.5, we have $h_\sigma \in C_\sigma^{a/2}$. Passing to the limit $m \rightarrow \infty$ in the previous relation, we obtain

$$\left\| \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)} - h_\sigma \right\|_\infty \leq C_\# \nu^n,$$

for all $n \geq 2n_3$, and all $\sigma \in \Sigma^{\mathbb{Z}}$. Using the same reasoning, we have for any pair $f, f' \in C_{\tau-n\sigma}^a$,

$$(9.4) \quad \left\| \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)} - \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f'}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f')} \right\|_\infty \leq C_\# \nu^n,$$

which proves that the limit h_σ does not depend on the choice of $f \in C_{\text{inf}}^a$. By Lemma 9.11, $\widehat{\mathcal{L}}_{\tau-n\sigma}^n f \geq \varrho \Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)$ for all $n \geq n_3$, whence we obtain $\inf h_\sigma \geq \varrho$. Remark that (9.3) implies $\|h_\sigma\|_\infty \leq 1+a$. When $n < 2n_3$, we have

$$\left\| \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)} - h_\sigma \right\|_\infty \leq \frac{M^{2n_3}}{\delta_\star^{2n_3}} \sup f + (1+a) \leq (1+a) \left(\frac{M^{2n_3}}{\delta_\star^{2n_3}} + 1 \right) \leq C_\# \nu^n,$$

where $M > 1$ and $0 < \delta_\star < 1$ are defined in Remark 9.10. For $n \geq n_3$ and $f \in C_{\text{inf}}^a$, since $\inf h_\sigma \geq \varrho$ and $\inf \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)} \geq \varrho$ by Lemma 9.11, we have

$$\left\| \frac{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)}{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f} - h_\sigma^{-1} \right\|_\infty \leq \varrho^{-2} \left\| \frac{\widehat{\mathcal{L}}_{\tau-n\sigma}^n f}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n f)} - h_\sigma \right\|_\infty \leq C_\# \nu^n.$$

We handle the case $n < n_3$ as previously, since $\|h_\sigma\|_\infty \leq 1+a$. \square

Lemma 9.16. *For all $\sigma \in \Sigma^{\mathbb{Z}}$, there exists $\lambda_\sigma \geq \rho_\sigma$ such that $\widehat{\mathcal{L}}_\sigma^n h_\sigma = \lambda_\sigma \cdots \lambda_{\tau^{n-1}\sigma} h_{\tau^n\sigma}$ for all $n \geq 1$.*

Proof. Applying Lemma 9.6 with $h = \widehat{\mathcal{L}}_{\tau-n\sigma}^n \mathbf{1}$, we have by definition of $\rho_\sigma = \Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_\sigma^1 \mathbf{1})$:

$$\rho_\sigma \leq \frac{\Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_{\tau-n\sigma}^{n+1} \mathbf{1})}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau-n\sigma}^n \mathbf{1})} \leq B\rho_\sigma.$$

Consequently, there exist a subsequence $\{n_j\}$ and $\lambda_\sigma \in [\rho_\sigma, B\rho_\sigma]$ such that

$$\lambda_\sigma = \lim_j \frac{\Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j+1} \mathbf{1})}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j} \mathbf{1})}.$$

We can now compute

$$\begin{aligned} \widehat{\mathcal{L}}_\sigma^1 h_\sigma &= \lim_j \widehat{\mathcal{L}}_\sigma^1 \frac{\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j} \mathbf{1}}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j} \mathbf{1})} = \lim_j \frac{\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j+1} \mathbf{1}}{\Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j+1} \mathbf{1})} \frac{\Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j+1} \mathbf{1})}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j} \mathbf{1})} \\ &= \lim_j \frac{\widehat{\mathcal{L}}_{\tau^{-(n_j+1)}\sigma}^{n_j+1} \mathbf{1}}{\Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_{\tau^{-(n_j+1)}\sigma}^{n_j+1} \mathbf{1})} \frac{\Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j+1} \mathbf{1})}{\Lambda_\sigma(\widehat{\mathcal{L}}_{\tau^{-n_j}\sigma}^{n_j} \mathbf{1})} \\ &= \lambda_\sigma h_{\tau\sigma}. \end{aligned}$$

The general case $n \geq 1$ is obtained by a simple induction. \square

Lemma 9.17. *For all $\sigma \in \Sigma^{\mathbb{N}}$, the functional Λ_σ (restricted to BV) is linear, positive, and enjoys the property $\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f) = \rho_\sigma \cdots \rho_{\tau^{n-1}\sigma} \Lambda_\sigma(f)$ for all $f \in \text{BV}$ and $n \geq 1$. Moreover, $\lambda_\sigma = \rho_\sigma$ and $\left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - \Lambda_\sigma(f) \right\|_\infty \leq C_\# \nu^n \|f\|_{\text{BV}}$ for all $f \in \text{BV}$.*

Proof. For $f \in C_{\text{inf}}^a$, we can write

$$\frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} = \frac{\widehat{\mathcal{L}}_\sigma^n f}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)} \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})} \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}}.$$

So

$$\begin{aligned} \left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})} \right\|_\infty &= \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})} \left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)} \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - 1 \right\|_\infty \\ &\leq \|f\|_\infty \left(\left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)} - h_\sigma \right\|_\infty \left\| \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} \right\|_\infty + \|h_\sigma\|_\infty \left\| \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - h_\sigma^{-1} \right\|_\infty \right). \end{aligned}$$

Since $\left\| \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} \right\|_\infty \leq \varrho^{-1}$ for $n \geq n_3$ by Lemma 9.11, we get, using Lemma 9.15, for all $f \in C_{\text{inf}}^a$ and $n \geq n_3$:

$$(9.5) \quad \left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})} \right\|_\infty \leq C_\# \nu^n \|f\|_\infty.$$

But, $\Lambda_\sigma(f) = \lim_{n \rightarrow \infty} \inf \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}}$ by definition, and, since $\frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})}$ are constants, we deduce that $\lim_{n \rightarrow \infty} \frac{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n f)}{\Lambda_{\tau^n\sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})} = \Lambda_\sigma(f)$ and

$$\lim_{n \rightarrow \infty} \left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - \Lambda_\sigma(f) \right\|_\infty = 0.$$

Now, if $f \in \text{BV}$, we have $f + c \in C_{\text{inf}}^a$ for $c = (1 + a^{-1})\|f\|_{\text{BV}}$, so we get that $\Lambda_\sigma(f) = \lim_{n \rightarrow \infty} \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}}$ in L^∞ for all $f \in \text{BV}$, since $\Lambda_\sigma(f + c) = \Lambda_\sigma(f) + c$. The linearity of Λ follows from the linearity of the limit.

Next, as $\widehat{\mathcal{L}}_\sigma^1 f \in \text{BV}$, we know that

$$\Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_\sigma^1 f) = \lim_{n \rightarrow \infty} \frac{\widehat{\mathcal{L}}_\sigma^{n+1} f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} = \lim_{n \rightarrow \infty} \frac{\widehat{\mathcal{L}}_\sigma^{n+1} f \widehat{\mathcal{L}}_\sigma^{n+1} \mathbf{1}}{\widehat{\mathcal{L}}_\sigma^{n+1} \mathbf{1} \widehat{\mathcal{L}}_\sigma^n \mathbf{1}} = \Lambda_\sigma(f) \Lambda_{\tau\sigma}(\widehat{\mathcal{L}}_\sigma^1 \mathbf{1}) = \rho_\sigma \Lambda_\sigma(f).$$

But then $\rho_\sigma = \lambda_\sigma$ is obtained by taking $f = h_\sigma$, since $\Lambda_\sigma(h_\sigma) = 1$. In particular, we have $\Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n f) = \rho_\sigma \cdots \rho_{\tau^{n-1} \sigma} \Lambda_\sigma(f)$ for all $f \in \text{BV}$, so $\frac{\Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n f)}{\Lambda_{\tau^n \sigma}(\widehat{\mathcal{L}}_\sigma^n \mathbf{1})} = \Lambda_\sigma(f)$. But if we look back to (9.5), the above implies that for all $f \in C_{\text{inf}}^a$ and $n \geq n_3$:

$$\left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - \Lambda_\sigma(f) \right\|_\infty \leq C_\# \nu^n \|f\|_\infty.$$

This can be easily extended to all $n \geq 0$, since $\left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - \Lambda_\sigma(f) \right\|_\infty \leq 2\|f\|_\infty$. We can again cover the general case $f \in \text{BV}$ using the fact that $\widehat{f} + c \in C_{\text{inf}}^a$ for $c = (1 + a^{-1})\|f\|_{\text{BV}}$, which finally implies

$$\left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\widehat{\mathcal{L}}_\sigma^n \mathbf{1}} - \Lambda_\sigma(f) \right\|_\infty \leq C_\# \nu^n \|f + c\|_\infty \leq C_\# \nu^n \|f\|_{\text{BV}}.$$

□

Remark 9.18. *Following closely the ideas of [17], it is possible to prove that Λ_σ can be interpreted as a non-atomic measure μ_σ , i.e. $\Lambda_\sigma(f) = \int f d\mu_\sigma$ for all $f \in \text{BV}$, and that the measure ν_σ defined by $d\nu_\sigma = h_\sigma d\mu_\sigma$ satisfies $(T_\sigma^1)_* \nu_\sigma = \nu_{\tau\sigma}$. Since we will not make use of these facts, we leave their proofs to the interested reader.*

The main properties of Λ_σ being proved, we can now improve Lemma 9.15 by extending it to general functions in BV and deduce Proposition 8.1:

Proof of Proposition 8.1. By (9.4) with $f' = h_\sigma$, for any $f \in C_\sigma^a$, we get using Lemma 9.17

$$\left\| \frac{\widehat{\mathcal{L}}_\sigma^n f}{\rho_\sigma \cdots \rho_{\tau^{n-1} \sigma}} - \Lambda_\sigma(f) h_{\tau^n \sigma} \right\|_\infty \leq C_\# \nu^n \Lambda_\sigma(f) \leq C_\# \nu^n \|f\|_\infty.$$

Now, if $f \in \text{BV}$, we have $f + ch_\sigma \in C_\sigma^a$ for all $\sigma \in \Sigma^{\mathbb{Z}}$ with $c = 2(1 + a^{-1})\|f\|_{\text{BV}}$. Indeed, since $\bigvee h_\sigma \leq \frac{a}{2} \Lambda_\sigma(h_\sigma) = \frac{a}{2}$ by Lemma 9.15, we have

$$\bigvee (f + ch_\sigma) \leq \bigvee f + c \bigvee h_\sigma \leq \bigvee f + \frac{ac}{2},$$

and

$$\Lambda_\sigma(f + ch_\sigma) = \Lambda_\sigma(f) + c \geq \inf f + c.$$

So $f + ch_\sigma \in C_\sigma^a$ if $c \geq 2(a^{-1} \bigvee f - \inf f)$, which is the case for our particular choice of c . Consequently, we have

$$\left\| \frac{\widehat{\mathcal{L}}_\sigma^n (f + ch_\sigma)}{\rho_\sigma \cdots \rho_{\tau^{n-1} \sigma}} - \Lambda_\sigma(f + ch_\sigma) h_{\tau^n \sigma} \right\|_\infty \leq C_\# \nu^n \|f + ch_\sigma\|_\infty \leq C_\# \nu^n \|f\|_{\text{BV}},$$

which leads to (8.1) after simplifications, since

$$\widehat{\mathcal{L}}_\sigma^n (f + ch_\sigma) = \widehat{\mathcal{L}}_\sigma^n f + c \rho_\sigma \cdots \rho_{\tau^{n-1} \sigma} h_{\tau^n \sigma}$$

and $\Lambda_\sigma(f + ch_\sigma) = \Lambda_\sigma(f) + c$. □

Proof of Lemma 8.3. By Remark 9.10, we have $\rho_\sigma \leq \|\widehat{\mathcal{L}}_\sigma^1 \mathbf{1}\|_\infty \leq M$, and so $\sup_\sigma \rho_\sigma \leq M < \infty$. The remaining statements are immediate consequences of Lemmata 9.15, 9.16 and 9.17. \square

Proof of Lemma 8.4. This follows immediately from Lemma 9.6 and Remark 9.13. \square

REFERENCES

- [1] G. Atkinson. Recurrence of co-cycles and random walks. *Journal of the London Mathematical Society*, s2-13(3):486–488, 1976.
- [2] L. A. Bunimovich, Y. G. Sinaĭ, and N. I. Chernov. Statistical properties of two-dimensional hyperbolic billiards. *Uspekhi Mat. Nauk*, 46(4(280)):43–92, 192, 1991.
- [3] J.-P. Conze. Sur un critère de récurrence en dimension 2 pour les marches stationnaires, applications. *Ergodic Theory Dynam. Systems*, 19(5):1233–1245, 1999.
- [4] G. Cristadoro, M. Lenci, and M. Seri. Erratum: “Recurrence for quenched random Lorentz tubes” [Chaos 20, 023115 (2010)] [mr2741899]. *Chaos*, 20(4):049903, 1, 2010.
- [5] G. Cristadoro, M. Lenci, and M. Seri. Recurrence for quenched random Lorentz tubes. *Chaos*, 20(2):023115, 7, 2010.
- [6] J. De Simoi and C. Liverani. The martingale approach after Varadhan and Dolgopyat. In *Hyperbolic dynamics, fluctuations and large deviations*, volume 89 of *Proc. Sympos. Pure Math.*, pages 311–339. Amer. Math. Soc., Providence, RI, 2015.
- [7] D. Dolgopyat, D. Szász, and T. Varjú. Recurrence properties of planar Lorentz process. *Duke Math. J.*, 142(2):241–281, 2008.
- [8] D. Dolgopyat, D. Szász, and T. Varjú. Limit theorems for locally perturbed planar Lorentz processes. *Duke Math. J.*, 148(3):459–499, 2009.
- [9] N. Dunford, J. Schwartz, W. Bade, and R. Bartle. *Linear Operators: General theory. Part. I*. Linear Operators. Interscience Publishers, 1988.
- [10] M. Lenci. Aperiodic Lorentz gas: recurrence and ergodicity. *Ergodic Theory Dynam. Systems*, 23(3):869–883, 2003.
- [11] M. Lenci. Typicality of recurrence for Lorentz gases. *Ergodic Theory Dynam. Systems*, 26(3):799–820, 2006.
- [12] M. Lenci. Central limit theorem and recurrence for random walks in bistochastic random environments. *Journal of Mathematical Physics*, 49(12):125213, 2008.
- [13] M. Lenci and S. Troubetzkoy. Infinite-horizon Lorentz tubes and gases: recurrence and ergodic properties. *Phys. D*, 240(19):1510–1515, 2011.
- [14] C. Little. Deterministically driven random walks in a random environment on \mathbb{Z} . *Discrete and Continuous Dynamical Systems - A*, 36:5555–5578, 2016.
- [15] C. Liverani. Decay of correlations. *Ann. of Math. (2)*, 142(2):239–301, 1995.
- [16] C. Liverani. Decay of correlations for piecewise expanding maps. *Journal of Statistical Physics*, 78(3):1111–1129, 1995.
- [17] C. Liverani and V. Maume-Deschamps. Lasota-Yorke maps with holes: conditionally invariant probability measures and invariant probability measures on the survivor set. *Ann. Inst. H. Poincaré Probab. Statist.*, 39(3):385–412, 2003.
- [18] C. Liverani, B. Saussol, and S. Vaienti. Conformal measure and decay of correlation for covering weighted systems. *Ergodic Theory Dynam. Systems*, 18(6):1399–1420, 1998.
- [19] J. Marklof. The low-density limit of the Lorentz gas: periodic, aperiodic and random. In *Proceedings of the ICM 2014, Seoul, Vol. III*, pages 623–646. 2004.
- [20] J. Marklof and A. Strömbergsson. The Boltzmann-Grad limit of the periodic Lorentz gas. *Ann. of Math. (2)*, 174(1):225–298, 2011.
- [21] A. Mohapatra and W. Ott. Memory loss for nonequilibrium open dynamical systems. *Discrete Contin. Dyn. Syst.*, 34(9):3747–3759, 2014.
- [22] W. Rudin. *Real and complex analysis*. Mathematics series. McGraw-Hill, 1987.
- [23] K. Schmidt. On joint recurrence. *C. R. Acad. Sci. Paris Sér. I Math.*, 327(9):837–842, 1998.
- [24] K. Schmidt. Recurrence of cocycles and stationary random walks. In *Dynamics & stochasticity*, volume 48 of *IMS Lecture Notes Monogr. Ser.*, pages 78–84. Inst. Math. Statist., Beachwood, OH, 2006.

- [25] M. Seri, M. Lenci, M. degli Esposti, and G. Cristadoro. Recurrence and higher ergodic properties for quenched random Lorentz tubes in dimension bigger than two. *J. Stat. Phys.*, 144(1):124–138, 2011.
- [26] T. Simula and M. Stenlund. Deterministic walks in quenched random environments of chaotic maps. *Journal of Physics A: Mathematical and Theoretical*, 42(24):245101, 2009.
- [27] O. Zeitouni. Random walks in random environment. In *Lectures on probability theory and statistics*, volume 1837 of *Lecture Notes in Math.*, pages 189–312. Springer, Berlin, 2004.

ROMAIN AIMINO, DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE, 687, 4169-007 PORTO, PORTUGAL.

E-mail address: `romain.aimino@fc.up.pt`

CARLANGELLO LIVERANI, DIPARTIMENTO DI MATEMATICA, II UNIVERSITÀ DI ROMA (TOR VERGATA), VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA, ITALY.

E-mail address: `liverani@mat.uniroma2.it`