

# PRINCIPAL BUNDLES ON 2-DIMENSIONAL CW-COMPLEXES WITH DISCONNECTED STRUCTURE GROUP

ANDRÉ OLIVEIRA

**ABSTRACT.** Given any topological group  $G$ , the topological classification of  $G$ -principal bundles over a finite CW-complex  $X$  is long-known to be given by the set of free homotopy classes of maps from  $X$  to the corresponding classifying space  $BG$ . This classical result has been long-used to provide such classification in terms of explicit characteristic classes. However, even when  $X$  has dimension 2, it seems there is a case in which such explicit classification has not been explicitly considered. This is the case where  $G$  is a Lie group, whose group of components acts non-trivially on its fundamental group  $\pi_1 G$ . In this note we deal with this case by obtaining the classification, in terms of characteristic classes, of  $G$ -principal bundles over a finite CW-complex of dimension 2, with  $G$  is a Lie group such that  $\pi_0 G$  is abelian.

## 1. INTRODUCTION

It is a classical fact that, for a path-connected CW-complex  $X$  and for a topological group  $G$ , the set of topological types of principal  $G$ -bundles over  $X$  is in bijection with the set  $[X, BG]$  of free homotopy classes of maps from  $X$  to the classifying space  $BG$  of  $G$ . It is, however, many times useful to have a more explicit classification of such bundles, that is, to have a more detailed description of the set  $[X, BG]$ , for example in terms of characteristic classes.

In this note we provide such a description in a setting which is quite classical, and hence which ought to be very well-known. We obtain an explicit and complete classification of  $G$ -bundles over a 2-dimensional connected finite CW-complex  $X$ , for any Lie group  $G$  with  $\pi_0 G$  abelian<sup>1</sup>, in terms of cohomology classes. This includes two classical cases. Firstly, if  $G$  is connected, such bundles are classified by the cohomology group  $H^2(X, \pi_1 G)$ . Secondly, there is a natural action of  $\pi_0 G$  on  $\pi_1 G$ , and if this action is trivial, then  $G$ -bundles over  $X$  are classified by the product  $H^1(X, \pi_0 G) \times H^2(X, \pi_1 G)$ . Nonetheless, such  $\pi_0 G$ -action can be non-trivial and, somehow surprisingly, this case does not seem to have been explicitly treated so far. An instance where this phenomenon occurs is the case of the projective orthogonal group of even rank, which we explicitly deal with as an example.

The importance of the topological classification of  $G$ -principal bundles over any finite, 2-dimensional, CW-complex by itself or by its connections with other areas of Mathematics and Physics is obvious, so we believe that the mentioned untreated case deserves to be considered. This is the purpose of this article. Just to give an example where such topological classification plays an important role, consider the following. Suppose  $G$  is semisimple and consider the space  $\mathcal{R}(X, G) = \text{Hom}^{red}(\pi_1 X, G)/G$  of reductive representations  $\rho: \pi_1 X \rightarrow G$  modulo the  $G$ -action by conjugation. These spaces, or spaces deeply related to them, arise from several different natural points of view (Physics, Hyperbolic Geometry, etc.) and have been intensively studied in the past decades, mostly in the case  $X$  is a closed oriented surface, where deep connections arise with the theory of holomorphic vector bundles [13] and also of Higgs bundles under the so-called non-abelian Hodge theory (there are many references for this subject: the seminal paper is [8] and an overview and other references may be found for instance in [21]). However,

---

*Date:* December 7, 2020.

*2000 Mathematics Subject Classification.* Primary 14H60, 14H40; Secondary 14D20.

The author is grateful to Gustavo Granja (Instituto Superior Técnico, Lisboa) for his generous support and for introducing him to these subjects. Author partially supported by CMUP (UIDB/00144/2020) with national funds.

<sup>1</sup>The abelian condition for  $\pi_0 G$ , besides being naturally verified for most Lie groups, also makes several of the technical arguments much easier to handle with.

the spaces  $\mathcal{R}(X, G)$  have also been studied for compact, non-orientable surfaces [9, 1, 2, 17] and even for any finite 2-dimensional CW-complex [3]. Now, the topological type of  $G$ -bundles over  $X$  provides a way to distinguish certain connected components of  $\mathcal{R}(X, G)$  and to give a lower bound on the number of such components (for example, it is known that if  $G$  is a complex Lie group, such bound is exact [5]). Hence, the theorem of this article gives a way to achieve this for any Lie group  $G$  with  $\pi_0 G$  abelian.

To give a hint on the classification theorem, note first that if a  $G$ -bundle over  $X$  is given by the homotopy class of a map  $X \rightarrow BG$ , then the homotopy class of composite  $X \rightarrow BG \rightarrow B\pi_0 G$  defines a topological invariant  $\mu_1 \in [X, B\pi_0 G] \cong H^1(X, \pi_0 G)$  of the given  $G$ -bundle, measuring the obstruction to reduce its structure group to the identity component of  $G$ . Having this, the classification result (see Theorem 2.2) states that there is a bijection between the subset of  $[X, BG]$  consisting of isomorphism classes of principal  $G$ -bundles over  $X$  with invariant  $\mu_1 \in H^1(X, \pi_0 G)$  and the quotient set  $H^2(X, \pi_1 \mathcal{G}_{\mu_1})/\pi_0 G$ . Here  $\pi_1 \mathcal{G}_{\mu_1}$  denotes the local system obtained from  $\mu_1$  and from a natural action of  $\pi_0 G$  in  $\pi_1 G$  and where  $\pi_0 G$  acts on  $H^2(X, \pi_1 \mathcal{G}_{\mu_1})$  also via the same action of  $\pi_0 G$  in  $\pi_1 G$ .

## 2. THE CLASSIFICATION AND EXAMPLES

**2.1. A first topological invariant.** Let  $X$  be a path-connected 2-dimensional CW-complex. Let also  $G$  a Lie group, with (discrete) group of connected components  $\pi_0 G$ . Recall that isomorphism classes  $G$ -principal bundles over  $X$  are topologically classified by the set  $[X, BG]$  of free homotopy classes of maps from  $X$  to the classifying space  $BG$ . There is a natural map

$$(2.1) \quad \chi : [X, BG] \rightarrow [X, B\pi_0 G], \quad \chi([f]) = [p_{0,*} \circ f],$$

where  $p_{0,*} : BG \rightarrow B\pi_0 G$  is the map induced from the canonical projection  $p_0 : G \rightarrow \pi_0 G$ .

From here one readily defines a first topological invariant of a  $G$ -bundle  $E$  over  $X$ .

**Definition 2.1.** Let  $E$  be a  $G$ -bundle over  $X$  represented by a map  $f : X \rightarrow BG$ . Define

$$\mu_1(E) = \chi([f]) \in [X, B\pi_0 G].$$

Since  $B\pi_0 G$  is the Eilenberg-MacLane space  $K(\pi_0 G, 1)$ , it follows that  $[X, B\pi_0 G] \cong H^1(X, \pi_0 G)$ .  $\mu_1(E)$  is obviously a topological invariant of  $E$  and it represents the isomorphism class of the  $\pi_0 G$ -bundle on  $X$  obtained from  $E$  through the projection  $G \rightarrow \pi_0 G$ .

From now on we fix an element  $\mu_1 \in [X, B\pi_0 G]$  and our task is to classify  $G$ -bundles  $E$  on  $X$  such that  $\mu_1(E) = \mu_1$ .

**2.2. The classification theorem and examples.** Notice that the action

$$\Psi : G \longrightarrow \text{Aut}(G)$$

of  $G$  on itself by conjugation induces an action of  $\pi_0 G$  on the higher homotopy groups  $\pi_i G$  (by  $\pi_i G$  we always mean homotopy groups based at the identity of  $G$ ): for  $a \in G$ , consider the induced automorphism  $\Psi(a)_* : \pi_i G \rightarrow \pi_i G$  and, if  $b \in G$  is in the same component as  $a$ , then  $\Psi(a)$  is homotopic to  $\Psi(b)$  via a path joining  $a$  and  $b$ , hence  $\Psi(a)_* = \Psi(b)_*$ .

In particular, we are interested in the action of  $\pi_0 G$  on  $\pi_1 G$ , which we will denote by

$$(2.2) \quad \Psi(-)_* : \pi_0 G \longrightarrow \text{Aut}(\pi_1 G).$$

Assume now, once and for all, that the group  $\pi_0 G$  is abelian and consider the homomorphism in  $\pi_1$ ,  $\mu_{1*} : \pi_1 X \rightarrow \pi_0 G$ , induced from the invariant  $\mu_1 : X \rightarrow B\pi_0 G$  that we have previously fixed. Given this, we have an induced action of  $\pi_1 X$  in  $\pi_1 G$

$$(2.3) \quad \Psi(-)_* \circ \mu_{1*} : \pi_1 X \longrightarrow \text{Aut}(\pi_1 G)$$

and so this action defines a local system  $\pi_1 \mathcal{G}_{\mu_1}$  on  $X$ .

Let  $H^*(X, \pi_1 \mathcal{G}_{\mu_1})$  be the cohomology of  $X$  with values in the local system  $\pi_1 \mathcal{G}_{\mu_1}$ . Recall that this is the cohomology of the cochain complex  $C_{\mathbb{Z}\pi}^*(\tilde{X}, \pi_1 G) = \text{Hom}_{\mathbb{Z}\pi}(C_*(\tilde{X}), \pi_1 G)$ , where  $\mathbb{Z}\pi = \mathbb{Z}[\pi_1 X]$  is the group ring of  $\pi_1 X$  and  $C_*(\tilde{X})$  is the  $\mathbb{Z}\pi$ -module of chains on the universal cover  $\tilde{X}$  of  $X$ .

Note now that  $\Psi(-)_*$  in (2.2) induces, by composition, a  $\pi_0 G$ -action on the  $\mathbb{Z}\pi$ -module  $C_{\mathbb{Z}\pi}^2(\tilde{X}, \pi_1 G)$ . Explicitly,  $a \in \pi_0 G$  acts on a 2-cochain  $\tau \in C_{\mathbb{Z}\pi}^2(\tilde{X}, \pi_1 G)$  as

$$(2.4) \quad a \cdot \tau = \Psi(a)_* \circ \tau.$$

Since  $\tilde{X}$  has dimension 2, every 2-cochain is a cocycle hence, passing to the quotient, (2.4) yields a  $\pi_0 G$ -action on  $H^2(X, \pi_1 \mathcal{G}_{\mu_1})$ . Let

$$H^2(X, \pi_1 \mathcal{G}_{\mu_1}) / \pi_0 G$$

be the corresponding quotient set.

We are now ready to state the result concerning the topological classification of principal  $G$ -bundles over  $X$ .

**Theorem 2.2.** *Let  $G$  be a Lie group with  $\pi_0 G$  abelian and let  $X$  be a 2-dimensional connected CW-complex. There is a bijection between the set of isomorphism classes of continuous principal  $G$ -bundles over  $X$  with invariant  $\mu_1 \in H^1(X, \pi_0 G)$  and the (non-empty) quotient set  $H^2(X, \pi_1 \mathcal{G}_{\mu_1}) / \pi_0 G$ .*

This means that the map

$$\chi : [X, BG] \rightarrow H^1(X, \pi_0 G)$$

defined in (2.1) is such that there is bijection of sets  $\chi^{-1}(\mu_1) \simeq H^2(X, \pi_1 \mathcal{G}_{\mu_1}) / \pi_0 G$ .

*Remark 2.3.* Of course if one has a topological space  $M$  which is homotopically equivalent to a finite 2-dimensional CW-complex  $X$ , then  $[M, BG] \cong [X, BG]$ , so the above theorem also applies for the classification of  $G$ -principal bundles over  $M$ . For instance,  $M$  could be the complement of a finite number of points in a compact 3-dimensional manifold.

*Remark 2.4.* Using different methods, Theorem 2.2 has actually been proved before in [14], in the particular case  $X$  is a closed oriented surface.

We will prove the theorem in section 5. In the next sections, we will briefly recall some notions (of obstruction theory and of Postnikov sections) which will be required for it. Before that, let us now consider some explicit examples (most of them quite classical) of applications of this theorem.

*Example 2.5.* Let  $X$  be any finite connected 2-dimensional CW-complex and  $G$  a Lie group with  $\pi_0 G$  abelian.

- (1) If  $G$  is connected, then  $G$ -bundles over  $X$  are classified by

$$[X, BG] \cong H^2(X, \pi_1 G)$$

as it is well-known (for surfaces, see for example [15], Proposition 5.1).

- (2) If  $\pi_0 G$  acts trivially on  $\pi_1 G$ , then  $G$ -bundles over  $X$  are classified by

$$[X, BG] \cong H^1(X, \pi_0 G) \times H^2(X, \pi_1 G).$$

Because  $\pi_0 G$  is abelian, this case yields

$$[X, BG] \cong \pi_0 G^{2g} \times \pi_1 G$$

when  $X$  is a closed (i.e., compact and without boundary) oriented surface of genus  $g$ , and yields

$$[X, BG] \cong (\pi_0 G)_2 \times \pi_0 G^{k-1} \times \pi_1 G / 2\pi_1 G$$

when  $X$  is a closed non-orientable surface which is a connected sum of  $k$  copies of the projective plane  $\mathbb{R}P^2$ . Here  $(\pi_0 G)_2$  denotes the 2-torsion subgroup of  $\pi_0 G$ .

- (3) When  $G = O(n)$ , the group of orthogonal transformations of  $\mathbb{R}^n$ , with  $n \geq 3$ , then we are in the situation of the previous item and the classification is given by the first and second Stiefel-Whitney classes  $w_1, w_2$ .

- (4) If  $X$  is simply-connected, then principal  $G$ -bundles over  $X$  are classified by

$$[X, BG] \cong H^2(X, \pi_1 G) / \pi_0 G.$$

In particular, this holds for the 2-sphere  $S^2$  and gives  $[S^2, BG] \cong \pi_1 G / \pi_0 G$ . Again this is quite classical: see [20], Section 18.

- (5) Here is another classical example, in which  $\pi_0 G$  acts non-trivially on  $\pi_1 G$ , and so which we can deduce from the above theorem. Consider  $G = \mathrm{O}(2)$ , the group of orthogonal transformations of  $\mathbb{R}^2$ .

Then  $\pi_0 \mathrm{O}(2) = \mathbb{Z}_2$  acts non-trivially on  $\pi_1 \mathrm{O}(2) = \mathbb{Z}$ , by changing the sign of the generator. Indeed, such generator can be represented by the loop  $\gamma(\theta) = \begin{pmatrix} \cos(2\pi\theta) & \sin(2\pi\theta) \\ -\sin(2\pi\theta) & \cos(2\pi\theta) \end{pmatrix}$ , with  $\theta \in [0, 1]$ , and the non-trivial element of  $\pi_0 \mathrm{O}(2)$ , which we can represent by  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ , takes it to the inverse loop  $\gamma(-\theta)$ .

- (a) Suppose  $X$  is a closed oriented surface of genus  $g$ . Then we have the isomorphism  $H^2(X, \mathbb{Z}) \cong \mathbb{Z}$  by cap product with the fundamental class. Since the action of  $\pi_0 \mathrm{O}(2) = \mathbb{Z}_2$  on  $H^2(X, \mathbb{Z})$  by (2.4) is by post-composition on the coefficients, the isomorphism  $H^2(X, \mathbb{Z}) \cong \mathbb{Z}$  is  $\pi_0 \mathrm{O}(2)$ -equivariant with the action on the right-hand-side given by (2.2). Note that this argument always applies for any coefficients  $\pi_1 G$ , and hence applies generally and not just for  $G = \mathrm{O}(2)$ .

Hence,  $\mathrm{O}(2)$ -bundles over  $X$  with  $\mu_1 = 0$ , i.e.  $w_1 = 0$  (hence which reduce to  $\mathrm{SO}(2) \cong S^1$ -bundles), are classified by  $H^2(X, \mathbb{Z}) / \mathbb{Z}_2 \cong \mathbb{Z} / \mathbb{Z}_2 \cong \mathbb{Z}_{\geq 0}$ .

Consider now  $\mathrm{O}(2)$ -bundles over  $X$  with first Stiefel-Whitney class  $w_1 = \mu_1 \neq 0$  (so non-orientable rank 2 orthogonal bundles on  $X$ ). These are hence classified by  $H^2(X, \mathbb{Z}_{w_1}) / \mathbb{Z}_2$ . Then, by Poincaré duality with local coefficients (see [18], Theorem 10.4) and by [4, Proposition 5.14 (1)] (or [22, Theorem 3.2, VI]),

$$(2.5) \quad H^2(X, \mathbb{Z}_{w_1}) \cong H_0(X, \mathbb{Z}_{w_1}) \cong \mathbb{Z} / H_{w_1} = \pi_1 \mathrm{O}(2) / H_{w_1},$$

where

$$(2.6) \quad H_{w_1} = \langle x - \gamma \cdot x \mid x \in \pi_1 \mathrm{O}(2), \gamma \in \pi_1 X \rangle$$

and  $\gamma \cdot x$  is the action of  $\gamma$  on  $x$  determined by  $w_1$  and by (2.3). From the calculation at the beginning of the example,  $H_{w_1} \cong 2\mathbb{Z}$ , so  $H^2(X, \mathbb{Z}_{w_1}) \cong \mathbb{Z}_2$ . Now we have to look at the action of  $\mathbb{Z}_2$  on the set  $H^2(X, \mathbb{Z}_{w_1})$  induced by (2.4). As above, the isomorphism  $H^2(X, \mathbb{Z}_{w_1}) \cong \mathbb{Z}_2 = \{0, 1\}$  is equivariant with respect to the  $\pi_0 \mathrm{O}(2)$ -actions (2.4) and (2.2), because duality is also given by cap product and also because the isomorphism of [4] Proposition 5.14 (1) is  $\pi_0 G$ -equivariant. Again this holds in general and not just for  $G = \mathrm{O}(2)$ . But the  $\mathbb{Z}_2 = \pi_0 \mathrm{O}(2)$ -action on  $\mathbb{Z}_2 = \{0, 1\}$  changes the sign of the generator, hence preserves the parity, thus is trivial. So  $H^2(X, \mathbb{Z}_{w_1}) / \mathbb{Z}_2 \cong \mathbb{Z}_2$ .

We conclude then the following common knowledge fact: rank 2 orthogonal bundles over a closed oriented surface  $X$  of genus  $g$  are topologically classified by the characteristic classes in

$$(\mu_1, \mu_2) \in (\{0\} \times \mathbb{Z}_{\geq 0}) \cup ((\mathbb{Z}_2)^{2g} \setminus \{0\}) \times \mathbb{Z}_2$$

where  $\mu_1$  is the first Stiefel-Whitney class and  $\mu_2$  is the (non-negative) degree when  $\mu_1 = 0$  and the second Stiefel-Whitney class when  $\mu_1 \neq 0$ .

- (b) Suppose now that  $X$  a closed but non-orientable surface. Then  $H^2(X, \mathbb{Z}) \cong \mathbb{Z} / 2\mathbb{Z} \cong \mathbb{Z}_2$ , and we have to look how  $\mathbb{Z}_2 = \pi_0 \mathrm{O}(2)$  acts via (2.4) on  $H^2(X, \mathbb{Z})$ . This is the same as looking to the action (2.2) of  $\mathbb{Z}_2 = \pi_0 \mathrm{O}(2)$  on  $\mathbb{Z} / 2\mathbb{Z} \cong \mathbb{Z}_2$  because the isomorphism  $H^2(X, \mathbb{Z}) \cong \mathbb{Z} / 2\mathbb{Z}$  is equivariant by arguments similar to the orientable case (just take instead the fundamental class of  $X$ , in homology with local coefficients, cf. [7], Example 3H.3); this holds in general and not only for  $\mathrm{O}(2)$ . From above, such action changes the sign of the generator, hence preserves the parity. So  $H^2(X, \mathbb{Z}) / \mathbb{Z}_2 \cong (\mathbb{Z} / 2\mathbb{Z}) / \mathbb{Z}_2 \cong \mathbb{Z} / 2\mathbb{Z} \cong \mathbb{Z}_2$ .

Suppose now that  $\mu_1 = w_1 \neq 0$ . Write

$$(2.7) \quad w_X : \pi_1 X \rightarrow \text{Aut}(\mathbb{Z}) \cong \mathbb{Z}_2$$

for the corresponding non-trivial orientation character associated to the oriented double cover (this is really the first Stiefel-Whitney class of  $X$  i.e. of its tangent bundle), and  $\mathbb{Z}_{w_X}$  for the associated local system on  $X$ . Then, as above, using Poincaré duality (see [18, Theorem 10.4]), and [4, Proposition 5.14 (1)], we find that

$$(2.8) \quad H^2(X, \mathbb{Z}_{w_1}) \cong H_0(X, \mathbb{Z}_{w_X} \otimes_{\mathbb{Z}\pi} \mathbb{Z}_{w_1}) \cong (\mathbb{Z}/H_{w_X} \otimes_{\mathbb{Z}} \mathbb{Z}/H_{w_1}) \cong \mathbb{Z}_2 \otimes_{\mathbb{Z}} \mathbb{Z}_2 \cong \mathbb{Z}_2,$$

where  $H_{w_1}$  is given analogously to (2.6) and  $H_{w_X} = \langle x - \gamma \cdot x \mid x \in \mathbb{Z}, \gamma \in \pi_1 X \rangle$  with  $\pi_1 X$  acting by the non-trivial orientation character  $w_X$  in (2.7) (thus so that  $H_{w_X} \cong 2\mathbb{Z}$ ). Here we used again the two first isomorphisms are  $\pi_0 G$ -equivariant with respect to (2.4) and (2.2) (and this is true for any group, not just  $O(2)$ ).

Concluding, rank 2 orthogonal bundles over a non-orientable surface  $X$  which is a connected sum of  $k$  copies of  $\mathbb{R}\mathbb{P}^2$ , are topologically classified by

$$(\mu_1, \mu_2) \in (\{0\} \times \mathbb{Z}_2) \cup (\mathbb{Z}_2^k \setminus \{0\} \times \mathbb{Z}_2) = \mathbb{Z}_2^k \times \mathbb{Z}_2.$$

- (6) Let us see now a perhaps slightly less known case. We will use, without reference, some of the results we used in the previous example, which hold in general, such as the various versions of Poincaré duality and the  $\pi_0 G$ -equivariance of the isomorphisms from the various  $H^2$  under the actions (2.2) and (2.4).

Let  $n \geq 4$  be even and consider the projective orthogonal group  $\text{PO}(n) = \text{O}(n)/\mathbb{Z}_2$ , where  $\mathbb{Z}_2 = \{\pm I_n\}$  is the center of  $\text{O}(n)$ . (If  $n$  is odd  $\text{PO}(n)$  is connected.) Since  $n$  is even,  $\pi_0 \text{PO}(n) \cong \mathbb{Z}_2$  and

$$\pi_1 \text{PO}(n) = \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_2 & \text{if } n \equiv 0 \pmod{4} \\ \mathbb{Z}_4 & \text{if } n \equiv 2 \pmod{4}. \end{cases}$$

More precisely, the universal cover of  $\text{PO}(n)$  is the  $\text{Pin}(n)$  and, if  $p : \text{Pin}(n) \rightarrow \text{PO}(n)$  is the covering projection, then, as a set (and using the abelian notation)  $\ker(p) = \{0, 1, \omega_n, -\omega_n\}$  where  $\omega_n = e_1 \cdots e_n$  is the oriented volume element of  $\text{Pin}(n)$  in the standard construction of this group via the Clifford algebra  $\text{Cl}(n)$  (see, for example, [10]). So, if  $n \equiv 0 \pmod{4}$ ,  $\pm\omega_n$  are elements of order 2, while if  $n \equiv 2 \pmod{4}$ ,  $\pm\omega_n$  have order 4. It turns out that  $\pi_0 \text{PO}(n)$  acts on  $\pi_1 \text{PO}(n)$  by leaving 0 and 1 fixed and identifying  $\pm\omega_n$ . More precisely, recall that  $\text{Pin}(n)$  is a group with two connected components,  $\text{Pin}(n)^-$  and  $\text{Spin}(n)$ , where  $\text{Pin}(n)^-$  denotes the component which does not contain the identity. We have that  $\pm\omega_n$  do not lie in the centre of  $\text{Pin}(n)$  (which equals to  $\{0, 1\}$ ). In fact,  $\omega_n$  commutes with elements in  $\text{Spin}(n)$  and anti-commutes with elements in  $\text{Pin}(n)^-$ . This explains that  $\pi_1 \text{PO}(n)/\pi_0 \text{PO}(n) = \{0, 1, [\omega_n]\}$  where we  $[\omega_n]$  denotes the class of  $\omega_n \in \pi_1 \text{PO}(n)$  in  $\pi_1 \text{PO}(n)/\pi_0 \text{PO}(n)$ , consisting by  $\pm\omega$ .

- (a) Suppose now that  $X$  is a closed oriented surface of genus  $g$ .

Hence,  $\text{PO}(n)$ -bundles on  $X$  which reduce to  $\text{PSO}(n)$  (so with  $\mu_1 = 0$ ) are classified by

$$H^2(X, \pi_1 \text{PO}(n))/\mathbb{Z}_2 \cong \pi_1 \text{PO}(n)/\mathbb{Z}_2 \cong \{0, 1, [\omega_n]\}.$$

If  $\mu_1 \neq 0$ , then reasoning in a similar manner to the  $O(2)$  case in (2.5), one shows that  $H^2(X, (\mathbb{Z}_4)_{\mu_1}) \cong \{[0], [\omega_n]\} \cong \mathbb{Z}_2$  if  $n \equiv 2 \pmod{4}$ . So 0 and 1 are identified in this twisted cohomology. In addition, 0 and  $\omega_n$  are not in the same orbit under the  $\pi_0 \text{PO}(n)$ -action, so the action of  $\mathbb{Z}_2 = \pi_0 \text{PO}(n)$  in  $\mathbb{Z}_2 = H^2(X, (\mathbb{Z}_4)_{\mu_1})$ , induced by (2.4), is trivial. So  $H^2(X, (\mathbb{Z}_4)_{\mu_1})/\mathbb{Z}_2 \cong \mathbb{Z}_2$  for  $\mu_1 \neq 0$  and  $n \equiv 2 \pmod{4}$ . The same argument gives the same result for  $n$  multiple of 4. Concluding, if  $n \geq 4$  is even,  $\text{PO}(n)$ -bundles over the surface  $X$  are classified by characteristic classes

$$(\mu_1, \mu_2) \in (\{0\} \times \{0, 1, [\omega_n]\}) \cup ((\mathbb{Z}_2^{2g} \setminus \{0\}) \times \mathbb{Z}_2).$$

So we see that, when  $n \geq 4$  is even, there are precisely  $2^{2g+1} + 1$  isomorphism classes of topological  $\mathrm{PO}(n)$ -bundles over such surface. These invariants have interpretations in terms of obstruction theory of bundles [14, Proposition 3.2]. Namely, a  $\mathrm{PO}(n)$ -bundle with  $\mu_1 = 0$  (so actually a  $\mathrm{PSO}(n)$ -bundle) lifts to  $\mathrm{SO}(n)$ -bundle if and only if  $\mu_2 = 0, 1$  and lifts to  $\mathrm{Spin}(n)$ -bundle if and only if  $\mu_2 = 0$ . A  $\mathrm{PO}(n)$ -bundle with  $\mu_1 \neq 0$  lifts to  $\mathrm{O}(n)$ -bundle if and only if it lifts to a  $\mathrm{Pin}(n)$ -bundle and this happens if and only if  $\mu_2 = 0$ . Indeed, if a  $\mathrm{PO}(n)$ -bundle with  $\mu_1 \neq 0$  lifts to  $\mathrm{O}(n)$ , then one can always choose such a lift to have vanishing second Stiefel-Whitney class; see Remark 3.6 of [14].

(b) Suppose now that  $X$  is closed and non-orientable. Then

$$H^2(X, \pi_1 \mathrm{PO}(n)) \cong \pi_1 \mathrm{PO}(n) / 2\pi_1 \mathrm{PO}(n).$$

Here,  $2\pi_1 \mathrm{PO}(n) = \{0\}$ , if  $n = 0 \pmod{4}$ , while  $2\pi_1 \mathrm{PO}(n) = \{0, 1\}$  for  $n = 2 \pmod{4}$ , so

$$H^2(X, \pi_1 \mathrm{PO}(n)) \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_2 & \text{if } n = 0 \pmod{4} \\ \{[0], [\omega_n]\} \cong \mathbb{Z}_2 & \text{if } n = 2 \pmod{4}. \end{cases}$$

By acting further by  $\mathbb{Z}_2 = \pi_0 \mathrm{PO}(n)$  as above, we see that  $\mathrm{PO}(n)$ -bundles on the non-orientable surface  $X$  which reduce to  $\mathrm{PSO}(n)$ , so with  $\mu_1 = 0$ , are classified by

$$H^2(X, \pi_1 \mathrm{PO}(n)) / \mathbb{Z}_2 \cong \begin{cases} \{0, 1, [\omega_n]\} & \text{if } n = 0 \pmod{4} \\ \{[0], [\omega_n]\} \cong \mathbb{Z}_2 & \text{if } n = 2 \pmod{4}. \end{cases}$$

Consider now  $\mu_1 \neq 0$  and, as above, let  $w_X : \pi_1 X \rightarrow \mathrm{Aut}(\mathbb{Z}) \cong \mathbb{Z}_2$  be the non-trivial orientation character, with  $\mathbb{Z}_{w_X}$  the corresponding local system on  $X$ . Then, as in (2.8) and in Example 5(a),

$$\begin{aligned} H^2(X, \pi_1 \mathrm{PO}(n)_{\mu_1}) &\cong H_0(X, \mathbb{Z}_{w_X} \otimes_{\mathbb{Z}\pi} \pi_1 \mathrm{PO}(n)_{\mu_1}) \\ &\cong \mathbb{Z}_2 \otimes_{\mathbb{Z}} \{[0], [\omega_n]\} \cong \mathbb{Z}_2 \otimes_{\mathbb{Z}} \mathbb{Z}_2 \cong \mathbb{Z}_2, \end{aligned}$$

independently of the residue of  $n$  modulo 4. The next action of  $\mathbb{Z}_2 = \pi_0 \mathrm{PO}(n)$  in  $\mathbb{Z}_2 \cong H^2(X, \pi_1 \mathrm{PO}(n)_{\mu_1})$  is trivial as in Example 5(a).

Concluding, projective orthogonal bundles of rank  $n \geq 4$  even, over a non-orientable surface  $X$  which is a connected sum of  $k$  copies of  $\mathbb{R}\mathbb{P}^2$ , are topologically classified by

$$(\mu_1, \mu_2) \in \begin{cases} (\{0\} \times \{0, 1, [\omega_n]\}) \cup (\mathbb{Z}_2^k \setminus \{0\} \times \mathbb{Z}_2) & \text{if } n = 0 \pmod{4} \\ \mathbb{Z}_2^k \times \mathbb{Z}_2 & \text{if } n = 2 \pmod{4}. \end{cases}$$

*Remark 2.6.* The explicit examples (3), (5) and (6) are valid not only for the stated groups but also for any group which is homotopically equivalent to it. For example, respectively,  $G = \mathrm{GL}(n, \mathbb{R})$ ,  $n \geq 3$ ,  $G = \mathrm{GL}(2, \mathbb{R})$  and  $G = \mathrm{PGL}(n, \mathbb{R})$ , with  $n \geq 4$  even.

### 3. REMAINDER OF OBSTRUCTION THEORY IN FIBRATIONS

Here we briefly sketch the theory of obstructions in fibrations which will be used in the proof of Theorem 2.2. Some good references, among several others, on this material are [4, 20, 22]. We always take based spaces.

Consider a fibration  $p : E \rightarrow Y$  with fibre  $F$  and let  $f : X \rightarrow Y$  be a continuous map. The group  $\pi_1 F$  acts on  $\pi_k F$  (cf. Section 6.16 of [4] or Section 4A of [7]) through free homotopies of based maps  $S^k \rightarrow F$  and the quotient  $\pi_k F / \pi_1 F$  is  $[S^k, F]$ . Assume that  $F$  is *simple*, i.e., that the action of  $\pi_1 F$  on  $\pi_k F$  is trivial for every  $k$  (in the course of the proof of Theorem 2.2, we will be in this case).

**3.1. Obstruction to extending a map.** Let  $X_1$  be the 1-skeleton of our two-dimensional CW-complex  $X$  and let

$$f_1 : X_1 \longrightarrow E$$

be a partial lift of  $f$ :

$$\begin{array}{ccccc} & & & & F \\ & & & & \downarrow \\ & & & & E \\ & \nearrow g & & & \downarrow p \\ X_1 & \hookrightarrow X & \xrightarrow{f} & Y. & \end{array}$$

The obstruction to extending  $g$  to a lift  $\tilde{f} : X \rightarrow E$  is called the *obstruction 2-cochain*, and is the the 2-cochain

$$(3.1) \quad c^2(g) \in C_{\mathbb{Z}\pi}^2(\tilde{X}, \pi_1 F)$$

defined as follows. Take any 2-cell  $e_i^2$  of  $X$ , consider a cell  $\tilde{e}_i^2$  of  $\tilde{X}$  above  $e_i^2$  and let  $h_i : \mathbb{D} \rightarrow \tilde{X}$  be the corresponding characteristic map ( $\mathbb{D} \subset \mathbb{R}^2$  denotes the closed unit disc) and define  $\varphi_i = h_i|_{S^1} : S^1 \rightarrow \tilde{X}_1$  to be the attaching map. The value of  $c^2(g)$  on  $\tilde{e}_i^2$  is

$$c^2(g)(\tilde{e}_i^2) = [g \circ \pi \circ \varphi_i] \in \pi_1 F,$$

where  $\pi : \tilde{X} \rightarrow X$  is the universal cover of  $X$ :

$$\begin{array}{ccccccc} & & & & & & F \\ & & & & & & \downarrow \\ & & & & & & E \\ & & & \nearrow g & & & \downarrow p \\ S^1 & \xrightarrow{\varphi_i} & \tilde{X} & \xrightarrow{\pi} & X_1 & \hookrightarrow X & \xrightarrow{f} Y. \end{array}$$

The reason for which  $[g \circ \pi \circ \varphi_i] \in \pi_1 F$  is the following. The composite  $p \circ g \circ \pi \circ \varphi_i : S^1 \rightarrow Y$  is null-homotopic because it equals to  $f \circ \pi \circ \varphi_i$  which extends to  $\mathbb{D} \rightarrow Y$ . By lifting this null-homotopy ( $p$  is a fibration) we obtain a homotopy between  $g \circ \pi \circ \varphi_i$  and a map  $S^1 \rightarrow F$ . So  $[g \circ \pi \circ \varphi_i] \in \pi_1 F$ . Then  $c^2(g)$  extends linearly.

We have then that  $g$  extends to a lift  $\tilde{f} : X \rightarrow E$  of  $f$  if and only if  $c^2(g) = 0$ . In particular if  $F$  is simply-connected, the existence of such lift  $\tilde{f}$  is granted.

Even though we will not need this fact here, note that the cochain  $c^2(g)$  is in fact a cocycle so represents a class in  $H^2(X, \pi_1 \mathcal{F}_{\alpha_1})$ , for a certain local system  $\pi_1 \mathcal{F}_{\alpha_1}$ . It turns out that such class vanishes if and only if  $g$  can be redefined (relative to the 0-skeleton) in such a way that it can then be extended to a lift  $\tilde{f} : X \rightarrow E$  of  $f$ .

Similarly, if one has  $g' : X_0 \rightarrow E$  defined on the 0-skeleton of  $X$  such that  $p \circ g' = f|_{X_0}$  (obviously, this is always possible), then there is an *obstruction 1-cochain*

$$(3.2) \quad c^1(g') \in C_{\mathbb{Z}\pi}^1(\tilde{X}, \pi_0 F)$$

analogously defined, which vanishes precisely when  $g'$  extends to a partial lift  $g : X_1 \rightarrow E$  of  $f$ . In particular, this holds if  $F$  is connected.

**3.2. Obstruction to the existence of vertical homotopies.** As usual, write  $I = [0, 1]$ .

**Definition 3.1.** Let  $p : E \rightarrow Y$  be a fibration and  $f : X \rightarrow Y$  a map. Two lifts  $f_0, f_1 : X \rightarrow E$  of  $f : X \rightarrow Y$  are *vertically homotopic*, if there is a *vertical homotopy* between them, i.e., a homotopy  $K : X \times I \rightarrow E$  between  $f_0$  and  $f_1$  such that  $p \circ K : X \times I \rightarrow Y$  is the constant homotopy of  $f$ .

In other words, the homotopy  $K$  preserves the fibres of  $p$ . This is a stronger relation than simply being homotopic: two lifts of  $f$  may be homotopic but not vertically homotopic.

Let  $f_0$  and  $f_1$  be two lifts of  $f : X \rightarrow Y$  and let  $K_1$  be a vertical homotopy between them on  $X_1$ , that is between their restrictions  $f_0|_{X_1}$  and  $f_1|_{X_1}$ . We can ask whether  $f_0$  and  $f_1$  are vertically homotopic, i.e., if we can extend  $K_1$  to a vertical homotopy on  $X$ . There is indeed an obstruction to the existence of such a vertical homotopy  $K$ , called the *difference 2-cochain*

$$(3.3) \quad d^2(f_0, K_1, f_1) \in C_{\mathbb{Z}\pi}^2(\tilde{X}, \pi_2 F),$$

which is nothing but a slight modification (an increase of dimension) of the obstruction 2-cochain defined in section (3.1). More precisely, the difference cochain (3.3) is an obstruction cochain to extend to  $X \times I$  the map on the 2-skeleton of  $X \times I$

$$T = f_0 \cup K_1 \cup f_1 : X \times \{0\} \cup X_1 \times I \cup X \times \{1\} \longrightarrow E$$

lifting the constant homotopy  $H_f : X \times I \rightarrow Y$ ,  $H_f(x, t) = f(x)$ .

To define  $d^2(f_0, K_1, f_1)$  more explicitly, take any 2-cell  $e_i^2$  of  $X$ , so that  $e_i^2 \times I$  is a 3-cell of  $X \times I$ . Consider a cell  $\tilde{e}_i^2 \times I$  of  $\tilde{X} \times I$  mapping to  $e_i^2 \times I$  by the covering map  $\pi \times 1 : \tilde{X} \times I \rightarrow X \times I$ , and take the corresponding attaching  $\varphi_i : S^2 \rightarrow \tilde{X} \times \{0\} \cup \tilde{X}_1 \times I \cup \tilde{X} \times \{1\}$  be the attaching map. So,

$$S^2 \xrightarrow{\varphi_i} \tilde{X} \times \{0\} \cup \tilde{X}_1 \times I \cup \tilde{X} \times \{1\} \xrightarrow{\pi \times 1} X \times \{0\} \cup X_1 \times I \cup X \times \{1\} \hookrightarrow X \times I \xrightarrow{H_f} Y.$$

$$\begin{array}{c} F \\ \downarrow \\ E \\ \downarrow p \\ Y \end{array}$$

(A dashed arrow labeled  $T$  points from the middle of the top row to the  $E$  in the right diagram.)

Then

$$(3.4) \quad d^2(f_0, K_1, f_1)(e_i^2) = c^3(T)(\tilde{e}_i^2 \times I) = [T \circ (\pi \times 1) \circ \varphi_i] \in \pi_2 F$$

where  $c^3(T)$  is the obstruction cochain analogous to (3.1) but in dimension 3. As above, the reason why  $[T \circ (\pi \times 1) \circ \varphi_i] \in \pi_2 F$  comes basically from the fact that  $\mathbb{D} \times I$  is contractible. We then have that the partial vertical homotopy  $T$  extends to a vertical homotopy  $X \times I \rightarrow E$  if and only if the 2-cochain  $d^2(f_0, K_1, f_1)$  vanishes.

Note finally that, since  $X$  has dimension 2,  $d^2(f_0, K_1, f_1)$  is actually a cocycle, therefore defines cohomology class

$$(3.5) \quad \delta^2(f_0, K_1, f_1) \in H^2(X, \pi_2 \mathcal{F}_\alpha),$$

where  $\pi_2 \mathcal{F}_\alpha$  the local system on  $X$  with fibre  $\pi_2 F$ , given by

$$(3.6) \quad \alpha : \pi_1 X \xrightarrow{f_*} \pi_1 Y \longrightarrow \text{Aut}(\pi_2 F),$$

where  $\pi_1 Y \rightarrow \text{Aut}(\pi_2 F)$  is induced by the monodromy representation  $\pi_1 Y \rightarrow \text{Aut}_*^h(F)$ , where  $\text{Aut}_*^h(F)$  denotes the space of based homotopy self-equivalences of  $F$ .

Again, there is also a similar *difference 1-cochain* in dimension 1,

$$(3.7) \quad d^1(f'_0, K_0, f'_1) \in C_{\mathbb{Z}\pi}^1(\tilde{X}, \pi_1 F)$$

which is the obstruction to the existence of vertical homotopies of lifts  $f'_0, f'_1 : X_1 \rightarrow E$  of  $f|_{X_1}$ , when we are given a partial homotopy, over the 0-skeleton  $X_0$  of  $X$ ,  $K_0 : X_0 \times I \rightarrow E$  between  $f'_0|_{X_0}$  and  $f'_1|_{X_0}$ . So  $d^1(f'_0, K_0, f'_1) = 0$  if and only if  $T' = f'_0 \cup K_0 \cup f'_1 : X \times \{0\} \cup X_0 \times I \cup X \times \{1\} \rightarrow E$  can be extended to a vertical homotopy  $X_1 \times I \rightarrow E$  between  $f'_0, f'_1$ . In particular, this is always possible if  $F$  is simply-connected.

#### 4. A REMAINDER OF POSTNIKOV SECTIONS

The material in this section is also well-known. We provide references in due course.

**4.1. Universal fibrations of Eilenberg-MacLane spaces.** We work within the category of compactly generated based spaces. Let  $A$  be an abelian group and, for  $n \geq 1$ , let  $K(A, n)$  be the corresponding Eilenberg-MacLane space.

There is a specific model for  $K(A, n)$  which is a topological abelian group. It is constructed by taking the geometrical realization of the simplicial abelian group associated, via the Dold-Kan correspondence [6], to the chain complex  $A[-n]$  which has  $A$  in dimension  $n$  and 0 elsewhere. In all that follows we use this model for  $K(A, n)$  and we choose for base point the zero of its group structure.

Consider the space  $\text{Map}(X, K(A, n))$  of all continuous maps from  $X$  to  $K(A, n)$ , with the topology determined by the compact-open topology, and so that  $\text{Map}(X, K(A, n))$  is compactly generated (see Definition 6.2 and Section 6.1.3 of [4]). For the given model of  $K(A, n)$ ,  $\text{Map}(X, K(A, n))$  is a topological abelian group.

A *group-like monoid* is a topological monoid such that the monoid of its arcwise connected components is a group (the difference from a topological group is the possible non-existence of inverses).

Let  $\text{Aut}^h(K(A, n))$  denote the group-like monoid of all self-homotopy equivalences of  $K(A, n)$ . The model of  $K(A, n)$  is functorial, thus a group homomorphism  $A \rightarrow B$  induces a continuous map  $K(A, n) \rightarrow K(B, n)$  which is also a group homomorphism. It follows that an automorphism of  $A$  induces an automorphism of the topological abelian group  $K(A, n)$ . Hence  $\text{Aut}(A)$  acts on  $K(A, n)$  by group automorphisms, so we can take their semidirect product  $\text{Aut}(A) \rtimes K(A, n)$ . This produces an inclusion

$$(4.1) \quad \text{Aut}(A) \subset \text{Aut}^h(K(A, n))$$

given by the composite  $\text{Aut}(A) \hookrightarrow \text{Aut}(A) \rtimes K(A, n) \hookrightarrow \text{Aut}^h(K(A, n))$  where the second map is the inclusion

$$(4.2) \quad \nu : \text{Aut}(A) \rtimes K(A, n) \hookrightarrow \text{Aut}^h(K(A, n)), \quad \nu(\varphi, x)(z) = \varphi(z) + x,$$

with  $+$  denoting the group operation in  $A$ . It turns out that the inclusion (4.2) is a weak homotopy equivalence.

Now, there exists [19, 11] a unique (up to homotopy equivalence) CW-complex  $B\text{Aut}^h(K(A, n))$  which is a *classifying space* for fibre homotopy equivalence classes of fibrations over  $X$ , with fibre homotopically equivalent to  $K(A, n)$ . So there is a corresponding *universal fibration*  $\text{Aut}^h(K(A, n)) \rightarrow E\text{Aut}^h(K(A, n)) \rightarrow B\text{Aut}^h(K(A, n))$ , which has similar properties to the classifying space  $BG$  and to the universal  $G$ -bundle  $EG \rightarrow BG$ , for a topological group  $G$ . In particular, the set of is in bijection with the set of free (i.e. non-based) homotopy classes of maps  $X \rightarrow B\text{Aut}^h(K(A, n))$ , i.e.,  $[X, B\text{Aut}^h(K(A, n))]$ .

Since (4.2) is a weak homotopy equivalence, and since both  $B\text{Aut}^h(K(A, n))$  and  $B(\text{Aut}(A) \rtimes K(A, n))$  are CW-complexes, it follows that  $B\text{Aut}^h(K(A, n))$  is homotopically equivalent to  $B(\text{Aut}(A) \rtimes K(A, n))$ . In other words,  $B(\text{Aut}(A) \rtimes K(A, n))$  is a model for  $B\text{Aut}^h(K(A, n))$ .

**4.2. Postnikov sections.** Write

$$P_k BG$$

for the  $k^{\text{th}}$  Postnikov section of  $BG$ . By definition (see for example [7] Chapter 4) these CW-complexes fit in a commutative diagram as below, called the *Postnikov tower* of  $BG$

$$\begin{array}{ccc}
 & & \vdots \\
 & & \downarrow \\
 & & P_2 BG \\
 & \nearrow & \downarrow \\
 BG & \longrightarrow & P_1 BG
 \end{array}$$

where the vertical maps are fibrations and such that the maps  $BG \rightarrow P_k BG$  induce isomorphisms  $\pi_i BG \cong \pi_i P_k BG$  if  $i \leq k$  and, for  $i > k$ ,  $\pi_i P_k BG = 0$ . For each  $k$ , the homotopy fibre of the fibrations  $P_k BG \rightarrow P_{k-1} BG$  is the Eilenberg-MacLane space  $K(\pi_{k-1} G, k)$ .

Again, each Postnikov section  $P_k BG$  is unique up to homotopy equivalence, hence so is the Postnikov tower. In particular,  $P_1 BG = B\pi_0 G = K(\pi_0 G, 1)$ .

Regarding the second Postnikov section  $P_2 BG$ , there is [16] a model for  $P_2 BG \rightarrow B\pi_0 G$  which is a fiber bundle with structure group  $\pi_0 G$ . To briefly see this, consider the group-like monoid  $\text{Aut}^h(K(\pi_1 G, 2))$  of self-homotopy equivalences of  $K(\pi_1 G, 2)$ . Consider the universal fibration

$$\text{Aut}^h(K(\pi_1 G, 2)) \longrightarrow E\text{Aut}^h(K(\pi_1 G, 2)) \longrightarrow B\text{Aut}^h(K(\pi_1 G, 2))$$

as defined in the previous section. Recall that  $B(\text{Aut}(\pi_1 G) \times K(\pi_1 G, 2))$  is a model for the classifying space  $B\text{Aut}^h(K(\pi_1 G, 2))$ . Let

$$(4.3) \quad \mathcal{U} = E\text{Aut}^h(K(\pi_1 G, 2)) \times_{\text{Aut}^h(K(\pi_1 G, 2))} K(\pi_1 G, 2) \longrightarrow B(\text{Aut}(\pi_1 G) \times K(\pi_1 G, 2))$$

be the  $K(\pi_1 G, 2)$ -fibration obtained from the above universal fibration.

The splitting

$$\text{Aut}(\pi_1 G) \rightarrow \text{Aut}(\pi_1 G) \times K(\pi_1 G, 2), \quad \varphi \mapsto (\varphi, 0)$$

of the exact sequence  $K(\pi_1 G, 2) \hookrightarrow \text{Aut}(\pi_1 G) \times K(\pi_1 G, 2) \rightarrow \text{Aut}(\pi_1 G)$  yields a section

$$e : B\text{Aut}(\pi_1 G) \rightarrow B(\text{Aut}(\pi_1 G) \times K(\pi_1 G, 2))$$

of the fibration  $B(\text{Aut}(\pi_1 G) \times K(\pi_1 G, 2)) \rightarrow B\text{Aut}(\pi_1 G)$  induced by the above exact sequence.

Hence the pullback of  $\mathcal{U}$  in (4.3) under the composition

$$B\pi_0 G \rightarrow B\text{Aut}(\pi_1 G) \xrightarrow{e} B(\text{Aut}(\pi_1 G) \times K(\pi_1 G, 2))$$

is the fiber bundle

$$E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2) \rightarrow B\pi_0 G,$$

with fiber  $K(\pi_1 G, 2)$ , structure group  $\pi_0 G$  and where  $E\pi_0 G \rightarrow B\pi_0 G$  is the universal  $\pi_0 G$ -principal bundle. Here  $\pi_0 G$  acts on  $K(\pi_1 G, 2)$ -factor via the composition

$$(4.4) \quad \pi_0 G \xrightarrow{\Psi^{(-)*}} \text{Aut}(\pi_1 G) \hookrightarrow \text{Aut}^h(K(\pi_1 G, 2)).$$

Thus, we use the functorial construction of the Eilenberg-MacLane space  $K(\pi_1 G, 2)$  explained in the preceding section. On the other hand, the  $\pi_0 G$ -action on the  $E\pi_0 G$ -factor is by the monodromy action of  $\pi_0 G = \pi_1(B\pi_0 G)$  (based on the base point of  $B\pi_0 G = K(\pi_0 G, 1)$ ). So, if  $\gamma : S^1 \rightarrow B\pi_0 G$  is a loop representing a class in  $\pi_0 G$  and  $v \in E\pi_0 G$ , then  $\gamma \cdot v = v + \tilde{\gamma}(1)$  where  $\tilde{\gamma}$  is the lift of  $\gamma$  through  $v$  and  $+$  is the group operation on  $E\pi_0 G$  (recall that  $E\pi_0 G$  is an abelian group because  $B\pi_0 G$  is so).

Concluding, using (4.4), if  $[a, b] \in E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)$  and  $g = \gamma \in \pi_0 G = \pi_1(B\pi_0 G)$ , then

$$(4.5) \quad [a, b] = [a + \tilde{\gamma}(1), \Psi_*(-g)(b)].$$

This is the model for  $P_2 BG$  we will use:

$$(4.6) \quad P_2 BG = E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2) \rightarrow B\pi_0 G.$$

## 5. THE PROOF

Now that we have provided the necessary background, we prove the classification Theorem 2.2. This will be done by a sequence of lemmas.

5.1. **Replacement of  $BG$  by  $P_2BG$ .** Note that the map  $BG \rightarrow P_1BG = B\pi_0G$  is  $p_{0,*}$  as defined in (2.1). The homotopy fibre of the fibration  $P_2BG \rightarrow P_1BG$  is  $K(\pi_1G, 2)$ , i.e., we obtain the fibration

$$(5.1) \quad K(\pi_1G, 2) \longrightarrow P_2BG \xrightarrow{p} B\pi_0G$$

and the commutative diagram

$$\begin{array}{ccc} & & K(\pi_1G, 2) \\ & & \downarrow \\ & & P_2BG \\ & \nearrow & \downarrow p \\ BG & \xrightarrow{p_{0,*}} & B\pi_0G. \end{array}$$

Since the second homotopy group of any Lie group vanishes (cf. [12] Chapter VI, Theorem 4.17), we have  $\pi_3BG = 0 = \pi_3P_2BG$  so actually the map  $BG \rightarrow P_2BG$  induces isomorphisms  $\pi_iBG \cong \pi_iP_2BG$ , for  $i \leq 3$  i.e. is a 3-equivalence. Since  $\dim X = 2 < 3$ , we conclude by [22], Chapter IV, Theorem 7.16 that  $BG \rightarrow P_2BG$  induces a bijection

$$(5.2) \quad [X, BG] \simeq [X, P_2BG].$$

Hence in order to classify  $G$ -bundles over  $X$ , it suffices to consider the set

$$[X, P_2BG]$$

instead of  $[X, BG]$ .

As we have seen above, obtaining the classification of  $G$ -bundles with fixed invariant  $\mu_1 \in [X, B\pi_0G]$  is equivalent to describing the fibre over  $\mu_1$  of the map  $\chi : [X, BG] \rightarrow [X, B\pi_0G]$  defined in (2.1).

Now, consider the totally similar map induced by the fibration  $p$  in (5.1)

$$(5.3) \quad \chi' : [X, P_2BG] \longrightarrow [X, B\pi_0G], \quad \chi'([h]) = [p \circ h].$$

From (5.2) and the previous diagram, we have the commutative diagram

$$\begin{array}{ccc} & & [X, P_2BG] \\ & \nearrow & \downarrow \chi' \\ [X, BG] & \xrightarrow{\chi} & [X, B\pi_0G] \end{array}$$

so the next lemma follows immediately.

**Lemma 5.1.** *There is a bijection between the set of isomorphism classes of continuous principal  $G$ -bundles over  $X$  with invariant  $\mu_1 \in [X, B\pi_0G]$  and the fibre over  $\mu_1$  of  $\chi'$ .*

Let us now begin applying the obstruction theory sketched in section 3 to our situation, thus to the fibration (5.1) and to the next diagram

$$\begin{array}{ccc} & & K(\pi_1G, 2) \\ & & \downarrow \\ & & P_2BG \\ & \nearrow & \downarrow p \\ X & \xrightarrow{f} & B\pi_0G \end{array}$$

where  $f$  is a representative of a class  $\mu_1 \in [X, B\pi_0G]$ .

We want to analyse the fibre  $\chi'^{-1}(\mu_1)$  and a first question is if it is non-empty. Being non-empty means that we can lift  $f$  to  $P_2BG$ . To see this, first notice that, since  $K(\pi_1G, 2)$  is connected, we can always find a partial lift  $f'$  of  $f$  over the 1-skeleton, i.e. a lift of  $f|_{X_1}$ ,

because (3.2) vanishes. In addition, as  $K(\pi_1 G, 2)$  is simply-connected, then the obstruction 2-cochain (3.1) is again zero and hence we can lift  $f$  to a map  $X \rightarrow P_2 BG$ . So:

**Lemma 5.2.** *The map  $\chi'$  is surjective.*

**5.2. Homotopy classes of lifts and vertical homotopy classes of lifts.** Our next task is then to describe the fibre of  $\chi'$  over the class  $\mu_1 \in [X, B\pi_0 G]$ . This fibre is in general just a set, namely the set of homotopy classes in  $[X, B\pi_0 G]$  which project to a representative of the class  $\mu_1$ , hence to a map homotopic to  $f : X \rightarrow B\pi_0 G$ . We will describe *all* such homotopy classes of lifts of  $f$  in terms of *vertical* homotopy classes of lifts of  $f$ . Denote the latter set by

$$[X, P_2 BG]_f.$$

First note that if  $g : X \rightarrow B\pi_0 G$  is another representative of the class  $\mu_1$ , then by lifting a homotopy between  $f$  and  $g$  we obtain a bijection  $[X, P_2 BG]_f \cong [X, P_2 BG]_g$ . So we loose nothing by choosing any representative  $f$  of  $\mu_1$ .

Then take the group of homotopy classes of self-homotopies of  $f$ , i.e.,  $\pi_1(\text{Map}(X, B\pi_0 G), f)$ . It acts on  $[X, P_2 BG]_f$  also by lifting these self-homotopies, that is, if  $f_0$  is a lift of  $f$ , and  $H \in \pi_1(\text{Map}(X, B\pi_0 G), f)$  is a self-homotopy of  $f$ , then  $H$  acts on  $f_0$  as  $H \cdot f_0 = \tilde{H}_1$ , where  $\tilde{H} : X \times I \rightarrow P_2 BG$  is the unique lift of  $H$  such that  $\tilde{H}_0 = f_0$ .

Since  $\pi_0 G$  is abelian,  $B\pi_0 G = K(\pi_0 G, 1)$  is a topological abelian group, hence so is  $\text{Map}(X, B\pi_0 G)$ . It follows that  $\pi_1(\text{Map}(X, B\pi_0 G), f)$  also does not depend on  $f$ .

So the orbit space

$$[X, P_2 BG]_f / \pi_1(\text{Map}(X, B\pi_0 G), f)$$

is independent on the choice of the representative  $f$  of  $\mu_1$ .

**Lemma 5.3.** *Given  $\mu_1 \in [X, B\pi_0 G]$ , there is a bijection of sets*

$$\chi'^{-1}(\mu_1) \cong [X, P_2 BG]_f / \pi_1(\text{Map}(X, B\pi_0 G), f),$$

with  $f : X \rightarrow B\pi_0 G$  representing  $\mu_1$ .

*Proof.* By definition  $\chi'^{-1}(\mu_1) \subset [X, P_2 BG]$ . Consider the natural map

$$[X, P_2 BG]_f \rightarrow \chi'^{-1}(\mu_1)$$

which sends a vertical homotopy class of a lift  $g : X \rightarrow P_2 BG$  of  $f$  to its (not necessarily vertical) homotopy class. This is clearly surjective: take any class  $g$  in  $\chi'^{-1}(\mu_1)$  and consider the class in  $[X, P_2 BG]_f$  of the maps which are equivalent to  $g$  by vertical homotopies. The image of such class is the class of  $g$  in  $\chi'^{-1}(\mu_1)$ .

Take lifts  $f_0$  and  $f_1$  of  $f$ , which are homotopic, hence represent the same point in  $\chi'^{-1}(\mu_1)$ . Projecting the homotopy through  $p : P_2 BG \rightarrow B\pi_0 G$ , yields a self-homotopy of  $f$ , whose lift is the homotopy between  $f_0$  and  $f_1$  we started with. Hence  $f_0$  and  $f_1$  represent classes in  $[X, P_2 BG]_f$  which lie in the same orbit of  $\pi_1(\text{Map}(X, B\pi_0 G), f)$ ; note that  $f_0$  and  $f_1$  are vertically homotopic precisely if the induced self-homotopy of  $f$  is trivial. Conversely, by definition of the action of  $\pi_1(\text{Map}(X, B\pi_0 G), f)$  by lifting self-homotopies of  $f$ , we identify the lifts of  $f$  which are homotopic. This shows that the above map induces an injective map

$$[X, P_2 BG]_f / \pi_1(\text{Map}(X, B\pi_0 G), f) \rightarrow \chi'^{-1}(\mu_1).$$

Since  $[X, P_2 BG]_f \rightarrow \chi'^{-1}(\mu_1)$  is surjective, then so is the induced map, and we are done.  $\square$

Now we use the model (4.6) for  $P_2 BG$ , so that we have the next diagram

$$(5.4) \quad \begin{array}{ccc} & & K(\pi_1 G, 2) \\ & & \downarrow \\ & & E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2) \\ & \nearrow & \downarrow p \\ X & \xrightarrow{f} & B\pi_0 G. \end{array}$$

The remaining part of the paper will always refer to the above diagram.

Let  $\pi_0 G = \pi_1(B\pi_0 G)$  act on the set  $[X, E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)]_f$  of vertical homotopy classes of lifts of  $f$  as in (4.4), so via the inclusion  $\text{Aut}(\pi_1 G) \subset \text{Aut}(K(\pi_1 G, 2))$ . More precisely, a lift  $f_0$  of  $f$  representing a class in  $[X, E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)]_f$  can be written as  $f_0(x) = [h_1(x), h_2(x)] \in E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)$ , where the  $h_i$  are defined up to the simultaneous  $\pi_0 G$ -action on both factors as in (4.5). Then  $g \in \pi_0 G$  acts on the representative  $f_0$  as

$$(5.5) \quad (g \cdot f_0)(x) = [h_1(x), \Psi(g)_*(h_2(x))]$$

for  $x \in X$  and this induces an action on the homotopy classes of lifts of  $f$ . Indeed  $g \cdot f_0$  is another lift of  $f$  since  $[h_1(x), \Psi(g)_*(h_2(x))]$  and  $[h_1(x), h_2(x)]$  lie in the same fiber of  $p$ .

The next lemma shows that the fibre that we are studying is in bijection with the quotient set for the action (5.5).

**Lemma 5.4.** *Let  $[X, E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)]_f$  denote the set of vertical homotopy classes of lifts of  $f$  in diagram (5.4), together with the  $\pi_0 G$ -action (5.5). Then there is a bijection*

$$\chi'^{-1}(\mu_1) \simeq [X, E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)]_f / \pi_0 G.$$

*Proof.* Using our model  $P_2 BG = E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)$ , we know from Lemma 5.3 that

$$(5.6) \quad \chi'^{-1}(\mu_1) \simeq [X, E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)]_f / \pi_1(\text{Map}(X, B\pi_0 G), f),$$

where the group  $\pi_1(\text{Map}(X, B\pi_0 G), f)$  acts by lifting self-homotopies of  $f$  in (5.4).

Now we compute  $\pi_1(\text{Map}(X, B\pi_0 G), f)$ . As we said before, this is independent of  $f$ , because  $\text{Map}(X, B\pi_0 G)$  is a topological abelian group. So we compute  $\pi_1 \text{Map}(X, B\pi_0 G)$  taking for base point the constant map

$$\text{const} \in \text{Map}(X, B\pi_0 G)$$

equal to the base point of  $B\pi_0 G = K(\pi_0 G, 1)$ . In this case,

$$(5.7) \quad \begin{aligned} \pi_1(\text{Map}(X, B\pi_0 G), \text{const}) &= [(X \times S^1)/(X \times 1), B\pi_0 G] \\ &= [SX \vee S^1, B\pi_0 G] \\ &= [S^1, B\pi_0 G] \\ &= \pi_0 G. \end{aligned}$$

Here,  $SX = (X \times I)/(X \times 0 \cup X \times 1)$  is the unreduced suspension of  $X$ , which is simply connected because  $X$  is connected, and  $\vee$  is the wedge sum. Notice that in (5.7) is not relevant to distinguish classes of based or unbased maps because  $B\pi_0 G$  is a simple space (it is 1-simple since  $\pi_0 G$  is abelian). Moreover, the third equality holds since a homotopy class of a (based) map  $f : SX \vee S^1 \rightarrow B\pi_0 G$  is defined by two homotopy classes of (based) maps  $f_1 : SX \rightarrow B\pi_0 G$  and  $f_2 : S^1 \rightarrow B\pi_0 G$ , but  $[SX, B\pi_0 G] = [SX, K(\pi_0 G, 1)] = H^1(SX, \pi_0 G) = 0$  because  $\pi_1 SX = 0$ .

Denote by  $+$  the group operation on  $B\pi_0 G = K(\pi_0 G, 1)$  and let

$$g \in \pi_0 G \cong \pi_1(B\pi_0 G).$$

If we represent  $g$  by a loop

$$\gamma : S^1 \longrightarrow B\pi_0 G$$

then the self-homotopy of  $f$  associated to  $g$  is  $H : X \times S^1 \rightarrow B\pi_0 G$  defined by

$$(5.8) \quad H(x, t) = f(x) + \gamma(t).$$

Here, the operation of summing  $f$  is the homeomorphism which identifies the component of the constant map  $\text{const} : X \rightarrow B\pi_0 G$  in  $\text{Map}(X, B\pi_0 G)$  with that of  $f$ . Now we use diagram (5.4). The element  $g \in \pi_0 G$  acts on the class of  $f_1$  by lifting the self-homotopy  $H$  of  $f$  given in (5.8) to  $E\pi_0 G \times_{\pi_0 G} K(\pi_1 G, 2)$  and taking its final value. As  $B\pi_0 G$  is an abelian group, then  $E\pi_0 G$  is also an abelian group and we denote again the operation by  $+$ . A lift of  $H$  is given by

$$\tilde{H}(x, t) = [h_1(x) + \tilde{\gamma}(t), h_2(x)]$$

where  $\tilde{\gamma}$  is a lift of  $\gamma$ . So, by (4.5),

$$\tilde{H}(x, 1) = [h_1(x) + \tilde{\gamma}(1), h_2(x)] = [h_1(x), \Psi(g)_*(h_2(x))].$$

Hence the action of  $\pi_1(\text{Map}(X, B\pi_0G), f)$  by lifting self-homotopies is compatible with the action (5.5) of  $\pi_0G$  and with the identification  $\pi_1(\text{Map}(X, B\pi_0G), f) \cong \pi_1(\text{Map}(X, B\pi_0G), \text{const}) \cong \pi_0G$  in (5.7). This, together with (5.6), completes the proof of the lemma.  $\square$

**5.3. Bijection with twisted cohomology.** Consider the fibration  $p$  in (5.4). The corresponding difference cochain (3.7) is zero so there are always vertical homotopies between lifts of  $f$  on  $X_1$ . Thus, given two lifts  $f_0$  and  $f_1$  of  $f$ , we can suppose that  $f_0|_{X_1} = f_1|_{X_1}$  and that  $K_1$  is the constant homotopy between them. We will hence write

$$\delta^2(f_0, f_1)$$

instead of  $\delta^2(f_0, K_1, f_1)$  for the cocycle (3.5) determined by the difference cochain (3.3). Moreover, the action  $\alpha$  defined in (3.6) becomes

$$\alpha : \pi_1 X \xrightarrow{f_*} \pi_1 B\pi_0G \longrightarrow \text{Aut}(\pi_2(K(\pi_1G, 2))),$$

that is

$$\alpha : \pi_1 X \xrightarrow{f_*} \pi_0G \longrightarrow \text{Aut}(\pi_1G).$$

But  $p$  is a fiber bundle with  $\pi_0G$  as the structure group, with  $\pi_0G$  acting on the fiber  $K(\pi_1G, 2)$  as in (4.4), so via  $\Psi(-)_*$ . Hence the action  $\alpha$  coincides with  $\Psi(-)_* \circ \mu_{1*}$  in (2.3). Hence we have that

$$\delta^2(f_0, f_1) \in H^2(X, \pi_1\mathcal{G}_{\mu_1}).$$

with  $\pi_1\mathcal{G}_{\mu_1}$  being the local system determined by (2.3).

The following result (Corollary 6.16 of Chapter VI of [22]) shows that there is a close relation between  $[X, E\pi_0G \times_{\pi_0G} K(\pi_1G, 2)]_f$  and  $H^2(X, \pi_1\mathcal{G}_{\mu_1})$ .

**Proposition 5.5.** *Let  $f_0 : X \rightarrow E\pi_0G \times_{\pi_0G} K(\pi_1G, 2)$  be any lift of  $f : X \rightarrow B\pi_0G$ . Then the map*

$$\phi : [X, E\pi_0G \times_{\pi_0G} K(\pi_1G, 2)]_f \longrightarrow H^2(X, \pi_1\mathcal{G}_{\mu_1}), \quad \phi(f_1) = \delta^2(f_0, f_1)$$

*is a bijection.*

Now we want to see the  $\pi_0G$ -quotient of Lemma 5.4 on the side of twisted cohomology  $H^2(X, \pi_1\mathcal{G}_{\mu_1})$ , via the correspondence of the previous proposition and via the action of  $\pi_0G$  in  $H^2(X, \pi_1\mathcal{G}_{\mu_1})$  induced from the action in  $C_{\mathbb{Z}\pi}^2(\tilde{X}, \pi_1G)$  defined in (2.4).

This is the content of the next and final lemma in the proof of Theorem 2.2.

**Lemma 5.6.** *The bijection between  $[X, E\pi_0G \times_{\pi_0G} K(\pi_1G, 2)]_f$  and  $H^2(X, \pi_1\mathcal{G}_{\mu_1})$  is  $\pi_0G$ -equivariant with respect to (5.5) and (2.4), respectively. Hence it factors through a bijection  $\pi_0G$ :*

$$[X, E\pi_0G \times_{\pi_0G} K(\pi_1G, 2)]_f / \pi_0G \simeq H^2(X, \pi_1\mathcal{G}_{\mu_1}) / \pi_0G.$$

*Proof.* Recall that we are always dealing with the situation represented by diagram (5.4). For the correspondence in Proposition 5.5, one has a choice of a lift  $f_0$  of  $f$  and the natural one is to choose the zero map, that is,  $f_0(x) = [g_1(x), 0]$ , with  $g_1(x)$  such that  $p([g_1(x), 0]) = f(x)$ .

Given another lift  $f_1$  of  $f$ , we have already seen that we can suppose that  $f_1|_{X_1} = f_0|_{X_1} = 0$ . With these choices, the cocycle  $\delta^2(f_0, f_1)$  is given by (3.3) and (3.4), with  $T = f_1$ , because  $f_0$  is the zero map and  $f_1|_{X_1} = f_0|_{X_1} = 0$  (so  $K_1$  is the constant homotopy equal to zero), so that we can collapse  $\partial\tilde{e}_i^2 \times I \cup \tilde{e}_i^2 \times 0$  to one point:

$$\delta^2(f_1)(e_i^2) = [f_1 \circ \pi \circ \varphi_i] \in \pi_2K(\pi_1G, 2) \cong \pi_1G.$$

On the other hand, by the definition of the action of  $\pi_0G$  in  $C_{\mathbb{Z}\pi}^2(\tilde{X}, \pi_1G)$  in (2.4), we have that, for  $g \in \pi_0G$ ,

$$(5.9) \quad (g \cdot \delta^2(f_1))(e_i^2) = \Psi(g)_*([f_1 \circ \pi \circ \varphi_i]) \in \pi_1G \cong \pi_2(K(\pi_1G, 2)).$$

But by (4.4), if we write  $f_1(x) = [h_1(x), h_2(x)]$ , we conclude that

$$(5.10) \quad \Psi(g)_*([f_1 \circ \pi \circ \varphi_i]) = [f'_1 \circ \pi \circ \varphi_i]$$

where  $f'_1(x) = [h_1(x), \Psi(g)_*(h_2(x))]$ .

By (5.5), (5.9) and (5.10), we have that

$$g \cdot \delta^2(f_1) = \delta^2(g \cdot f_1).$$

We conclude that the bijection stated in Proposition 5.5 is  $\pi_0 G$ -equivariant and this completes the proof.  $\square$

Finally, the proof Theorem 2.2 follows from Lemmas 5.1, 5.2, 5.4 and 5.6.

## REFERENCES

- [1] T. Baird, Moduli spaces of flat  $SU(2)$ -bundles over non-orientable surfaces, *Quart. J. Math.* **61**, no. 2 (2010), 141–170.
- [2] T. Baird, Moduli spaces of flat  $SU(3)$ -bundles over a Klein bottle, *Math. Res. Lett.* **17** (2010), 843–849.
- [3] G. Daskalopoulos, C. Mese, G. Wilkin, Higgs bundles over cell complexes and representations of finitely presented groups, *Pacific J. Math.* **296**, No.1 (2018), 31–55.
- [4] J. F. Davis, P. Kirk, *Lecture Notes in Algebraic Topology*, Graduate Studies in Mathematics **35**, American Mathematical Society, 2001.
- [5] O. García-Prada, A. Oliveira, Connectedness of Higgs bundle moduli for complex reductive Lie groups, *Asian J. Math.* **21**, No.5 (2017), 791–810.
- [6] P. G. Goerss, J. F. Jardine, *Simplicial Homotopy Theory*, Progress in Mathematics **174**, Birkhäuser-Verlag, 1999.
- [7] A. Hatcher, *Algebraic Topology*, Cambridge University Press, 2002.
- [8] N. J. Hitchin, The self-duality equations on a Riemann surface, *Proc. London Math. Soc.* (3) **55** (1987), 59–126.
- [9] N-K. Ho, C-C. M. Liu, Connected components of the space of surface group representations, *Int. Math. Res. Not.* **2003**, Issue 44 (2003), 2359–2372.
- [10] H. B. Lawson, M. L. Michelsohn, *Spin Geometry*, Princeton Mathematical Series **38**, Princeton University Press, 1989.
- [11] J. P. May, *Classifying Spaces and Fibrations*, Mem. Amer. Math. Soc. **1**, (1975), issue 1, no.155.
- [12] M. Mimura, H. Toda, *Topology of Lie Groups, I and II*, Translations of Mathematical Monographs **91**, American Mathematical Society, 2000.
- [13] M. S. Narasimhan, C. S. Seshadri, Stable and unitary vector bundles on a compact Riemann surface, *Ann. Math.* **82** (1965), 540–567.
- [14] A. Oliveira, Representations of surface groups in the projective general linear group, *Int. J. Math.* **2** (2011), 223–279.
- [15] A. Ramanathan, Stable principal bundles on a compact Riemann surface, *Math. Ann.* **213** (1975), 129–152.
- [16] C. A. Robinson, Moore-Postnikov systems for non-simple fibrations, *Illinois J. Math.* **16**, Issue 2 (1972), 234–242.
- [17] F. Schaffhauser, Moduli spaces of vector bundles over a Klein surface, *Geom. Dedicata* **151** (2011), 187–206.
- [18] E. Spanier, Singular homology and cohomology with local coefficients and duality for manifolds, *Pacific J. Math.* **160** (1993), 165–200.
- [19] J. Stasheff, A classification theorem for fibre spaces, *Topology* **2** (1963), 239–246.
- [20] N. Steenrod, *The Topology of Fibre Bundles*, Princeton Mathematical Series **14**, Princeton University Press, 1999.
- [21] R. Wentworth, Higgs Bundles and Local Systems on Riemann Surfaces, In: Alvarez Consul L., Andersen J., Mundet i Riera I. (eds) *Geometry and Quantization of Moduli Spaces*. Advanced Courses in Mathematics - CRM Barcelona. Birkhäuser, Cham. (2016), 165–219.
- [22] G. W. Whitehead, *Elements of Homotopy Theory*, Graduate Texts in Mathematics **61**, Springer-Verlag, 1978.

### André G. Oliveira

Centro de Matemática da Universidade do Porto, CMUP

Faculdade de Ciências, Universidade do Porto

Rua do Campo Alegre 687, 4169-007 Porto, Portugal

[www.fc.up.pt](http://www.fc.up.pt)

email: [andre.oliveira@fc.up.pt](mailto:andre.oliveira@fc.up.pt)

<https://sites.google.com/view/aoliveira>

*On leave from:*

Departamento de Matemática, Universidade de Trás-os-Montes e Alto Douro, UTAD

Quinta dos Prados, 5000-911 Vila Real, Portugal  
[www.utad.pt](http://www.utad.pt)  
email: [agoliv@utad.pt](mailto:agoliv@utad.pt)