

# LARGE DEVIATIONS FOR NON-UNIFORMLY EXPANDING MAPS

V. ARAÚJO AND M. J. PACIFICO

ABSTRACT. We obtain large deviation results for non-uniformly expanding maps with non-flat singularities or criticalities and for partially hyperbolic non-uniformly expanding attracting sets. That is, given a continuous function we consider its space average with respect to a physical measure and compare this with the time averages along orbits of the map, showing that the Lebesgue measure of the set of points whose time averages stay away from the space average decays to zero exponentially fast with the number of iterates involved. As easy by-products we deduce escape rates from subsets of the basins of physical measures for these types of maps.

## CONTENTS

1. Introduction	2
1.1. Statement of the results	3
1.2. Partially hyperbolic diffeomorphisms	6
1.3. Escape rates	8
1.4. Organization of the paper	8
2. Examples of application	9
2.1. Quadratic maps and infinite-modal maps	9
2.2. Piecewise smooth one-dimensional expanding maps	9
2.3. Non-uniformly expanding local diffeomorphisms	10
2.4. Viana maps	10
2.5. Partially hyperbolic non-uniformly expanding diffeomorphisms	10
3. Hyperbolic times	11
3.1. Coverings by hyperbolic pre-balls	12
3.2. The partially hyperbolic setting	15
3.3. The volume of dynamical balls	19
4. Hyperbolic times and large deviations	21
4.1. Upper bound for large deviations	21
4.2. Upper bound for partially hyperbolic diffeomorphisms	23
4.3. Upper bound with singular/critical set	23
5. Strictly negative upper bound	25
5.1. The local diffeomorphism and partially hyperbolic case	26

---

*Date:* January 18, 2006.

*2000 Mathematics Subject Classification.* 37D25, 37A50, 37B40, 37C40.

*Key words and phrases.* non-uniform expansion, physical measures, hyperbolic times, large deviations.

V.A. was partially supported by FCT-Portugal through CMUP and POCI/MAT/61237/2004. M.J.P. was partially supported by CNPq-Brazil/FAPERJ-Brazil/Pronex Dynamical Systems.

5.2. The case with singular/critical set	26
5.3. Ruelle's Inequality for maps with non-flat singularities	27
References	31

## 1. INTRODUCTION

Smooth Ergodic Theory provides asymptotic information of the behavior of a dynamical system, given by a smooth transformation, when times goes to infinity. This statistical approach to Dynamics has provided valuable insights into many phenomena: from the remarkable result of Jakobson [24] (see also [10, 11]) showing the existence of many (positive Lebesgue measure of) parameters  $a \in (0, 2)$  for which the corresponding map of the quadratic family  $x \mapsto a - x^2$  has positive Lyapunov exponent along almost every orbit; a different set of ideas in higher dimensions provided the first clue to the nature of the Hénon attractor [11, 31] or the existence of robust classes of maps which are not uniformly expanding but exhibit several distinct positive Lyapunov exponents [43], to the study of the statistical properties of these and other classes of systems [35, 12, 47, 14, 2, 16, 6].

The basic ideas can be traced back to the Boltzman Ergodic Hypothesis from Statistical Mechanics which was the main motivation behind the celebrated Birkhoff's Ergodic Theorem ensuring the equality between temporal and spatial averages with respect to a probability measure  $\mu$  invariant under a measurable transformation  $f : M \rightarrow M$  of a compact manifold  $M$ , i.e. for every continuous map  $\varphi : M \rightarrow \mathbb{R}$  we have

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{j=0}^{n-1} \varphi(f^j(x)) = \int \varphi d\mu \quad (1)$$

for  $\mu$  almost every point  $x \in M$ . Defining  $B(\mu)$ , the *ergodic basin* of  $\mu$ , to be the set of points for which (1) holds for every continuous function  $\varphi$ , the Ergodic Theorem says that  $\mu(B(\mu)) = 1$  for all ergodic  $f$ -invariant probability measures  $\mu$ . Since ergodic measures can be, for instance, Dirac masses concentrated on periodic orbits, the Ergodic Theorem in itself does not always provide information about the asymptotic behavior of “big” subsets of points. The notion of “big” can arguably be taken as meaning “having positive Lebesgue measure (or positive volume)”, since such sets are in principle “observable sets” when interpreting  $f : M \rightarrow M$  as a model of physical, biological or economic phenomena. Correspondingly invariant probability measures  $\mu$  for which  $B(\mu)$  has positive volume are called *physical* (or Sinai-Ruelle-Bowen) measures.

This kind of measures was first constructed for (uniformly) hyperbolic diffeomorphisms by Sinai, Ruelle and Bowen [42, 39, 17]. Such measures for non-uniformly hyperbolic maps were obtained more recently [35, 12, 13, 2].

We say that a local diffeomorphism  $f$  of a compact manifold is (uniformly) *expanding* if there exists  $n \geq 1$  such that for all  $x$  and every non-zero tangent vector  $v$  at  $x$

$$\|Df^n(x)v\| \geq 2\|v\|. \quad (2)$$

For diffeomorphisms of compact manifolds, the notion of *hyperbolicity* requires the existence of two complementary directions given by two (continuous) subbundles  $E$  and  $F$  of the tangent bundle admitting  $n \geq 1$  such that for all points  $x$  and non-zero tangent vectors  $(u, v) \in E_x \oplus F_x$

$$\|Df^n(x)u\| \leq \frac{1}{2}\|u\| \quad \text{and} \quad \|Df^n(x)v\| \geq 2\|v\|. \quad (3)$$

The statistical properties of physical measures are an object of intense study, see e.g. [17, 47, 14, 3, 5, 4, 7, 21]. The leitmotif is that the sequence  $\{\varphi \circ f^n\}_{n \geq 0}$  behaves like an i.i.d. random variable, at least asymptotically.

Here we are concerned with the rate of convergence of the time averages (1) for non-uniformly expanding maps and partially hyperbolic non-uniformly expanding diffeomorphisms (where condition (2) and the right hand side condition of (3) are replaced by an asymptotic one, see the statement of results below), extending some of the large deviation results in [46] (see also [19, 20] for a different presentation).

This again strenghtens in a definite sense the idea that non-uniformly hyperbolic systems are *chaotic*: they satisfy a version of the classical large deviation results for i.i.d. random variables. More precisely, if we set  $\delta > 0$  as an acceptable error margin and consider

$$B_n = \left\{ x \in M : \left| \frac{1}{n} \sum_{j=0}^{n-1} \varphi(f^j(x)) - \int \varphi d\mu \right| > \delta \right\}$$

then we are interested in knowing whether the Lebesgue measure of  $B_n$  decays to zero exponentially fast, i.e. wheather there are constants  $C, \xi > 0$  such that

$$\text{Leb}(B_n) \leq Ce^{-\xi n} \quad \text{for all } n \geq 1. \quad (4)$$

We are able to obtain such rates for non-uniformly expanding local diffeomorphisms and also for endomorphisms and maps with non-flat singularities and criticalities under a condition on the rate of approximation of most orbits to the critical/singular set. In particular we are able to obtain an exponential decay rate as above for piecewise expanding maps with infinitely many smoothness domains, for quadratic maps corresponding to a positive Lebesgue measure subset of parameters and for a class of maps with infinitely many critical points. Moreover we also obtain the same kind of rates for partially hyperbolic attracting sets with a non-uniformly expanding direction.

The values of  $C, \xi > 0$  in (4) depend on  $\delta, \varphi$  and on global invariants for the map  $f$  which are also the object of study of Smooth Ergodic Theory, such as the metric entropy and the pressure function of  $f$ , as detailed below.

**1.1. Statement of the results.** We denote by  $\|\cdot\|$  a Riemannian norm on the compact boundaryless manifold  $M$ , by  $d$  the induced distance and by  $\text{Leb}$  a Riemannian volume form, which we call *Lebesgue measure* or *volume* and assume to be normalized:  $\text{Leb}(M) = 1$ .

We start by describing one of the class of maps that we are going to consider. Let  $f : M \rightarrow M$  be a map of the compact manifold  $M$  which is a  $C^2$  local diffeomorphism outside a set  $\mathcal{S} \subset M$  with zero Lebesgue measure.

We assume that  $f$  behaves like a power of the distance close to  $\mathcal{S}$ : there are constants  $B > 1$  and  $\beta > 0$  for which

$$(S1) \quad \frac{1}{B}d(x, \mathcal{S})^\beta \leq \frac{\|Df(x)v\|}{\|v\|} \leq Bd(x, \mathcal{S})^{-\beta};$$

$$(S2) \quad |\log \|Df(x)^{-1}\| - \log \|Df(y)^{-1}\|| \leq B \frac{d(x, y)}{d(x, \mathcal{S})^\beta};$$

$$(S3) \quad |\log |\det Df(x)^{-1}| - \log |\det Df(y)^{-1}|| \leq B \frac{d(x, y)}{d(x, \mathcal{S})^\beta};$$

for every  $x, y \in M \setminus \mathcal{S}$  with  $d(x, y) < d(x, \mathcal{S})/2$  and  $v \in T_x M \setminus \{0\}$ . The singular set  $\mathcal{S}$  may be thought of as containing those points  $x$  where  $Df(x)$  is either not defined or else is non-invertible. Note in particular that  $\mathcal{S}$  contains the set  $\mathcal{C}$  of critical points of  $f$ , i.e. the set of points (which may be empty) where  $Df(x)$  is not invertible. We refer to this kind of singular sets as *non-flat* since conditions (S1) to (S3) above are natural generalizations to arbitrary dimensions of the notion of non-flat critical point from one-dimensional dynamics, see e.g.[18].

In what follows we write  $S_n \varphi(x)$  for  $\sum_{i=0}^{n-1} \varphi(f^i(x))$  and a function  $\varphi : M \rightarrow \mathbb{R}$ . We say that  $f$  as above is *non-uniformly expanding* if there exists  $c > 0$  such that

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} S_n \psi(x) \leq -c \quad \text{where} \quad \psi(x) = \log \|Df(x)^{-1}\|, \quad (5)$$

for Lebesgue almost every  $x \in M$ . We need to control the rate of approximation of most orbits to the singular set. We say that  $f$  has *slow recurrence to the singular set*  $\mathcal{S}$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} S_n \Delta_\delta(x) < \varepsilon \quad \text{with} \quad \Delta_\delta(x) = |\log d_\delta(x, \mathcal{S})| \quad (6)$$

for Lebesgue almost every  $x \in M$ , where for any given  $\delta > 0$  we define the *smooth  $\delta$ -truncated distance* from  $x \in M$  to  $\mathcal{S}$  by

$$d_\delta(x, \mathcal{S}) = \xi_\delta(d(x, \mathcal{S})) \cdot d(x, \mathcal{S}) + 1 - \xi_\delta(d(x, \mathcal{S}))$$

where  $\xi_\delta : \mathbb{R} \rightarrow [0, 1]$  is a standard  $C^\infty$  auxiliary function satisfying

$$\xi_\delta(x) = 1 \text{ if } |x| \leq \delta \text{ and } \xi_\delta(x) = 0 \text{ if } |x| \geq 2\delta.$$

Observe that  $\Delta_\delta$  is non-negative and continuous away from  $\mathcal{S}$  and identically zero  $2\delta$ -away from  $\mathcal{S}$ .

These notions were presented in [6] for higher dimensional maps abstracted from similar notions from one-dimensional maps [18] and previous work on maps with singularities [25], and in [6, 1] the following result on existence of finitely many physical measures was obtained.

**Theorem 1.1.** *Let  $f : M \rightarrow M$  be a  $C^2$  local diffeomorphism outside a non-flat singular set  $\mathcal{S}$ . Assume that  $f$  is non-uniformly expanding with slow recurrence to  $\mathcal{S}$ . Then there are finitely many physical (or Sinai-Ruelle-Bowen) measures  $\mu_1, \dots, \mu_k$  whose basins cover the manifold Lebesgue almost everywhere, that is  $B(\mu_1) \cup \dots \cup B(\mu_k) = M$ ,  $\text{Leb} - \text{mod } 0$ .*

We say that  $f$  is a *regular map* if  $f_* \text{Leb} \ll \text{Leb}$ , that is, if  $E \subset M$  is such that  $\text{Leb}(E) = 0$ , then  $\text{Leb}(f^{-1}(E)) = 0$ . We denote by  $\mathcal{M}_f$  the family of

all invariant probability measures with respect to  $f$ , by  $\mathcal{M}_f^e$  the family of all *ergodic*  $f$ -invariant probability measures, and define

$$B(x, n, \varepsilon) = \{y \in M : d(f^i(x), f^i(y)) < \varepsilon, i = 0, \dots, n-1\}$$

the  $(n, \varepsilon)$ -dynamical ball around  $x \in M$ . Large deviation statements are usually related to *entropies*: for any finite Borel measure  $m$  on  $M$  we define

$$h_m(f)(x) = \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} -\frac{1}{n} \log m(B(x, n, \varepsilon))$$

and for  $\nu \in \mathcal{M}_f$

$$h_m(f, \nu) = \nu - \text{ess sup } h_m(f).$$

**Theorem A.** *Let  $f : M \rightarrow M$  be a regular  $C^{1+\alpha}$  local diffeomorphism outside a non-flat singular set  $\mathcal{S}$ , for some  $\alpha \in (0, 1)$ . Assume that  $f$  is non-uniformly expanding with slow recurrence to  $\mathcal{S}$ . Then writing  $J = \log |\det Df|$ , given  $c \in \mathbb{R}$  and a continuous function  $\varphi : M \rightarrow \mathbb{R}$*

(1) *if  $h_{\text{top}}(f) < \infty$ , then*

$$\begin{aligned} \liminf_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb} \left( \left\{ x \in M : \frac{1}{n} S_n \varphi(x) > c \right\} \right) \\ \geq \sup \left\{ h_\nu(f) - h_{\text{Leb}}(f, \nu) : \nu \in \mathcal{M}_f^e, \int \varphi d\nu > c \right\}; \end{aligned}$$

(2) *if  $\mathcal{S} = \emptyset$  ( $f$  is a local diffeomorphism) then*

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb} \left( \left\{ x \in M : \frac{1}{n} S_n \varphi(x) \geq c \right\} \right) \\ \leq \sup \left\{ h_\nu(f) - \int J d\nu : \nu \in \mathcal{M}_f, \int \varphi d\nu \geq c \right\}. \end{aligned}$$

(3) *in general for any given  $\eta > 0$  there exists  $\varepsilon, \delta > 0$  such that*

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb} \left( \left\{ x \in M : \frac{1}{n} S_n \varphi(x) \geq c \text{ and } \frac{1}{n} S_n \Delta_\delta(x) \leq \varepsilon \right\} \right) \\ \leq \eta + \sup \left\{ h_\nu(f) - \int J d\nu : \nu \in \mathcal{M}_f, \int \varphi d\nu \geq c \text{ and } \Delta_\delta \in L^1(\nu) \right\}. \end{aligned}$$

We say that a measure  $\nu \in \mathcal{M}_f$  is an *equilibrium state* for  $f$  with respect to  $J$  (or just an *equilibrium state* in what follows) if

$$h_\nu(f) = \nu(J) = \int J d\nu.$$

As the above statement shows, equilibrium states are involved in the determination of the asymptotic rates of deviation. Given  $\varepsilon, \delta > 0$  we write  $\mathbb{E} = \mathbb{E}_{\varepsilon, \delta}$  for the family of all equilibrium states  $\mu$  of  $f$  with respect to  $J$  such that  $\mu(\Delta_\delta) \leq \varepsilon$  and, given a continuous  $\varphi : M \rightarrow \mathbb{R}$ , we define  $\mathbb{E}(\varphi) = \{\nu(\varphi) : \nu \in \mathbb{E}\}$ .

From Theorem A we are able to deduce that the supremum above is strictly negative for non-uniformly expanding maps with slow recurrence to the singular set.

**Theorem B.** *Let  $f : M \rightarrow M$  be a local diffeomorphism outside a non-flat singular set  $\mathcal{S}$  which is non-uniformly expanding and has slow recurrence to  $\mathcal{S}$ . For  $\omega > 0$  and a continuous function  $\varphi : M \rightarrow \mathbb{R}$  there exists  $\varepsilon, \delta > 0$  arbitrarily close to 0 such that, writing*

$$A_n = \{x \in M : \frac{1}{n} S_n \Delta_\delta(x) \leq \varepsilon\}$$

and

$$B_n = \left\{ x \in M : \inf \left\{ \left| \frac{1}{n} S_n \varphi(x) - \eta(\varphi) \right| : \eta \in \mathbb{E} \right\} > \omega \right\} \quad (7)$$

we get

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb}(A_n \cap B_n) < 0. \quad (8)$$

Clearly if  $\mathcal{S} = \emptyset$  ( $f$  is a local diffeomorphism) then  $A_n = M$  and we obtain an asymptotic large deviation rate for the sets  $B_n$ . Otherwise to get a similar upper bound for  $\text{Leb}(B_n)$  we need an extra assumption on the decay of the measure of the *tail sets*  $M \setminus A_n$ .

**Corollary C.** *In the setting of Theorem B with  $\mathcal{S} \neq \emptyset$ , if  $f$  also satisfies*

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Leb}(M \setminus A_n) < 0 \quad (9)$$

then we have also

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Leb}(B_n) < 0.$$

**Remark 1.2.** Observe that if  $\mu$  is a  $f$ -ergodic absolutely continuous probability measure, then the slow recurrence condition (6) is the same as saying that  $\log d(x, \mathcal{S})$  is  $\mu$ -integrable.

Note that for any  $C^2$  endomorphism  $f$  (i.e. the singular set  $\mathcal{S}$  of  $f$  coincides with the critical set  $\mathcal{C}$  of  $f$ ) we have  $|\log d(x, \mathcal{C})| \geq \Delta_\delta(x)$  and, as shown in [27], the function  $\log d(x, \mathcal{C})$  is  $\mu$ -integrable for every  $f$ -invariant probability measure. However we need to deal with families of invariant probability measures for which  $\log d(x, \mathcal{C})$  is *uniformly integrable* so that the proofs of Theorems A and B can be carried out with our arguments. This is why we need the sets  $A_n$  in the previous statements. To the best of our knowledge no such general integrability result for  $\log d(x, \mathcal{S})$  exists with respect to invariant probability measures for maps with non-flat singularities.

**1.2. Partially hyperbolic diffeomorphisms.** Let now  $f : M \rightarrow M$  be a  $C^2$  diffeomorphism. We say that a compact  $f$ -invariant set  $\Lambda$  is an *attracting set* if it admits a *trapping region*, that is, an open neighborhood  $U \subset \Lambda$  such that  $\overline{f(U)} \subset U$  and  $\Lambda = \bigcap_{n \geq 0} f^n(U)$ . Note that we may have  $\Lambda = U = M$  (where  $M$  is connected).

As shown in [46], for every attracting set  $\Lambda$  and every physical probability measure  $\nu$  supported in  $\Lambda$ , given  $\delta > 0$  and a continuous  $\varphi : \overline{U} \rightarrow \mathbb{R}$  we have

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \log \text{Leb} \left\{ \left| \frac{1}{n} S_n \varphi - \int \varphi d\mu \right| > \delta \right\} \geq \\ \sup \left\{ h_\nu(f) - \int \Sigma^+ d\nu : \nu \in \mathcal{M}_e, \left| \int \varphi d\nu - \int \varphi d\mu \right| \geq \delta \right\}. \end{aligned}$$

Here  $\Sigma^+$  denotes the sum of the positive Lyapunov exponents at a given point of  $M$ . Recall that Ruelle's Inequality  $h_\mu(f) \leq \int \Sigma^+ d\mu$  is true of every  $C^1$ -diffeomorphism [40].

An attracting set  $\Lambda$  is *partially hyperbolic* (see e.g. [35, 15]) if there exists a continuous splitting  $E \oplus F$  of the tangent bundle of  $M$  over  $\Lambda$  along two complementary vector subbundles satisfying

- $Df$ -invariance:  $Df(E_x) = E_{f(x)}$  and  $Df(F_x) = F_{f(x)}$  for all  $x \in \Lambda$ ;
- domination: there exists  $n \geq 1$  such that

$$\|Df^n|E_x\| \cdot \|(Df^n|F_x)^{-1}\| \leq \frac{1}{2} \quad \text{for all } x \in \Lambda;$$

- $E$  is uniformly contracting: there is  $n \geq 1$  such that  $\|Df^n|E_x\| \leq \frac{1}{2}$  for all  $x \in \Lambda$ .

In this setting we denote by  $J$  the Jacobian along the centre-unstable direction  $J(x) = |\det Df|F_x|$  and by  $\mathbb{E}$  the family of all *equilibrium states* with respect to  $J$ , i.e. the set of all  $f$ -invariant probability measures  $\nu$  such that  $h_\nu(f) = \nu(J)$ .

We will assume further that the  $F$  direction only has positive Lyapunov exponents in the following sense, introduced in [6]. We say that a partially hyperbolic attractor with trapping region  $U$  is *non-uniformly expanding* if there exists  $c > 0$  such that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \log \|(Df|F_{f^j(x)})^{-1}\| \leq -c$$

for Lebesgue almost every point  $x \in U$ . In [6] the following was obtained.

**Theorem 1.3.** *Let  $\Lambda$  be a partially hyperbolic non-uniformly expanding attracting set for a  $C^2$  diffeomorphism  $f$  with trapping region  $U$ . Then there are finitely many equilibrium states which are physical measures supported in  $\Lambda$ , and whose basins cover  $U$  except for a subset of zero Lebesgue measure.*

We are able to obtain an upper bound entirely analogous to item 2 of Theorem A replacing  $M$  by the points in the trapping region  $U$  of a partially hyperbolic non-uniformly expanding attracting set  $\Lambda$  for a  $C^2$  diffeomorphism. Then for the same kind of attracting sets we obtain an upper bound for the subset corresponding to (7).

**Theorem D.** *Let  $f : M \rightarrow M$  be a  $C^2$  diffeomorphism exhibiting a partially hyperbolic non-uniformly expanding attracting set  $\Lambda$  with isolating neighborhood  $U \supset \Lambda$ . Given  $\omega > 0$  and a continuous  $\varphi : \bar{U} \rightarrow \mathbb{R}$ , define*

$$B_n = \left\{ x \in U : \inf \left\{ \left| \frac{1}{n} S_n \varphi(x) - \eta(\varphi) \right| : \eta \in \mathbb{E} \right\} > \omega \right\}.$$

*Then*

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Leb}(B_n) < 0.$$

**1.3. Escape rates.** Using the estimates obtained above and the observation that for any compact subset  $K$  and a given  $\varepsilon > 0$  we can find an open set  $W \supset K$  and a continuous function  $\varphi : M \rightarrow \mathbb{R}$  such that

- $\text{Leb}(W \setminus K) < \varepsilon$ ;
- $0 \leq \varphi \leq 1$ ,  $\varphi|_K \equiv 1$  and  $\varphi|(M \setminus W) \equiv 0$ ,

we see that for  $n \geq 1$

$$\{x \in K : f(x) \in K, \dots, f^{n-1}(x) \in K\} \subset \left\{x \in M : \frac{1}{n} S_n \varphi(x) \geq 1\right\} \quad (10)$$

and so we get the following (recall the definition of  $A_n$  in the statement of Theorem B).

**Corollary E.** *Let  $f : M \rightarrow M$  be a local diffeomorphism outside a non-flat singular set  $\mathcal{S}$  which is non-uniformly expanding and has slow recurrence to  $\mathcal{S}$ . Let  $K$  be a compact subset such that  $\mu(K) < 1$  for all  $\mu$  in the weak\*-closure  $\overline{\mathbb{E}}$  of  $\mathbb{E}$ . Then for a pair  $\varepsilon, \delta > 0$  close to 0*

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb} \left( \{x \in K \cap A_n : f^j(x) \in K, j = 1, \dots, n-1\} \right) < 0.$$

Moreover if  $\limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Leb}(M \setminus A_n) < 0$  then

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb} \left( \{x \in K, f(x) \in K, \dots, f^{n-1}(x) \in K\} \right) < 0.$$

In the setting of a partially hyperbolic non-uniformly expanding attracting set we get, using the same reasoning as above

**Corollary F.** *Let  $f : M \rightarrow M$  be a diffeomorphism and  $\Lambda$  a partially hyperbolic non-uniformly expanding attracting set with isolating neighborhood  $U$ . Let  $K \subset U$  be a compact subset such that  $\mu(K) < 1$  for all  $\mu$  in the weak\*-closure  $\overline{\mathbb{E}}$  of  $\mathbb{E}$ . Then*

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb} \left( \{x \in K, f(x) \in K, \dots, f^{n-1}(x) \in K\} \right) < 0.$$

**Remark 1.4.** All the arguments use in fact that  $f$  is  $C^1$  and that its derivative  $Df$  is  $\alpha$ -Hölder continuous with respect to the fixed Riemannian norm on  $M$ , so that all we need is  $f$  to be a  $C^{1+\alpha}$  local diffeomorphism outside the singular set, for some  $\alpha \in (0, 1)$ .

**Remark 1.5.** Recently Pinheiro [36] has extended the statement of Theorem 1.1 replacing the limsup in condition (5) by liminf, keeping the same conclusions involving the existence of finitely many physical measures and of a positive density of hyperbolic times Lebesgue almost everywhere. Hence our statements are automatically valid in this more general setting.

**1.4. Organization of the paper.** We start by presenting some non-trivial classes of maps to which our results are applicable, in Section 2. In Section 3 we present preliminary technical results to be used in the following sections. Theorem A is then proved in Subsection 4.1 for local diffeomorphisms, in Subsection 4.2 for partially hyperbolic non-uniformly expanding diffeomorphisms and in Subsection 4.3 for maps with singularities or criticalities. In Section 5 we deduce Theorem B from Theorem A, first for local diffeomorphisms and for the partially hyperbolic case in Subsection 5.1, and then with

singularities or criticalities in Subsection 5.2, together with an extension of Ruelle's Inequality to maps with non-flat singularities in Subsection 5.3.

## 2. EXAMPLES OF APPLICATION

Here we show that there are many examples of maps in the conditions of Theorem B, Corollary C or Theorem D.

**2.1. Quadratic maps and infinite-modal maps.** In [8] the following  $C^\infty$  family of maps of  $I = [-1, 1]$  with infinitely many critical points was considered:

$$f_\mu(z) = \begin{cases} f(z) + \mu & \text{for } z \in (0, \varepsilon] \\ f(z) - \mu & \text{for } z \in [-\varepsilon, 0) \end{cases}$$

where  $f : I \rightarrow I$  is an expanding extension of

$$\hat{f} : [-\varepsilon, \varepsilon] \rightarrow [-1, 1], \quad \hat{f}(z) = \begin{cases} az^\alpha \sin(\beta \log(1/z)) & \text{if } z > 0 \\ -a|z|^\alpha \sin(\beta \log(1/|z|)) & \text{if } z < 0, \end{cases}$$

to  $I$  (i.e.  $|f'| \gg 1$  on  $I \setminus [-\varepsilon, \varepsilon]$ ), with  $a > 0$ ,  $0 < \alpha < 1$ ,  $\beta > 0$  and  $\varepsilon > 0$ . It was shown that there exists a positive Lebesgue measure subset  $P$  of parameters in  $(-\varepsilon, \varepsilon)$  such that for  $\mu \in P$  the map  $f_\mu$  is non-uniformly expanding and has slow recurrence to the non-flat infinite and denumerable singular set. Moreover for the same parameters the decay rate of the tail set is exponential, i.e. (9) is true and hence  $f_\mu$  for  $\mu \in P$  is in the setting of Corollaries C and E.

Analogous results hold for the quadratic family  $Q_a(x) = a - x^2$  (and also for general  $C^2$  unimodal families), so that Corollaries C and E apply to quadratic maps for a positive Lebesgue measure subset of parameters.

**2.2. Piecewise smooth one-dimensional expanding maps.** Let  $f : I \rightarrow I$  be a map admitting a sequence  $\mathcal{S} = \{a_n, n \geq 1\} \subset I = [-1, 1]$  such that for every connected component  $G$  of  $I \setminus \mathcal{S}$  we have that  $f|_G$  is  $C^1$  diffeomorphism with its image. Assume that  $\mathcal{S}$  is a non-flat singular set for  $f$  and that  $f$  admits a absolutely continuous ergodic invariant probability measure  $\mu$  with positive Lyapunov exponent and such that  $\log d(x, \mathcal{S})$  is  $\mu$ -integrable and  $\text{supp } \mu = I$ . Then  $f$  is in the setting of Theorem B.

Examples of this kind of maps are the Gauss map [44], and transitive piecewise one dimensional maps satisfying the conditions in [41] (see also [44]), that is there exists  $\kappa > 0$  such that for every connected component  $G$  of  $I \setminus \mathcal{S}$  we also have

$$\text{var}_G \frac{1}{|f'|} \leq \kappa \cdot \sup_G \frac{1}{|f'|} \quad \text{and} \quad \sum_G \sup_G \frac{1}{|f'|} \leq \kappa.$$

More concrete examples are Lorenz-like maps [26, 44] (even with criticalities [28]) and the maps introduced by Rovella [38, 30].

A proof of the exponential decay of the tail set for this class of maps is not available in the literature to the best of our knowledge but can be done as an application of the technique of exclusion of parameters introduced in [10] (the details will appear in forthcoming work), so that Corollaries C and E also hold for this type of maps.

**2.3. Non-uniformly expanding local diffeomorphisms.** Consider a local diffeomorphism  $f : M \rightarrow M$ , so that  $S = \emptyset$ , which satisfies

- $\|(Df)^{-1}\| \leq 1$  and
- $K_1 = \{x \in M : \|Df(x)^{-1}\| = 1\}$  is finite.

Then by the results in [9] we have that such  $f$  has a finite set  $\mathbb{E}$  of equilibrium states for  $\phi$ . Hence in this case Theorem B holds for every continuous function  $\varphi : M \rightarrow \mathbb{R}$ .

**2.4. Viana maps.** The following family of endomorphisms of the cylinder was introduced by Viana in [43]. Let  $a_0 \in (1, 2)$  be such that the critical point  $x = 0$  is pre-periodic for the quadratic map  $Q(x) = a_0 - x^2$ . Let  $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$  and  $b : \mathbb{S}^1 \rightarrow \mathbb{R}$  be a Morse function, for instance  $b(s) = \sin(2\pi s)$ . For fixed small  $\alpha > 0$ , consider

$$\begin{aligned} \hat{f} : \mathbb{S}^1 \times \mathbb{R} &\longrightarrow \mathbb{S}^1 \times \mathbb{R} \\ (s, x) &\longmapsto (\hat{g}(s), \hat{q}(s, x)) \end{aligned}$$

where  $\hat{g}$  is the uniformly expanding map of the circle defined by  $\hat{g}(s) = d \cdot s \pmod{\mathbb{Z}}$  for some  $d \geq 16$ , and  $\hat{q}(s, x) = a(s) - x^2$  with  $a(s) = a_0 + \alpha b(s)$ . For  $\alpha > 0$  small enough there exists an interval  $I \subset (-2, 2)$  such that  $\hat{f}(S^1 \times I)$  is contained in the interior of  $S^1 \times I$ . Hence any map  $f$  sufficiently  $C^0$  close to  $\hat{f}$  has  $S^1 \times I$  as a forward invariant region. We consider from here on these maps  $f$  close to  $\hat{f}$  restricted to  $\mathbb{S}^1 \times I$ .

In [43, 2, 3] a  $C^3$  neighborhood  $\mathcal{U}$  of  $\hat{f}$  was studied and it was proved that every  $f \in \mathcal{U}$  is non-uniformly expanding and has slow recurrence to the non-flat critical set  $\mathcal{C}$ . Hence every  $f \in \mathcal{U}$  is in the setting of Theorem B. Results in [7, 21] show that the tail set decays at least sub-exponentially fast, which is not enough to ensure that Corollaries C and E are true for the maps in  $\mathcal{U}$ . It is conjectured that the tail set indeed decays exponentially fast and with a uniform rate for all maps in  $\mathcal{U}$ .

## 2.5. Partially hyperbolic non-uniformly expanding diffeomorphisms.

We sketch the construction of a robust class of partially hyperbolic non-uniformly expanding diffeomorphisms, taking  $U$  equal to  $M$ , following [6]. This construction is closely related to the  $C^1$  open classes of transitive non-Anosov diffeomorphisms presented in [16, Section 6], as well as other robust examples from [29].

Start with a linear Anosov diffeomorphism  $\hat{f}$  on the  $d$ -dimensional torus  $M = \mathbb{T}^d$ ,  $d \geq 2$ . Write  $TM = E \oplus F$  the corresponding hyperbolic decomposition of the tangent bundle. Let  $V$  be a small closed domain in  $M$  for which there exist unit open cubes  $K^0$  and  $K^1$  in  $\mathbb{R}^d$  such that  $V \subset \pi(K^0)$  and  $\hat{f}(V) \subset \pi(K^1)$ , where  $\pi : \mathbb{R}^d \rightarrow \mathbb{T}^d$  is the canonical projection. Let now  $f$  be a diffeomorphism on  $\mathbb{T}^d$  such that

- (A)  $f$  admits invariant cone fields  $C_E$  and  $C_F$ , with small width  $a > 0$  and containing, respectively, the stable bundle  $E$  and the unstable bundle  $F$  of  $\hat{f}$ ;
- (B)  $f$  is *partially hyperbolic and volume expanding along the center-unstable direction*: there is  $\sigma_1 > 1$  so that

$$|\det(Df | T_x \mathcal{D}_F)| > \sigma_1 \quad \text{and} \quad \|Df | T_x \mathcal{D}_E\| < \sigma_1^{-1}$$

for any  $x \in M$  and any disks  $\mathcal{D}_F, \mathcal{D}_E$  tangent to  $C_F, C_E$ , respectively (see Subsection 3.2 for more on invariant cone fields and disks tangent to cone fields in this setting).

- (C)  $f$  is  $C^1$ -close to  $\tilde{f}$  in the complement of  $V$ , so that there exists  $\sigma_2 < 1$  satisfying

$$\|(Df | T_x \mathcal{D}_F)^{-1}\| < \sigma_2 \quad \text{and} \quad \|Df | T_x \mathcal{D}_E\| < \sigma_2$$

for any  $x \in (M \setminus V)$  and any disks  $\mathcal{D}_F, \mathcal{D}_E$  tangent to  $C_F, C_E$ , respectively. Moreover  $f(V)$  is also contained in the projection of a unit open cube.

- (D) there exist some small  $\delta_0 > 0$  satisfying

$$\|(Df | T_x \mathcal{D}_F)^{-1}\| < 1 + \delta_0$$

for any  $x \in V$  and any disk  $\mathcal{D}_F$  tangent to  $C_F$ .

If  $\tilde{f}$  is a torus diffeomorphism satisfying (A), (B), (D), and coinciding with  $\hat{f}$  outside  $V$ , then any map  $f$  in a  $C^1$  neighborhood of  $\tilde{f}$  satisfies all the previous conditions. Results in [6, Appendix] show in particular that for any  $f$  satisfying (A)–(D) there exist  $c_u > 0$  such that  $f$  is partially hyperbolic and non-uniformly expanding along its center-unstable direction, as defined in Subsection 1.2. Hence on a small  $C^2$  neighborhood  $\mathcal{U}$  of  $\tilde{f}$  every diffeomorphism  $f \in \mathcal{U}$  satisfies all the conditions of Theorem D.

### 3. HYPERBOLIC TIMES

The main technical tool used in the study of non-uniformly expanding maps is the notion of hyperbolic times, introduced in [37, 2]. We say that  $n$  is a  $(\sigma, \delta, b)$ -hyperbolic time of  $f$  for a point  $x$  if the following two conditions hold with  $0 < \sigma < 1$  and  $b, \delta > 0$

$$\prod_{j=n-k}^{n-1} \|Df(f^j(x))^{-1}\| \leq \sigma^k \quad \text{and} \quad d_\delta(f^k(x), \mathcal{S}) \geq e^{-bk} \quad (11)$$

for all  $k = 0, \dots, n-1$ .

We now outline the properties of these special times. For detailed proofs see [6, Proposition 2.8] and [3, Proposition 2.6, Corollary 2.7, Proposition 5.2].

**Proposition 3.1.** *There are constants  $C_1, \delta_1 > 0$  depending on  $(\sigma, \delta, b)$  and  $f$  only such that, if  $n$  is  $(\sigma, \delta, b)$ -hyperbolic time of  $f$  for  $x$ , then there are hyperbolic pre-balls  $V_k(x)$  which are neighborhoods of  $f^{n-k}(x)$ ,  $k = 1, \dots, n$ , such that*

- (1)  $f^k | V_k(x)$  maps  $V_k(x)$  diffeomorphically to the ball of radius  $\delta_1$  around  $f^n(x)$ ;

- (2) for every  $1 \leq k \leq n$  and  $y, z \in V_k(x)$

$$d(f^{n-k}(y), f^{n-k}(z)) \leq \sigma^{k/2} \cdot d(f^n(y), f^n(z));$$

- (3) for  $y, z \in V_k(x)$

$$\frac{1}{C_1} \leq \frac{|\det Df^{n-k}(y)|}{|\det Df^{n-k}(z)|} \leq C_1.$$

The following ensures existence of infinitely many hyperbolic times for Lebesgue almost every point for non-uniformly expanding maps with slow recurrence to the singular set. A complete proof can be found in [6, Section 5].

**Theorem 3.2.** *Let  $f : M \rightarrow M$  be a  $C^{1+\alpha}$  local diffeomorphism away from a non-flat singular set  $\mathcal{S}$ , for some  $\alpha \in (0, 1)$ , non-uniformly expanding and with slow recurrence to  $\mathcal{S}$ . Then there are  $\sigma \in (0, 1)$ ,  $\delta, b > 0$  and there exists  $\theta = \theta(\sigma, \delta, b) > 0$  such that Leb-a.e.  $x \in M$  has infinitely many  $(\sigma, \delta, b)$ -hyperbolic times. Moreover if we write  $0 < n_1 < n_2 < n_3 < \dots$  for the hyperbolic times of  $x$  then their asymptotic frequency satisfies*

$$\liminf_{N \rightarrow \infty} \frac{\#\{k \geq 1 : n_k \leq N\}}{N} \geq \theta \quad \text{for Leb-a.e. } x \in M.$$

### 3.1. Coverings by hyperbolic pre-balls.

**Lemma 3.3.** *Let  $B \subset M$ ,  $\theta > 0$  and  $g : M \rightarrow M$  be a local diffeomorphism outside a non-flat exceptional set  $\mathcal{S}$  such that  $g$  has density  $> \theta$  of hyperbolic times for every  $x \in B$ . Then, given any probability measure  $\nu$  on  $B$  and any  $m \geq 1$ , there exists  $n > m$  such that*

$$\nu(\{x \in B : n \text{ is a hyperbolic time of } g \text{ for } x\}) > \frac{\theta}{2}.$$

This is [33, Lemma 4.4] easily adapted to our setting. For completion we include its very short proof. This lemma shows that we can translate the density of hyperbolic times into the Lebesgue measure of the set of points which have a specific (large) hyperbolic time.

*Proof.* Let  $H$  be the set of pairs  $(x, n) \in B \times \mathbb{N}$  for which  $n$  is a hyperbolic time of  $g$  for  $x$ . For each  $k \geq 1$ , let  $\#_k$  be the normalized counting measure on  $\{m+1, m+2, \dots, m+k\}$ . Our assumption implies that for any given  $x \in B$  we have for big enough  $k \geq 1$

$$\#_k(H \cap (\{x\} \times \mathbb{N})) > 2\theta.$$

Given any probability measure  $\nu$  on  $B$ , fix  $k \geq 1$  large enough so that the above holds for  $C \subset B$  with  $\nu(C) > 1/2$ . By Fubini's Theorem

$$(\nu \times \#_k)(H) > \theta \text{ and thus } \nu(H \cap (B \times \{n\})) > \frac{\theta}{2}$$

for some  $m < n \leq m+k$ . This proves the lemma.  $\square$

Let  $f$  be a regular map in the setting of the Main Theorem with positive density of  $(\sigma, \delta)$ -hyperbolic times for Lebesgue almost everywhere. Let  $\mathcal{E} = \{B(x_i, \delta_1/8), i = 1, \dots, l\}$  be a finite open cover of  $M$  by  $\delta_1/8$ -balls. From this we define a finite partition  $\mathcal{P}$  of  $M$  as follows. We start by setting  $P_1 = B(x_1, \delta_1/8)$  as the first element of the partition. Then, assuming that  $P_1, \dots, P_k$  are already defined we set  $P_{k+1} = B(x_{k+1}, \delta_1/8) \setminus (P_1 \cup \dots \cup P_k)$  for  $k = 1, \dots, l-1$ . Note that if  $P_k \neq \emptyset$  then  $P_k$  has non-empty interior, diameter smaller than  $\delta_1/4$  and the boundary  $\partial P_k$  is a (finite) union of pieces of boundaries of balls in a Riemannian manifold, thus has zero Lebesgue measure. We define  $\mathcal{P}$  by the elements  $P_k$  constructed above which are non-empty.

Note that since  $f$  is regular the boundary of  $g(P)$  still has zero Lebesgue measure for every atom  $P \in \mathcal{P}$  and every inverse branch  $g$  of  $f^n$ , for any  $n \geq 1$ .

Let us choose one interior point in each atom  $P \in \mathcal{P}$  and form the set  $\mathcal{C}_0$  of representatives of the atoms of  $\mathcal{P}$ . Let  $d_0 = \min\{d(w, \partial\mathcal{P}), w \in \mathcal{C}_0\} > 0$  where  $\partial\mathcal{P} = \cup_{P \in \mathcal{P}} \partial P$  is the boundary of  $\mathcal{P}$ .

**Lemma 3.4.** *Let  $(\mu_n)_{n \geq 1}$  be a family of Borel probability measures on  $M$  and  $\mu$  some weak\* accumulation point of the sequence  $(\mu_n)$ . Then given  $0 < \varepsilon < d_0$  there exists a partition  $\mathcal{P}_\varepsilon$  with the same number of atoms of  $\mathcal{P}$ , whose atoms have non-empty interior, diameter smaller than  $\delta_1/2$  and whose boundaries have zero Lebesgue measure, such that*

- (1)  $\mu(\partial\mathcal{P}_\varepsilon) = 0$  and  $\mu_n(\partial\mathcal{P}_\varepsilon) = 0$  for all  $n \geq 1$ ;
- (2) each  $P \in \mathcal{P}_\varepsilon$  contains one, and only one, element of  $\mathcal{C}_0$ ;
- (3) given  $\delta > 0$  there is  $\varepsilon > 0$  small enough such that for each  $P \in \mathcal{P}_\varepsilon$  there is  $Q \in \mathcal{P}$  such that  $\text{Leb}(P \triangle Q) < \varepsilon < \delta \cdot \text{Leb}(Q)$ .

*Proof.* Let us take  $0 < \gamma < \min\{\varepsilon, \delta_1/8\}$  such that for all  $i = 1, \dots, l$

$$\text{Leb}\left(B(x_i, \frac{\delta_1}{8} + \gamma) \setminus B(x_i, \frac{\delta_1}{8})\right) < \frac{\varepsilon}{l} \quad (12)$$

and also for all  $n \geq 1$

$$\mu(\partial B(x_i, \frac{\delta_1}{8} + \gamma)) = 0 = \mu_n(\partial B(x_i, \frac{\delta_1}{8} + \gamma)). \quad (13)$$

Such value of  $\gamma$  exists since the set of values of  $\gamma > 0$  such that some of the expressions in (13) is positive for some  $i \in \{1, \dots, l\}$  and some  $n \geq 1$  is denumerable. Thus we may take  $\gamma > 0$  satisfying (13) arbitrarily close to zero, and so inequality (12) can also be obtained.

We consider now the finite open cover  $\mathcal{E}_\gamma = \{B(x_i, \delta_1/8 + \gamma), i = 1, \dots, l\}$  of  $M$  and construct the partition  $\mathcal{P}_\gamma$  induced by  $\mathcal{E}_\gamma$  by the same procedure as before. Since  $\gamma < \varepsilon < d_0$  we obtain  $d(w, \partial B(x_i, \delta_1/8 + \gamma)) \geq d_0 - \gamma > 0$  for all  $i = 1, \dots, l$  and every  $w \in \mathcal{C}_0$ . This shows that each  $w \in \mathcal{C}_0$  is contained in some atom  $P_w$  of  $\mathcal{P}_\gamma$ . Moreover there cannot be distinct  $w_1, w_2 \in \mathcal{C}_0$  such that  $w_2 \in P_{w_1}$ , because this would mean that for some  $i \in \{1, \dots, l\}$  we have  $w_2 \in B(x_i, \delta_1/8)$ ,  $w_1 \notin B(x_i, \delta_1/8)$  and  $w_1, w_2 \in B(x_i, \delta_1/8 + \gamma)$ , which contradicts the choice of  $\gamma < d_0$ .

Let us consider  $\{P_w, w \in \mathcal{C}_0\}$ . There might be other (finitely many) atoms  $P$  in  $\mathcal{P}_\gamma$  and, if so, we join them to some adjacent atom  $P_w$  (meaning  $\overline{P} \cap \overline{P_w} \neq \emptyset$ ) obtaining a new atom  $P \cup P_w$ . In this way we obtain a partition  $\mathcal{P}_\varepsilon$  with as many atoms as the elements of  $\mathcal{C}_0$  and satisfying items (1) and (2) of the statement of the lemma.

Clearly for any  $w \in \mathcal{C}_0$  the corresponding atoms  $P_w \in \mathcal{P}_\varepsilon$  and  $Q_w \in \mathcal{P}$  satisfy

$$\text{Leb}(P_w \triangle Q_w) \leq \sum_{i=1}^l \text{Leb}\left(B(x_i, \frac{\delta_1}{8} + \gamma)\right) < l \cdot \frac{\varepsilon}{l} = \varepsilon$$

and  $\text{diam}(P_w) \leq 2(\delta_1/8 + \gamma) < \delta_1/2$ . Since  $\mathcal{P}$  is a finite partition with  $\text{Leb}(\partial\mathcal{P}) = 0$  we have  $\iota = \min\{\text{Leb}(P) : P \in \mathcal{P}\} > 0$  and so given  $\delta > 0$  and

taking  $\varepsilon < \min\{\iota \cdot \delta, d_0\}$  we get

$$\text{Leb}(P_w \triangle Q_w) < \varepsilon = \iota \cdot \frac{\varepsilon}{\iota} < \iota \cdot \delta \leq \delta \cdot \text{Leb}(Q_w).$$

The proof is complete.  $\square$

Having this we can now obtain the following flexible covering lemma with hyperbolic pre-balls which will enable us to approximate the Lebesgue measure of a given set through the measure of families of hyperbolic pre-balls.

**Lemma 3.5.** *Let a measurable set  $E \subset M$ ,  $m \geq 1$  and  $\varepsilon > 0$  be given. Let  $\theta > 0$  be a lower bound for the density of hyperbolic times for Lebesgue almost every point. Then there are integers  $m < n_1 < \dots < n_k$  for  $k = k(\varepsilon) \geq 1$  and families  $\mathcal{E}_i$  of subsets of  $M$ ,  $i = 1, \dots, k$  such that*

- (1)  $\mathcal{E}_1 \cup \dots \cup \mathcal{E}_k$  is a finite pairwise disjoint family of subsets of  $M$ ;
- (2)  $n_i$  is a  $(\sigma/2, \delta/2)$ -hyperbolic time for every point in  $P$ , for every element  $P \in \mathcal{E}_i$ ,  $i = 1, \dots, k$ ;
- (3) every  $P \in \mathcal{E}_i$  is the pre-image of some element  $Q \in \mathcal{P}$  under an inverse branch of  $f^{n_i}$ ,  $i = 1, \dots, k$ ;
- (4) there is an open set  $U_1 \supset E$  containing the elements of  $\mathcal{E}_1 \cup \dots \cup \mathcal{E}_k$  with  $\text{Leb}(U_1 \setminus E) < \varepsilon$ ;
- (5)  $\text{Leb}\left(E \triangle \bigcup_i \mathcal{E}_i\right) \leq \left(1 - \frac{\theta}{4}\right)^k < \varepsilon$ .

The proof follows [33, Lemma 8.2] closely. We write  $\mathcal{C}_m$  the set of pairs  $(z, n_i)$  where  $f^{n_i}(z) = w \in \mathcal{C}_0$  and  $z \in P$  for all  $P \in \mathcal{E}_i$  and  $i = 1, \dots, k$  (such  $z$  exist by item (3) of Lemma 3.5).

**Remark 3.6.** Note that  $k$  depends on  $\varepsilon$  only and not on the set  $E$ .

*Proof.* By the non-uniformly expanding assumption on  $f$  we know that there exists  $\theta > 0$  such that Lebesgue almost every point has density  $> \theta$  of hyperbolic times of  $f$ .

Let  $U_1$  be an open set and  $K_1$  a compact set such that  $K_1 \subset E \subset U_1$  and  $\text{Leb}(U_1 \setminus K_1) < \varepsilon$  and  $\text{Leb}(K_1) > (1/2)\text{Leb}(U_1)$ . Using Lemma 3.3 with  $B = K_1$  and  $\nu = \text{Leb}/\text{Leb}(K_1)$  we can find  $n_1 > m$  such that  $e^{-cn_1} < d(K_1, M \setminus U_1)$  and the subset  $L_1$  of points of  $K_1$  for which  $n_1$  is a hyperbolic time satisfies  $\text{Leb}(L_1) \geq \frac{\theta}{2}\text{Leb}(K_1) \geq \frac{\theta}{4}\text{Leb}(E)$ .

Given  $x \in L_1$  let  $g : B(f^{n_1}(x), \delta_1) \rightarrow V_{n_1}(x)$  be the inverse branch of  $f^{n_1} | V_{n_1}(x)$ , recall that  $n_1$  is a hyperbolic time for  $x$  and see Proposition 3.1. By the choice of  $\mathcal{P}$  there exists a unique  $P \in \mathcal{P}$  such that  $f^{n_1}(x) \in P$ . Let us consider  $g(P)$  and let  $\mathcal{E}_1$  be the family of all such sets obtained as  $g(P)$  which intersect  $L_1$ , where  $g$  is an inverse branch of  $f^{n_1}$  corresponding to a hyperbolic time and  $P$  is an element of  $\mathcal{P}$ .

Note that the elements of  $\mathcal{E}_1$  are pairwise disjoint because  $\mathcal{P}$  is a partition. Moreover by the properties of hyperbolic times (Proposition 3.1) the diameter of  $P \in \mathcal{E}_1$  is smaller than  $e^{-cn_1}$ . Hence the union  $E_1$  of all the elements of  $\mathcal{E}_1$  is contained in  $U_1$  and by construction

$$\text{Leb}(E_1 \cap E) \geq \text{Leb}(L_1) \geq \frac{\theta}{4}\text{Leb}(E).$$

Now consider the open set  $U_2 = U_1 \setminus \overline{E_1}$  and set  $K_2 \subset E \setminus \overline{E_1}$  a compact set such that  $\text{Leb}(K_2) \geq (1/2)\text{Leb}(E \setminus E_1)$ . Observe that  $\text{Leb}(\overline{E_1} \setminus E_1) = 0$

since  $\partial\mathcal{P}$  has zero Lebesgue measure and this property is preserved under backward iteration by the regularity assumption on  $f$ . Reasoning as before, we can find  $n_2 > n_1$  such that  $e^{-cn_2} < d(K_2, M \setminus U_2)$  and a set  $L_2 \subset K_2$  such that  $\text{Leb}(L_2) \geq (\frac{\theta}{2}) \text{Leb}(K_2)$  and  $n_2$  is a hyperbolic time for every  $x \in L_2$ . Let  $\mathcal{E}_2$  be the family of elements  $g(P)$  which intersect  $L_2$ , where  $P \in \mathcal{P}$  and  $g$  is an inverse branch of  $f^{n_1}$  corresponding to a hyperbolic time.

Again  $\mathcal{E}_2$  is a pairwise disjoint family of sets whose diameters are smaller than  $e^{-cn_2}$ . Thus their union  $E_2$  is contained in  $U_2$ . Hence  $\mathcal{E}_1 \cup \mathcal{E}_2$  is also a pairwise disjoint family and, in addition

$$\text{Leb}(E_2 \cap (E \setminus E_1)) \geq \text{Leb}(L_2) \geq \frac{\theta}{2} \text{Leb}(K_2) \geq \frac{\theta}{4} \text{Leb}(E \setminus E_1).$$

Repeating this procedure we obtain families  $\mathcal{E}_i, i = 1, \dots, k$  of elements of  $\mathcal{P}_{n_i}$  which are pairwise disjoint and contained in  $U_1$ , and

$$\text{Leb}(E_{i+1} \cap (E \setminus (E_1 \cup \dots \cup E_i))) \geq \frac{\theta}{4} \text{Leb}(E \setminus (E_1 \cup \dots \cup E_i)) \quad (14)$$

for all  $i = 1, \dots, k-1$ , for some  $k \geq 1$ , where  $E_j = \cup \mathcal{E}_j$ . Hence

$$\text{Leb}\left(\bigcup_{i=1}^k E_i \setminus E\right) \leq \text{Leb}(U_1 \setminus E) < \varepsilon$$

and (14) ensures that

$$\text{Leb}\left(E \setminus \bigcup_{i=1}^k E_i\right) \leq \left(1 - \frac{\theta}{4}\right)^k \text{Leb}(E).$$

Therefore we can find  $k \geq 1$  such that  $\text{Leb}(E \triangle \bigcup_{i=1}^k \mathcal{E}_i) < \varepsilon$ , as stated.  $\square$

**Remark 3.7.** Note that the construction proving Lemma 3.5 gives a finite sequence of hyperbolic times, open sets  $U_1, \dots, U_k$  and closed sets  $\overline{E}_1, \dots, \overline{E}_k$ . Having these we can find small enough  $\delta > \varepsilon > 0$ , replace  $\mathcal{P}$  in the proof of Lemma 3.5 by any partition  $\mathcal{P}_\varepsilon$  obtained as in Lemma 3.4 (by slightly modifying  $\mathcal{P}$ ), and use the same inverse branches to obtain families  $\mathcal{E}'_i$  of pre-balls such that

$$\text{Leb}\left(\left(\bigcup_i \mathcal{E}_i\right) \triangle \left(\bigcup_i \mathcal{E}'_i\right)\right) \leq \sum_i C_1 \delta \text{Leb}(\mathcal{E}_i) < C_1 \delta \text{Leb}\left(\bigcup_i \mathcal{E}_i\right) \leq C_1 \delta$$

where  $C_1$  is the volume distortion constant (see Proposition 3.1). Hence after the modification of the initial partition we get

$$\text{Leb}\left(E \triangle \bigcup_i \mathcal{E}'_i\right) < \varepsilon + C_1 \delta < (1 + C_1) \delta$$

since  $\varepsilon < \delta$ . Moreover the set  $\mathcal{C}_m$  is unaffected since  $\mathcal{C}_0$  is fixed and the inverse branches are kept.

**3.2. The partially hyperbolic setting.** Here we state the main results needed to obtain an extension of the covering Lemma 3.5 to the setting of partially hyperbolic non-uniformly expanding attracting sets. As we indicate along the way, the proofs of most of them can be found in [6].

**3.2.1. Stable/Unstable cone fields.** Let  $\Lambda$  be a partially hyperbolic and non-uniformly expanding attracting set for a  $C^2$  diffeomorphism  $f : M \rightarrow M$  with a trapping region  $U \subset M$ . The existence of the dominated splitting  $E \oplus F$  of  $T_\Lambda M$  ensures the existence of a continuous extension  $\tilde{E} \oplus \tilde{F}$  of  $E \oplus F$  to a neighborhood of  $\Lambda$ , which we assume without loss to be  $U$ , and of the following cone fields:

**stable cones:**  $\mathbb{E}_x^a = \{(u, v) \in \tilde{E}(x) \oplus \tilde{F}(x) : \|v\| \leq a \cdot \|u\|\};$   
**unstable cones:**  $\mathbb{F}_x^b = \{(u, v) \in \tilde{E}(x) \oplus \tilde{F}(x) : \|u\| \leq b \cdot \|v\|\};$

for all  $x \in U$  and  $a, b \in (0, 1)$ , which are  $Df$ -invariant in the following sense (see e.g. [15, Appendix C])

- if  $x, f^{-1}(x) \in U$ , then  $Df^{-1}(\mathbb{E}_x^a) \subset \mathbb{E}_{f^{-1}(x)}^{\lambda a}$ ;
- if  $x, f(x) \in U$ , then  $Df(\mathbb{F}_x^b) \subset \mathbb{F}_{f(x)}^{\lambda b}$ ;

for some  $\lambda \in (0, 1)$ . Continuity enables us to unambiguously denote  $d_E = \dim(\tilde{E})$  and  $d_F = \dim(\tilde{F})$ , so that  $d = d_E + d_F = \dim(M)$ , and domination guarantees that the angles between the  $\tilde{E}$  and  $\tilde{F}$  directions are bounded from below away from zero at every point.

**3.2.2. Hyperbolic times.** In this setting, given  $\sigma > 1$  we say that  $n$  is a  $\sigma$ -hyperbolic time for  $x \in U$  if

$$\prod_{j=n-k+1}^n \|(Df|_{F_{f^j(x)}})^{-1}\| \leq \sigma^k \quad \text{for all } 1 \leq k \leq n.$$

**Remark 3.8.** This definition of hyperbolic time is entirely analogous to the one given in the local diffeomorphisms setting except that we restrict the derivatives to the  $F$ -direction. Hence the statement and proof of Lemma 3.3 carry over without change.

**3.2.3.  $E$ -disks and  $F$ -disks.** Let us fix the unit balls of dimensions  $d_E, d_F$

$$\mathbb{B}_E = \{w \in \mathbb{R}^{d_E} : \|w\|_2 \leq 1\} \quad \text{and} \quad \mathbb{B}_F = \{w \in \mathbb{R}^{d_F} : \|w\|_2 \leq 1\}$$

where  $\|\cdot\|_2$  is the standard Euclidean norm on the corresponding Euclidean space. We say that a  $C^{1+\alpha}$  embedding  $\Delta : \mathbb{B}_E \rightarrow M$  (respectively  $\Delta : \mathbb{B}_F \rightarrow M$ ) is a  $E$ -disk (resp.  $F$ -disk) if the image of  $D\Delta(w)$  is contained in  $\mathbb{E}_{\Delta(w)}^a$  for all  $w \in \mathbb{B}_E$  (resp.  $D\Delta(w)(\mathbb{R}^{d_F}) \subset \mathbb{F}_{\Delta(w)}^b$  for every  $w \in \mathbb{B}_F$ ), where  $\alpha \in (0, 1)$  is fixed.

**3.2.4. Curvature of  $E$ - and  $F$ -disks at hyperbolic times.** Let  $r_0 > 0$  be an injectivity radius of the exponential map on  $M$ , that is  $\exp_x : B(x, r_0) \rightarrow M$  is a diffeomorphism onto its image  $G(x, r_0) = \exp_x(B(x, r_0))$ , where  $B(x, r_0) = \{v \in T_x M : \|v\| < r_0\}$  is the  $r_0$ -neighborhood of 0 in  $T_x M$ . By the continuity of the splitting  $E \oplus F$  and the cone fields we can choose  $0 < r < \min\{r_0, \delta_1/4\}$  such that for every  $x \in \Lambda$  the subspace  $E_x$  is contained in all the images of the cone field  $\mathbb{E}_x^a$  under the exponential map  $\exp_x$  and analogously for the complementary direction, that is for every  $y \in G(x, r) \cap \Lambda$  we have

$$E_x \subset D(\exp_x^{-1})(\mathbb{E}_y^a) \quad \text{and} \quad F_x \subset D(\exp_x^{-1})(\mathbb{F}_y^b). \quad (15)$$

This ensures that every  $F$ -disk (respectively every  $E$ -disk)  $\Delta$  is such that its image on  $B(x, r)$  given by  $\exp_x^{-1}(\Delta \cap G(x, r))$  is transversal to the direction of  $E_x$  (resp.  $F_x$ ).

The “curvature” of  $E$ - and  $F$ -disks can be determined by the notion of Hölder variation of the tangent bundle as follows.

We write  $\Delta$  also for the image of the respective embedding for every  $E$ - or  $F$ -disk. Hence if  $\Delta$  is a  $E$ -disk and  $y = \Delta(w)$  for some  $w \in \mathbb{B}_E$ , then the tangent space of  $\Delta$  at  $y$  is the graph of a linear map  $A_x(y) : T_x\Delta \rightarrow F(x)$  for  $w \in \Delta^{-1}(V_x)$  (here  $T_x\Delta = D\Delta(x)(\mathbb{R}^{d_E})$ ). The same happens locally for a  $F$ -disk exchanging the roles of the bundles  $E$  and  $F$  above.

The domination condition on the splitting  $E \oplus F$  ensures the existence of  $\zeta \in (0, 1)$  such that for some  $n \geq 1$  and all  $x \in \Lambda$

$$\|Df^n | E_x\| \cdot \|(Df^n | F_x)^{-1}\|^{1+\zeta} \leq \frac{3}{4}.$$

Given  $C > 0$  we say that the *tangent bundle of  $\Delta$  is  $(C, \zeta)$ -Hölder* if

$$\|A_x(y)\| \leq C \operatorname{dist}_\Delta(x, y)^\zeta \quad \text{for all } y \in G(x, r) \cap \Delta \quad \text{and } x \in U, \quad (16)$$

where  $\operatorname{dist}_\Delta(x, y)$  is the *distance along  $\Delta$*  defined by the length of the shortest smooth curve from  $x$  to  $y$  inside  $\Delta$  calculated with respect to the Riemannian norm  $\|\cdot\|$  induced on  $TM$ .

For a  $E$ - or  $F$ -disk  $\Delta \subset U$  we define

$$\kappa(\Delta) = \inf\{C > 0 : T\Delta \text{ is } (C, \zeta)\text{-Hölder}\}. \quad (17)$$

The proof of the following result can be found in [6, Subsection 2.1]. The basic ingredients are the cone invariance and dominated decomposition properties for  $f$ .

**Proposition 3.9.** *There is  $C_2 > 0$  such that given a  $F$ -disk  $\Delta \subset U$*

- (1) *there exists  $n_1 \in \mathbb{N}$  such that  $\kappa(f^n(\Delta)) \leq C_2$  for all  $n \geq n_1$ ;*
- (2) *if  $\kappa(\Delta) \leq C_2$  then  $\kappa(f^n(\Delta)) \leq C_2$  for all  $n \geq 0$ ;*
- (3) *in particular, if  $\Delta$  is as in the previous item, then*

$$J_n : f^n(\Delta) \ni x \mapsto \log |\det(Df | T_x(f^n(\Delta)))|$$

*is  $(L_1, \zeta)$ -Hölder continuous with  $L_1 > 0$  depending only on  $C_2$  and  $f$ , for every  $n \geq 1$ .*

**3.2.5. Distortion bounds.** The following uniform backward contraction and distortion bounds are proved in [6, Lemma 2.7, Proposition 2.8].

**Proposition 3.10.** *There exist  $C_3, \delta_1 > 0$  depending only on  $f, \sigma$  such that, given any  $F$ -disk  $\Delta \subset U$ ,  $x \in \Delta$ , and  $n \geq 1$  a  $\sigma$ -hyperbolic time for  $x$ ,*

- (1)  *$\operatorname{dist}_{f^{n-k}(D)}(f^{n-k}(y), f^{n-k}(x)) \leq \sigma^{k/2} \operatorname{dist}_{f^n(D)}(f^n(y), f^n(x))$ , for all  $y \in \Delta$  with  $\operatorname{dist}(f^n(x), f^n(y)) \leq \delta_1$ ;*
- (2) *if  $\kappa(\Delta) \leq C_2$  then*

$$\frac{1}{C_3} \leq \frac{|\det Df^n | T_y\Delta|}{|\det Df^n | T_x\Delta|} \leq C_3$$

*for every  $y \in \Delta$  such that  $\operatorname{dist}(f^n(y), f^n(x)) \leq \delta_1$ .*

3.2.6. *The initial partition and the covering lemma.* Now we consider the following rectangle

$$\hat{R}(x, s) = \{(u, v) \in T_x M : \|u\| < s, \|v\| < s, u \in E_x, v \in F_x\}$$

where  $s$  is chosen so that  $\hat{R}(x, s) \subset B_x(r)$  for all  $x \in \Lambda$ . This defines an open cover  $\{\exp_x(\hat{R}(x, s))\}_{x \in \Lambda}$  of  $\Lambda$  which admits a finite subcover denoted by  $\{R_1 = R(x_1, s), \dots, R_h = R(x_h, s)\}$ . This finite cover will define the initial partition  $\mathcal{P}$  given as before by

$$\mathcal{P} = \{R_1, M \setminus R_1\} \vee \dots \vee \{R_h, M \setminus R_h\}.$$

We may assume without loss that  $\text{Leb}(\partial \mathcal{P}) = 0$  by slightly changing the initial cover. We choose an interior point in each element of  $\mathcal{P}$  which together define the set  $\mathcal{C}_0$ .

Now we adapt the covering Lemma 3.5 to the setting of partially hyperbolic non-uniformly expanding attracting sets as follows.

**Lemma 3.11.** *Let a measurable set  $E \subset U$ ,  $m \geq 1$  and  $\varepsilon > 0$  be given. Let  $\theta > 0$  be a lower bound for the density of hyperbolic times for Lebesgue almost every point on  $U$ . Then there are integers  $m < n_1 < \dots < n_k$  for  $k = k(\varepsilon) \geq 1$ , and families  $\mathcal{E}_i$  of subsets of  $M$ ,  $i = 1, \dots, k$  such that*

- (1)  $\mathcal{E}_1 \cup \dots \cup \mathcal{E}_k$  is a finite family of subsets of  $M$  and each  $\mathcal{E}_i$  is a pairwise disjoint family;
- (2)  $n_i$  is a  $(\sigma/2, \delta/2)$ -hyperbolic time for every point in  $P$ , for every element  $P \in \mathcal{E}_i$ ,  $i = 1, \dots, k$ ;
- (3) every  $P \in \mathcal{E}_i$  is the pre-image of some element  $Q \in \mathcal{P}$  under  $f^{-n_i}$ ,  $i = 1, \dots, k$ ;
- (4)  $\text{Leb}\left(E \setminus \bigcup_i \mathcal{E}_i\right) \leq \left(1 - \frac{\theta}{4}\right)^k < \varepsilon$ .

*Proof.* Let  $E \subset U$ ,  $\varepsilon > 0$  and  $m \geq 1$  be given. Set  $\nu = \text{Leb} / \text{Leb}(E)$  and apply Lemma 3.3 with  $B = E$  to obtain  $n_1 > m$  and  $L_1 \subset E$  such that  $n_1$  is a hyperbolic time for every point  $x \in L_1$  and  $\text{Leb}(L_1) \geq \frac{\theta}{2} \text{Leb}(E)$ .

Given  $x \in L_1$  let  $P_x$  be the unique element of the partition  $f^{-n_1} \mathcal{P}$  which contains  $x$  (recall that  $f$  is a diffeomorphism). Define  $\mathcal{E}_1 = \{P_x : x \in L_1\}$ . Then  $\mathcal{E}_1$  is a finite pairwise disjoint family of preimages of elements of  $\mathcal{P}$  corresponding to a hyperbolic time  $n_1$ . If  $E_1$  is the union of the elements of  $\mathcal{E}_1$ , then

$$\text{Leb}(E_1 \cap E) \geq \text{Leb}(L_1) \geq \frac{\theta}{2} \text{Leb}(E).$$

Now consider  $\hat{E}_2 = E \setminus \overline{E_1}$ . If  $\text{Leb}(\hat{E}_2) < \varepsilon$  then we are done, since then  $\text{Leb}(E \setminus E_1) < \varepsilon$  because  $\text{Leb}(\partial \mathcal{E}_1) = 0$  as  $f$  is regular map. Otherwise use again Lemma 3.3 to find  $n_2 > n_1$  and  $L_2 \subset \hat{E}_2$  such that  $n_2$  is a hyperbolic time for all points of  $L_2$  and  $\text{Leb}(L_2) \geq \frac{\theta}{2} \text{Leb}(\hat{E}_2)$ .

Let  $\mathcal{E}_2$  be the family of all elements of the partition  $f^{-n_2} \mathcal{P}$  which intersect  $\hat{E}_2$ . Then  $\mathcal{E}_2$  is a pairwise disjoint family and the union  $E_2$  of its elements satisfies

$$\text{Leb}(E_2 \cap (E \setminus E_1)) \geq \text{Leb}(L_2) \geq \frac{\theta}{2} \text{Leb}(\hat{E}_2) \geq \frac{\theta}{4} \text{Leb}(E \setminus E_1).$$

Repeating this procedure we get families  $\mathcal{E}_i, i = 1, \dots, k$  of elements of  $f^{-n_i}\mathcal{P}$  with  $m < n_1 < \dots < n_k$  satisfying the inequality (14). These families satisfy items (1)-(3) by construction and item (4) follows by (14) as in the proof of Lemma 3.5. This concludes the proof.  $\square$

Observe that we may apply Lemma 3.4 to  $\mathcal{P}$  to ensure that, for a given denumerable family of  $f$ -invariant probability measures, there is a partition  $\mathcal{P}_\varepsilon$  arbitrarily close to  $\mathcal{P}$ , with the same number of elements, such that the measure of the boundary of the elements of  $\mathcal{P}_\varepsilon$  is zero with respect to all measures of the family. Moreover as in the previous subsection, we write  $\mathcal{C}_m$  the set of pairs  $(z, n_i)$  where  $f^{n_i}(z) = w \in \mathcal{C}_0$  and  $z \in P$  for all  $P \in \mathcal{E}_i$  and  $i = 1, \dots, k$ . In addition, we can build the new partition  $\mathcal{P}_\varepsilon$  in such a way that the sets  $\mathcal{C}_n$  are unchanged.

**3.3. The volume of dynamical balls.** Here we show that the volume of dynamical balls on hyperbolic times is well controlled by  $S_n J$ , either in the local diffeomorphism case with or without singularities, or in the partially hyperbolic case.

**3.3.1. The local diffeomorphism case with singularities.** Note that by the properties of bounded distortion of volumes during hyperbolic times (item 3 of Proposition 3.1) we can write, if  $n$  is a hyperbolic time of  $f$  for  $x \in M$

$$\begin{aligned} \text{Leb}(B(f^k(x), n-k, \delta_1)) &= \int_{B(f^k(x), n-k, \delta_1)} \frac{dz}{|\det Df^{n-k}(z)|} \\ &\leq C_1 \frac{\text{Leb}(B(f^n(x), \delta_1))}{|\det Df^{n-k}(x)|}, \end{aligned}$$

then recalling that  $J = \log |\det Df|$  we get

$$\begin{aligned} \text{Leb}(B(f^k(x), n-k, \delta_1)) &\leq C_1 e^{-S_{n-k}J(f^k(x))} \text{Leb}(B(f^n(x), \delta_1)) \\ &\leq C_1 e^{-S_{n-k}J(f^k(x))}. \end{aligned}$$

Observe that by Proposition 3.1 if  $n$  is a hyperbolic time of  $f$  for  $x$  we get due to uniform backward contraction

$$S_{n-k}J(f^k(x)) = \log |\det Df^{n-k}(x)| \geq (n-k) \cdot \dim(M) \log \sigma/2 > 0$$

which will be used several times in what follows.

**3.3.2. The partially hyperbolic case with non-uniform expansion.** In the partially hyperbolic and non-uniformly expanding setting we recall the construction of the cover  $R_1, \dots, R_h$  and the initial partition  $\mathcal{P}$  from Subsection 3.2. Observe that if we take  $\delta_0$  to be the Lebesgue number of the covering  $R_1, \dots, R_h$  (see e.g. [32]), then for all  $0 < \delta < \delta_0$  we have for all  $x \in U$  and  $n \geq 1$  a hyperbolic time for  $x$

$$B(x, n, \delta) \subset f^{-n}\mathcal{P}(x),$$

where  $f^{-n}\mathcal{P}(x)$  denotes the element of  $f^{-n}\mathcal{P}$  which contains  $x$ . To find an upper bound for the volume of this dynamical ball it is enough to estimate the volume of  $f^{-n}\mathcal{P}(x)$  when  $n$  is a hyperbolic time for  $x$ .

Let  $P \in \mathcal{P}$  be such that  $f^{-n}(P)$  has a positive Lebesgue measure subset  $\tilde{P}$  of points for which  $n$  is a hyperbolic time and choose  $h$  such that  $R_h \supset P$ .

Let  $\tilde{Q} \in \mathcal{P}$  be such that  $Q = \tilde{Q} \cap \tilde{P}$  has positive Lebesgue measure and choose  $l$  such that  $R_l \supset Q$ .

We consider the projection of  $\hat{P} = \exp_{x_l}^{-1}(\tilde{P})$  on  $E_{x_l}$  parallel to  $F_{x_l}$ . Its diameter will be bounded a constant which is a function of  $f$  and  $s$  only, since the number of different  $R_l$  is finite. Projecting  $\tilde{Q}$  on the complementary direction  $F_{x_l}$  parallel to  $E_{x_l}$  we may use the backward contraction and bounded area distortion for hyperbolic times along  $F$ -disks to estimate the area along  $F$ -disks and integrate to deduce a volume estimate.

Indeed, observe that since the  $E$  direction is uniformly contracted by  $Df$ , if we fix a point  $x_0 \in Q$ , the corresponding point  $x_n = f^n(x_0) \in P \cap f^n(Q)$  and a  $E$ -disk  $\gamma$  which crosses  $R_h$ , then the connected component  $\hat{\gamma}$  of  $f^{-n}(\gamma) \cap R_l$  containing  $x_0$  is a  $E$ -disk which also crosses  $R_l$ . Moreover distances along  $\gamma$  are uniformly expanded by  $f^{-1}$ . Thus every point  $w_0 \in \hat{\gamma}$  is such that  $w_k = f^k(w_0)$  and  $x_k = f^k(x_0)$  satisfy

$$C \frac{\delta_1}{4} > Cs \geq \text{dist}(w_0, x_0) \geq C\lambda^{-k} \text{dist}(w_k, x_k), \quad (18)$$

for some constant  $C > 0$  depending on  $f$  only. Hence if we take  $s$  small enough then we can ensure that  $w_k$  is close enough to  $x_k$  for  $k = 1, \dots, n$  so that  $n$  is also an hyperbolic time for all  $w_0 \in \hat{\gamma}$ . Thus we can consider  $F$ -disks  $\beta_q$  through the points  $q$  of  $Q$  parallel to  $F$ , which are transversal to  $\hat{\gamma}$ . Then the images  $f^n(\beta_q)$  will be  $F$ -disks crossing  $R_l$  which together cover  $P \cap f^n(Q)$ , see Figure 1.




FIGURE 1. The diameter of the elements of  $\mathcal{E}_n$  through the use of  $E$ -disks and images of  $F$ -disks on a hyperbolic time.

The preimages  $f^{-n}(P \cap f^n(Q) \cap f^n(\beta_q))$  then form a cover of  $Q$  and these predisks are  $F$ -disks whose diameter is smaller than  $e^{-cn}$ .

Using Tonelli's Theorem, the Change of Variables Formula and the bounded area distortion along hyperbolic times in the partially hyperbolic setting given by Proposition 3.10, together with the bounded curvature of the images of  $F$ -disks given by Proposition 3.9, we arrive at

$$\begin{aligned} \text{Leb}(Q) &= \int_{\hat{\gamma}} m(Q \cap \beta_q) dq \\ &\leq \int_{\hat{\gamma}} C_3 e^{-S_n J(q)} m(f^n(Q) \cap f^n(\beta_q)) dq, \end{aligned}$$

where  $m$  denotes the  $d_F$ -dimensional Lebesgue measure induced by Leb on  $F$ -disks. But by (18) we see that every  $q \in \hat{\gamma} \cap Q$  satisfies for  $k = 0, \dots, n$

$$d(f^k(q), f^k(x)) \leq C\lambda^k \frac{\delta_1}{4}.$$

Hence because  $J$  is at least  $C^{1+\alpha}$  for some  $\alpha \in (0, 1)$  with Hölder constant  $C > 0$  (in fact we can take  $\alpha = 1$  if  $f$  is  $C^2$ ) the usual bounded distortion argument provides a constant  $C_0 > 0$  such that

$$\log \frac{|\det Df^n|_{F_q}|}{|\det Df^n|_{F_x}} = \sum_{j=0}^{n-1} \log \frac{|\det Df(f^j(q))|}{|\det Df(f^j(x))|} \leq \sum_{j=0}^{n-1} C d(f^j(q), f^j(x))^\alpha \leq C_0.$$

Thus  $|S_n J(q) - S_n J(x)| \leq C_0$  and by the above integration estimates we get

$$\text{Leb}(Q) \leq \int_{\hat{\gamma}} C_3 e^{C_0} e^{-S_n J(x)} m(f^n(Q) \cap f^n(\beta_q)) dq \leq C' e^{-S_n J(x)},$$

where  $C'$  is bounded by the  $d_E$ -dimensional area  $A_E$  of  $\hat{\gamma}$  (which is a function of  $s < \delta_1/4$ ) times a uniform bound  $A_F$  for the  $d_F$ -dimensional area of  $f^n(\beta_q)$  (which is a function of the curvature bound  $C_2$  and of  $\delta_1$ ) multiplied by the bounded distortion constants, that is  $C' \leq C_3 e^{C_0} A_E A_F$ .

This shows that we have the same kind of estimate for the volume of a dynamical ball as in the local diffeomorphism case, except for a different distortion constant and the fact that the Jacobian is calculated along the  $F$  direction.

#### 4. HYPERBOLIC TIMES AND LARGE DEVIATIONS

The statements of the main theorems and corollaries are consequences of the following more abstract result.

**Theorem 4.1.** *Let  $f : M \rightarrow M$  be a local diffeomorphism outside a non-flat singular set  $\mathcal{S}$  admitting  $\sigma \in (0, 1)$  and  $b, \delta > 0$  such that Lebesgue almost every point has positive density of  $(\sigma, \delta, b)$ -hyperbolic times. Then given  $c \in \mathbb{R}$  and a continuous function  $\varphi : M \rightarrow \mathbb{R}$  items (1)-(3) of Theorem A hold.*

Clearly Theorem A follows from Theorem 3.2 together with Theorem 4.1. Moreover item (1) in the statement of Theorem A is just item (1) of [46, Theorem 1] so it will not be proved here.

**4.1. Upper bound for large deviations.** Here we prove the upper bound in item 2 of Theorem 4.1.

Let  $\varphi : M \rightarrow \mathbb{R}$  be a fixed continuous function. Consider for  $n \geq 1$  and some fixed  $\varepsilon, \delta, c > 0$

$$A_n = A_n(\delta, \varepsilon) = \left\{ x : \frac{1}{n} S_n \Delta_\delta(x) \leq \varepsilon \right\} \text{ and } B_n = \left\{ x : \frac{1}{n} S_n \varphi(x) \geq c \right\}.$$

Since we want to bound a limit superior from above, we can assume without loss that  $\text{Leb}(A_n \cap B_n) > 0$  in what follows. We fix a partition  $\mathcal{P}$  of  $M$  as in Subsection 3.1 (whose diameter is smaller than  $\delta_1/4$ ) and use Lemma 3.5 with  $m = n$ ,  $E \subset U_1 \subset A_n \cap B_n$  such that  $U_1$  is open and

$$\text{Leb}((B_n \cap A_n) \setminus E) < \frac{1}{2n} \text{Leb}(B_n \cap A_n),$$

which can be done since  $S_n\varphi$  is continuous and  $S_n\Delta_\delta$  is upper-semicontinuous. Then we can find a family  $\mathcal{U}_n = \mathcal{E}_1 \cup \dots \cup \mathcal{E}_k$  of hyperbolic pre-balls contained in  $U_1$  satisfying

$$\text{Leb}(E \triangle \bigcup \mathcal{U}_n) \leq \left(1 - \frac{\theta}{4}\right)^k < \frac{1}{2n} \text{Leb}(A_n \cap B_n).$$

Note that  $\text{Leb}((A_n \cap B_n) \setminus \mathcal{U}_n) \leq \text{Leb}((A_n \cap B_n) \setminus E) + \text{Leb}(E \setminus \mathcal{U}_n) < \frac{1}{n} \text{Leb}(A_n \cap B_n)$  and so

$$\text{Leb}(A_n \cap B_n) < \frac{n}{n-1} \text{Leb}(\mathcal{U}_n). \quad (19)$$

Observe also that for any element  $P \in \mathcal{E}_i$  there exists  $x \in M$  and a hyperbolic time  $h_i$  of  $f$  for  $x$  such that  $P \subset B(x, h_i, \delta_1)$ , by construction, where  $i = 1, \dots, k_n$  and  $n < h_1 < \dots < h_{k_n}$ . Let  $\mathcal{C}_n$  be the set of all such pairs  $(x, h_i)$ , one for each element of  $\mathcal{U}_n$  and to simplify the notation we write  $h_n$  for  $h_{k_n}$ .

Following the arguments in the proof of [46, Thm.1(2)] we consider the measure

$$\sigma_n = \frac{1}{Z_n} \sum_{(x,l) \in \mathcal{C}_n} e^{-S_l J(x)} \cdot \delta_x \quad \text{where} \quad Z_n = \sum_{(x,l) \in \mathcal{C}_n} e^{-S_l J(x)}.$$

Note that by definition each element of the partition  $\bigvee_{i=0}^{h_n-1} f^{-i}\mathcal{P}$  contains at most the first coordinate of one element of  $\mathcal{C}_n$ . Thus using [45, Lemma 9.9] we have

$$H_{\sigma_n} \left( \bigvee_{i=0}^{h_n-1} f^{-i}\mathcal{P} \right) - \int S_{l(x)} J(x) d\sigma_n(x) = \log \sum_{(x,l) \in \mathcal{C}_n} e^{-S_l J(x)},$$

where we write  $l(x)$  for the unique integer  $l$  such that  $(x, l) \in \mathcal{C}_n$ . Since  $S_{l(x)-n} J(f^n(x)) > 0$  (see Subsection 3.3) and  $l(x) > n$  we get

$$H_{\sigma_n} \left( \bigvee_{i=0}^{h_n-1} f^{-i}\mathcal{P} \right) - \int S_n J d\sigma_n \geq \log \sum_{(x,l) \in \mathcal{C}_n} e^{-S_l J(x)}. \quad (20)$$

Setting  $\mu_n = \frac{1}{n} \sum_{i=0}^n f_*^i \sigma_n$  and  $\mu$  a weak\* accumulation point of  $\mu_n$ , we may modify the initial partition  $\mathcal{P}$  according to Lemma 3.4 and Remark 3.7 so that its diameter is smaller than  $\delta_1/2$  and  $\mu(\partial\mathcal{P}) = 0$  without loss, keeping  $\mathcal{C}_n$  unchanged. As in [45, pag. 220] from the above we can deduce that for every  $q \geq 1$

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log Z_n \leq \frac{1}{q} \limsup_{n \rightarrow +\infty} H_{\mu_n} \left( \bigvee_{i=0}^{q-1} f^{-i}\mathcal{P} \right) + \limsup_{n \rightarrow +\infty} \int -J d\mu_n \quad (21)$$

$$\leq h_\mu(f, \mathcal{P}) - \int J d\mu \leq h_\mu(f) - \int J d\mu \quad (22)$$

if  $f$  is a local diffeomorphism, ensuring that  $\mu$  is  $f$ -invariant and that  $J$  is a continuous function (in this case  $\mathcal{S} = \emptyset$  and  $\Delta_\delta$  plays no role, we may take  $\Delta_\delta \equiv 0$  and  $A_n = M$ ). Observe that since the points in  $\mathcal{C}_n$  are contained in  $B_n$  and  $\mu_n$  is a linear convex combination of measures of the form

$\frac{1}{n} \sum_{i=0}^{n-1} \delta_{f^i(x)}$ , we get for all  $n \geq 1$

$$\begin{aligned} \int \varphi d\mu_n &= \frac{1}{n} \sum_{j=0}^{n-1} \sigma_n(\varphi \circ f^j) = \frac{1}{Z_n} \sum_{(x,l) \in \mathcal{C}_n} e^{-S_l J(x)} \cdot \frac{1}{n} \sum_{j=0}^{n-1} \varphi(f^j(x)) \\ &\geq c \cdot \frac{1}{Z_n} \sum_{(x,l) \in \mathcal{C}_n} e^{-S_l J(x)} = c \end{aligned} \quad (23)$$

and hence  $\int \varphi d\mu \geq c$  also because  $\varphi$  is a continuous function.

Note that from (19) and by Subsection 3.3 we get for some constant  $C > 0$

$$\begin{aligned} \text{Leb}(B_n) &\leq \frac{n}{n-1} \text{Leb}(\mathcal{U}_n) \leq \frac{n}{n-1} \sum_{(x,l) \in \mathcal{C}_n} \text{Leb}\left(B(x, l, \delta_1)\right) \\ &\leq \frac{n}{n-1} \sum_{(x,l) \in \mathcal{C}_n} C e^{-S_l J(x)} = \frac{Cn}{n-1} Z_n. \end{aligned} \quad (24)$$

Therefore we have shown that there exists  $\mu \in \mathcal{M}_f$  such that  $\int \varphi d\mu \geq c$  and

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb}(B_n) \leq \limsup_{n \rightarrow +\infty} \frac{1}{n} \log Z_n \leq h_\mu(f) - \int J d\mu,$$

which completes the proof of item 2 in the statement of Theorem 4.1 and Theorem A.

**4.2. Upper bound for partially hyperbolic diffeomorphisms.** Here we show that a bound similar to the one in item 2 of Theorem A also holds in the case of a partially hyperbolic non-uniformly expanding attracting set.

Let  $f : M \rightarrow M$  be a diffeomorphism satisfying the conditions of Theorem D, let  $\varphi : M \rightarrow \mathbb{R}$  be a continuous function, fix a real number  $c$  and set  $J = \log |\det Df| \cdot F$ . Observe that since we have Lemma 3.11 we may argue exactly as in the previous subsection to arrive at an inequality just like (20).

Again as in the previous subsection we consider  $\mu_n = \frac{1}{n} \sum_{i=0}^{n-1} f_*^i \sigma_n$  and  $\mu$  a weak\* accumulation point of  $\mu_n$ . We also modify the partition  $\mathcal{P}$  in such a way that the boundaries of each atom have zero measure with respect to all measures  $\mu$  and  $\mu_n, n \geq 1$ .

The inequality (20) enables us to obtain inequalities (21) and (22) exactly as before. Together with the volume estimates obtained in Subsection 3.3.2 we can then arrive also at inequality (24) just by using a different distortion constant and replacing the Jacobian of  $f$  by the Jacobian of  $f$  along the  $F$  direction. Hence we obtain the upper bound given by item 2 of Theorem A also in the setting of partially hyperbolic non-uniformly expanding attracting sets. This will be very useful to deduce Theorem D in Subsection 5.1.

**4.3. Upper bound with singular/critical set.** To obtain an analogous result to (22) in the limit with a transformation  $f$  with non-flat singularities, thus proving item 3 from Theorem A and Theorem 4.1, we need some extra work. Note that the same arguments lead us to (21) as before and, since the points in  $\mathcal{C}_n$  are contained in  $A_n \cap B_n$ , by the same calculations (23) above we also get  $\int \Delta_\delta d\mu_n \leq \varepsilon$  for every  $n \geq 1$ .

**Lemma 4.2.** *The singular set  $\mathcal{S}$  has null  $\mu$ -measure.*

*Proof.* Arguing by contradiction, assume that  $\mu(\mathbb{S}) > 0$ . Then there exists  $a > 0$  such that  $\mu(B(\mathbb{S}, \eta)) \geq a$  for all  $\eta > 0$ . Let  $\eta > 0$  be chosen so that  $\mu(\partial B(\mathbb{S}, \eta)) = 0$  and  $\inf_{B(\mathbb{S}, \eta)} \Delta_\delta \geq 4\varepsilon/a$ .

On the one hand, since  $\mu$  is a weak\* limit point of  $\mu_n$ , there exists  $n_0$  such that for  $n > n_0$  we have  $\mu_n(B(\mathbb{S}, \eta)) \geq a/2$ . On the other hand, since  $\Delta_\delta \geq 0$  we get by the choice of  $\eta$

$$\frac{4\varepsilon}{a} \mu_n(B(\mathbb{S}, \eta)) \leq \mu_n(\Delta_\delta \cdot \chi_{B(\mathbb{S}, \eta)}) \leq \mu_n(\Delta_\delta) \leq \varepsilon,$$

where  $\chi_{B(\mathbb{S}, \eta)}$  is the characteristic function of  $B(\mathbb{S}, \eta)$ , from which we deduce that  $\mu_n(B(\mathbb{S}, \eta)) \leq a/4$ . This contradiction shows that  $\mu(\mathbb{S}) = 0$  and concludes the proof.  $\square$

**Lemma 4.3.** *The functions  $\Delta_\delta, J$  and  $\psi$  are  $\mu$ -integrable.*

*Proof.* Let us define the sequence of functions

$$\Delta_\delta^k = \xi_k \circ \Delta_\delta \text{ where } \xi_k(x) = \begin{cases} k & \text{if } |x| \geq k \\ x & \text{if } |x| < k \end{cases}, \quad k \geq 1.$$

For  $k > k_0$  with  $k_0 > |\log(\delta/2)|$  and fixing  $\eta > 0$ , since  $\Delta_\delta^k$  is continuous and  $\Delta_\delta \geq \Delta_\delta^k$  there is an integer  $n_0$  such that for all  $n > n_0$  we have

$$\mu(\Delta_\delta^k) \leq \mu_n(\Delta_\delta^k) + \eta \leq \mu_n(\Delta_\delta) + \eta \leq \varepsilon + \eta.$$

Since this holds for all  $k \geq k_0$  and  $\Delta_\delta(x) \rightarrow \infty$  when  $x \rightarrow \mathbb{S}$ , we have proved

$$\int_{M \setminus \mathbb{S}} \Delta_\delta d\mu < \infty.$$

Thus we get  $\Delta_\delta \in L^1(\mu)$  since  $\mu(\mathbb{S}) = 0$  by Lemma 4.2.

For the other functions, note that by conditions (S2) and (S3) on the singular set  $\mathbb{S}$  we see that there exists a constant  $\zeta > \beta$  such that on a small neighborhood  $V$  of  $\mathbb{S}$  we have

$$|\log \|Df(x)^{-1}\|| + |\log |\det Df(x)^{-1}|| \leq \zeta |\log d(x, \mathbb{S})| \quad (25)$$

and since  $f$  is a local diffeomorphism on  $M \setminus \mathbb{S}$ , the  $\mu$ -integrability of  $\Delta_\delta$  implies that of  $\psi$  and  $J$ . This concludes the proof of the lemma.  $\square$

**Lemma 4.4.** *The measure  $\mu$  is  $f$ -invariant.*

*Proof.* Since by Lemma 4.2  $\mu(\mathbb{S}) = 0$  we can find a sequence  $\eta_n \rightarrow 0$  of positive numbers such that  $\mu(\partial B(\mathbb{S}, \eta_n)) = 0$  for all  $n \geq 1$  and  $\mu(B(\mathbb{S}, \eta_n)) \rightarrow 0$  when  $n \rightarrow \infty$ .

Let us fix  $\eta > 0$  and a continuous function  $h : M \rightarrow \mathbb{R}$ . Take  $n_0$  such that

$$\mu(B(\mathbb{S}, \eta_n)) \cdot \sup |h| < \frac{\eta}{2}$$

for all  $n > n_0$  and fix  $n_1 > n_0$  such that

$$\frac{1}{2} \mu(B(\mathbb{S}, \eta_n)) \leq \mu_n(B(\mathbb{S}, \eta_n)) \leq 2\mu(B(\mathbb{S}, \eta_n))$$

for all  $n \geq n_1$ . Then if  $\tilde{f}$  is any continuous extension of  $f|_{M \setminus B(\mathbb{S}, \eta_n)}$  to  $M$  (which always exists by Tietze Extension Theorem, see e.g. [32]) we get

$$\int |h \circ f - h \circ \tilde{f}| d\mu_n \leq \sup |h| \cdot \mu_n(B(\mathbb{S}, \eta_n)) < \eta \quad (26)$$

for all  $n > n_1$ . Also note that (26) holds with  $\mu$  in the place of  $\mu_n$ . Since  $h \circ \tilde{f}$  is continuous there exists  $n_2 > n_1$  such that

$$\left| \int h \circ \tilde{f} d\mu_n - \int h \circ \tilde{f} d\mu \right| < \eta \quad \text{for every } n > n_2.$$

Hence for  $n > n_2$  we get

$$\begin{aligned} \left| \int h \circ \tilde{f} d\mu_n - \int h \circ \tilde{f} d\mu \right| &\leq |\mu(h \circ f) - \mu(h \circ \tilde{f})| + |\mu(h \circ \tilde{f}) - \mu_n(h \circ \tilde{f})| \\ &\quad + |\mu_n(h \circ \tilde{f}) - \mu_n(h \circ f)| \leq 3\eta. \end{aligned}$$

Since  $h$  was an arbitrary continuous function and  $\eta$  was any positive number, we have shown that  $f_*\mu_n \rightarrow f_*\mu$  in the weak\* topology when  $n \rightarrow \infty$ . This is exactly what is needed to show that  $\mu$  is  $f$ -invariant:

$$f_*\mu = \lim_n f_*\mu_n = \lim_n \left( \frac{1}{n} \sum_{j=0}^{n-1} f_*^j \sigma_n + \frac{f_*^n \sigma_n - \sigma_n}{n} \right) = \lim_n \mu_n = \mu,$$

concluding the proof.  $\square$

Now we consider  $\tilde{J}$  a continuous extension of  $J|_{\chi_{M \setminus B(\mathcal{S}, \rho)}}$  to  $M$  with the same range (this is Tietze's Extension Theorem) for  $0 < \rho < \delta$  and write

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mu_n(-J) &= \limsup_{n \rightarrow \infty} [\mu_n((-J + \tilde{J})\chi_{B(\mathcal{S}, \rho)}) + \mu_n(-\tilde{J})] \\ &\leq 2 \limsup_{n \rightarrow \infty} \mu_n(\zeta \Delta_\delta) + \mu(-\tilde{J}) \leq 2\zeta\varepsilon - \mu(\tilde{J}) \end{aligned}$$

since  $\tilde{J}$  is continuous and  $|-J + \tilde{J}|_{\chi_{B(\mathcal{S}, \rho)}} \leq 2|J|_{\chi_{B(\mathcal{S}, \delta)}} \leq 2\zeta\Delta_\delta$  by (25). Taking  $\rho \rightarrow 0$  we get  $\mu(\tilde{J}) \rightarrow \mu(J)$  because  $J \in L^1(\mu)$  and together with (21) we arrive at

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log Z_n \leq h_\mu(f, \mathcal{P}) - \int J d\mu + 2\zeta\varepsilon$$

for some  $\mu \in \mathcal{M}_f$  with  $\mu(\varphi) \geq c$  and  $\Delta_\delta \in L^1(\mu)$ , which is enough to prove item (3) of Theorem 4.1 and Theorem A.

## 5. STRICTLY NEGATIVE UPPER BOUND

Here we prove Theorem B and Theorem D. For a  $C^1$  endomorphism  $f$  it is known [40] that the following inequality (also known as *Ruelle's inequality*) holds for every  $f$ -invariant probability measure  $\mu$

$$h_\mu(f) \leq \int \Sigma^+ d\mu. \quad (27)$$

where  $\Sigma^+$  denotes the sum of the positive Lyapunov exponents at  $\mu$ -a.e. point. In Subsection 5.3 we present a proof of this inequality in the setting of maps which are local diffeomorphisms away from a non-flat singular set  $\mathcal{S}$  with zero Lebesgue measure, for invariant probability measures  $\mu$  such that  $\log d(x, \mathcal{S})$  is  $\mu$ -integrable.

We note that in [25] a similar result was proved under more general geometric assumptions but stricter analytic hypothesis, mostly due to the fact that in [25] the authors considered  $M$  to be a compact metric space admitting a finite dimensional manifold  $V$  as an open dense subset and  $\mathcal{S} = M \setminus V$ ,

which demands technical conditions on how the Riemannian metric on  $V$  and  $f$  behave (including the first and second derivatives on local charts) near  $\mathcal{S}$  for the proof to work. Our conditions are similar except that we only need the transformation  $f$  to be  $C^1$  but assume that  $\log d(x, \mathcal{S})$  is integrable, which is natural in our setting.

**5.1. The local diffeomorphism and partially hyperbolic case.** From Ruelle's Inequality (27) and from Subsection 3.3 it follows that we get a non-positive upper bound in item (2) of Theorem A since  $\int J d\mu$  equals the sum of the Lyapunov exponents of  $\mu$  [34]. Moreover let  $\mu \in \mathbb{E}$  be given. Then we have

$$\int J d\mu = h_\mu(f) \leq \int \Sigma^+ d\mu \leq \int J d\mu.$$

Hence if  $\mu \in \mathcal{M}_f$  is not in  $\mathbb{E}$  then the inequality (27) is strict.

To prove Theorem B we fix a continuous  $\varphi : M \rightarrow \mathbb{R}$  and replace  $B_n$  in Subsection 4.1 with

$$B_n = \left\{ x \in M : \inf \left\{ \left| \frac{1}{n} S_n \varphi(x) - \eta(\varphi) \right| : \eta \in \mathbb{E} \right\} > \omega \right\} \quad (28)$$

for some  $\omega > 0$ . Then  $B_n$  is an open subset of  $M$  and we can assume without loss that  $\text{Leb}(A_n \cap B_n) > 0$  in what follows, for otherwise the limit superior in (8) is smaller than any given real number and there is nothing to prove. Hence arguing as in Subsection 4.1 we obtain a measure  $\nu \in \mathcal{M}_f$  satisfying  $\inf \{ |\nu(\varphi) - \eta(\varphi)| : \eta \in \mathbb{E} \} > \omega$ , the bound of item (3) of Theorem A and  $\Delta_\delta \in L^1(\nu)$  with  $\nu(\Delta_\delta) \leq \varepsilon$ .

If  $f$  is a local diffeomorphism, i.e.  $\mathcal{S} = \emptyset$ , then we can use the bound given by item (2) of Theorem A and it is enough to show that  $h_\nu(f) - \nu(J)$  is strictly negative. But we cannot have  $h_\nu(f) - \nu(J) = 0$  since by construction  $\nu$  is not in  $\mathbb{E}$ , thus  $h_\nu(f) - \nu(J) < 0$ , completing the proof of Theorem B in the case of a local diffeomorphism.

For a partially hyperbolic non-uniformly expanding attracting set we obtain a negative upper bound following the same reasoning as above since we can use the same bound from item (2) of Theorem A, as shown in Subsection 4.2, and we can also apply Ruelle's Inequality. This completes the proof of Theorem D.

**5.2. The case with singular/critical set.** In the case  $\mathcal{S} \neq \emptyset$  we now show that the upper bound in item (3) of Theorem A must be strictly negative for some values of  $\eta, \varepsilon, \delta > 0$  and for some  $\nu \in \mathcal{M}_f$ . For that we argue by contradiction and take decreasing sequences  $\varepsilon_n, \delta_n \rightarrow 0$  such that the corresponding measures  $\nu_k$  obtained according to the proof of Theorem A with  $B_n$  as in (28) and

$$A_n^k = \{ x \in M : \frac{1}{n} S_n \Delta_{\delta_i} \leq \varepsilon_i, i = 1, \dots, k \}$$

in the place of  $A_n$ , for each  $k \geq 1$ , satisfy

- $\nu_k \in \mathcal{M}_f$ ,  $\Delta_{\delta_i} \in L^1(\nu_k)$  and  $\nu_k(\Delta_{\delta_i}) \leq \varepsilon_i$  for  $i = 1, \dots, k$ ;
- $\limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Leb}(A_n^k \cap B_n) \leq h_{\nu_k}(f, \mathcal{P}) - \int J d\nu_k + 2\zeta\varepsilon_k$ ;
- $h_{\nu_k}(f, \mathcal{P}) - \int J d\nu_k + 2\zeta\varepsilon_k \geq 0$ ; and
- $\inf \{ |\nu_k(\varphi) - \eta(\varphi)| : \eta \in \mathbb{E} \} > \omega$ ;

where  $\mathcal{P}$  is a partition obtained using Lemma 3.4 with the sequence  $\mu_k = \nu_k$  and  $\mu$  some weak\* accumulation point of the  $\nu_k$ .

Thus on the one hand we have for any fixed  $N \geq 1$

$$h_{\nu_k}(f, \mathcal{P}) = \inf_{j \geq 1} \frac{1}{j} H_{\nu_k} \left( \bigvee_{i=0}^{j-1} f^{-i} \mathcal{P} \right) \leq \frac{1}{N} H_{\nu_k} \left( \bigvee_{i=0}^{N-1} f^{-i} \mathcal{P} \right)$$

and since  $\mu(\partial \mathcal{P}) = 0$  we get

$$\limsup_{k \rightarrow \infty} h_{\nu_k}(f, \mathcal{P}) \leq \frac{1}{N} H_{\mu} \left( \bigvee_{i=0}^{N-1} f^{-i} \mathcal{P} \right).$$

But  $N \geq 1$  was arbitrarily fixed, so

$$\limsup_{k \rightarrow \infty} h_{\nu_k}(f, \mathcal{P}) \leq \inf_{N \geq 1} \frac{1}{N} H_{\mu} \left( \bigvee_{i=0}^{N-1} f^{-i} \mathcal{P} \right) = h_{\mu}(f, \mathcal{P}).$$

On the other hand, choosing  $J_i$  to be a continuous extension of  $J \chi_{B(\mathcal{S}, \delta_i)}$  to  $M$  with the same range,  $i \geq 1$ , we have

$$\begin{aligned} \limsup_{k \rightarrow \infty} \nu_k(-J) &= \limsup_{k \rightarrow \infty} [\nu_k((-J + J_i) \chi_{B(\mathcal{S}, \delta_i)}) + \nu_k(-J_i)] \\ &\leq 2 \limsup_{k \rightarrow \infty} \nu_k(\zeta \Delta_{\delta_i}) + \mu(-J_i) \leq 2\zeta \varepsilon_i - \mu(J_i) \end{aligned}$$

since  $J_i$  is continuous and  $|-J + J_i| \chi_{B(\mathcal{S}, \delta_i)} \leq 2|J| \chi_{B(\mathcal{S}, \delta_i)} \leq 2\zeta \Delta_{\delta_i}$  by definition of  $\Delta_{\delta_i}$  and by (25). Similar arguments to the ones proving Lemmas 4.2, 4.3 and 4.4 show that  $J, \psi, \Delta_{\delta}$  are  $\mu$ -integrable and that  $\mu$  is  $f$ -invariant. Because  $i \geq 1$  can be arbitrarily chosen above and both  $\varepsilon_i \rightarrow 0$  and  $\mu(J_i) \rightarrow \mu(J)$ , we conclude that  $\limsup_{k \rightarrow \infty} \nu_k(-J) \leq -\mu(J)$ . Hence we deduce

$$0 \leq \limsup_{k \rightarrow \infty} (h_{\nu_k}(f, \mathcal{P}) + \nu_k(-J) + 2\zeta \varepsilon_k) \leq h_{\mu}(f, \mathcal{P}) - \mu(J) \leq h_{\mu}(f) - \mu(J)$$

and also that  $\inf \{ |\mu(\varphi) - \eta(\varphi)| : \eta \in \mathbb{E} \} \geq \omega > 0$  by construction. By Ruelle's Inequality we also get  $h_{\mu}(f) - \mu(J) \leq 0$ , which yields a contradiction since this means  $\mu \in \mathbb{E}$ . This contradiction shows that for some  $k \geq 1$

$$h_{\nu_k}(f, \mathcal{P}) - \int J d\nu_k + 2\zeta \varepsilon_k < 0$$

which proves Theorem B, except for the Ruelle Inequality for maps with non-flat singularities, which is the content of the next subsection.

### 5.3. Ruelle's Inequality for maps with non-flat singularities.

**Theorem 5.1.** *Let  $f : M \setminus \mathcal{S} \rightarrow M$  be a  $C^1$  local diffeomorphism away from a non-flat singular set  $\mathcal{S}$  and  $\mu$  a  $f$ -invariant probability measure such that  $|\log d(x, \mathcal{S})|$  is  $\mu$ -integrable. Then*

$$h_{\mu}(f) \leq \int \Sigma^+ d\mu,$$

where  $\Sigma^+$  denotes the sum of the positive Lyapunov exponents at a regular point, counting multiplicities.

Observe that the  $\mu$ -integrability of  $|\log d(x, \mathcal{S})|$  implies the  $\mu$ -integrability of  $\log^+ \|Df\|$ , where  $\log^+ x = \max\{0, \log x\}$ , and thus the Lyapunov exponents of  $f$  are well defined  $\mu$ -almost everywhere by Oseledec's Theorem [34]. The proof we present here follows Mañé [29, Chap. IV] closely.

We start by taking the  $M$  as a compact submanifold of  $\mathbb{R}^N$  with the usual Euclidean norm and induced Riemannian structure, and considering  $W_0$  an open *normal tubular neighborhood* of  $M$  in  $\mathbb{R}^N$ , that is, there exists  $\Phi : W_0 \rightarrow W, (x, u) \mapsto x + u$  a  $(C^\infty)$  diffeomorphism from a neighborhood  $W_0$  of the zero section of the normal bundle  $TM^\perp$  of  $M$  to  $W$ . Let also  $\pi : W \rightarrow M$  be the associated projection:  $\pi(w)$  is the closest point to  $w$  in  $M$  for  $w \in W$ , so that the line through the pair of points  $w, \pi(w)$  is normal to  $M$  at  $\pi(w)$ , see e.g. [23] or [22]. Now we define for  $\rho \in (0, 1)$

$$F_0 : W_0 \setminus (T_{\mathcal{S}}M) \rightarrow W_0, \quad (x, u) \mapsto (f(x), \rho \cdot u)$$

and also

$$F : W \setminus \Phi(T_{\mathcal{S}}M) \rightarrow W, \quad w \mapsto (\Phi \circ F_0 \circ \Phi^{-1})(w).$$

Then clearly  $F$  is a local diffeomorphism outside  $\Phi(T_{\mathcal{S}}M)$ ,  $\overline{F(W)} \subset W$  and  $M = \bigcap_{n \geq 0} F^n(W)$ .

For each  $n \geq 1$  consider the partition of  $\mathbb{R}^N$  into dyadic cubes

$$\mathcal{P}_n = \left\{ \prod_{i=1}^N \left[ \frac{a_i}{2^n}, \frac{a_i+1}{2^n} \right) : a_i \in \mathbb{Z}, i = 1, \dots, N \right\}.$$

Up to a slight translation of the partitions  $\mathcal{P}_n$  we can assume that the probability measure  $\mu$  on  $M$  satisfies  $\mu(M \cap \partial \mathcal{P}) = 0$ , where  $\partial \mathcal{P} = \bigcup_{n \geq 1} \partial \mathcal{P}_n \cup \mathcal{S}$ . For  $x \in M \setminus \partial \mathcal{P}$  we define

$$v_n(x) = v_n^F(x) = \#\{P \in \mathcal{P}_n : F(\mathcal{P}_n(x)) \cap P \neq \emptyset\}$$

and

$$v(x) = v^F(x) = \limsup_{n \rightarrow \infty} v_n(x)$$

where  $\mathcal{P}_n(x)$  denotes the atom of the partition  $\mathcal{P}_n$  containing  $x$ .

**Lemma 5.2.** *Let  $Q = [-1, 1]^N$  and  $x \in M \setminus \partial \mathcal{P}$ . Then*

$$v(x) \leq \sup_{z \in \mathbb{R}^n} \#\{P \in \mathcal{P}_1 : (z + Dg(x)Q) \cap P \neq \emptyset\}$$

*Proof.* For  $x \in M \setminus \partial \mathcal{P}$  and  $n \geq 1$  define  $\varphi_n(y) = x + y/n$  on  $\mathbb{R}^N$  and  $W_n = \varphi_n^{-1}(W)$ . Let  $F_n : W_n \rightarrow F_n(W_n) \subset W_n$  be such that

$$\begin{array}{ccc} W_n & \xrightarrow{F_n} & W_n \\ \varphi_n \downarrow & & \downarrow \varphi_n \\ W & \xrightarrow{F} & W \end{array}$$

commutes. We have  $F(w) = F(x) + DF(x)(w - x) + p_x(w)$  where  $p_x : W \setminus \Phi(T_{\mathcal{S}}M) \rightarrow \mathbb{R}^N$  is  $C^1$  and  $\lim_{w \rightarrow x} \|p_x(w)\|/\|w - x\| = 0$ , where  $\|\cdot\|$  is the Euclidean norm on  $\mathbb{R}^N$ . Then we write  $F_n(y) = DF(x)(y) + q_n^x(y) + \alpha_n(x)$  where

$$\alpha_n(x) = n \cdot F(x) - x \quad \text{and} \quad q_n^x(y) = n \cdot p_x(y/n + x). \quad (29)$$

Note that for  $x \in M \setminus \partial \mathcal{P}$  we have  $q_n^x \rightarrow 0$  uniformly on compacta. Indeed if  $\|y\| < r$  for some  $r > 0$  there is, for each given  $\delta > 0$ , a  $n_0 \in \mathbb{N}$  such

that  $\|y/n\| < \delta, \forall n \geq n_0$  and then, by definition of  $p_x$ , for all  $\varepsilon > 0$  there is  $n_1 \in \mathbb{N}$  so that  $\forall n \geq n_1, \|p_x(y/n + x)\| < \varepsilon\|y/n\|$  which is the same as  $\|n \cdot p_x(y/n + x)\| < \varepsilon r$ , or  $\|q_n^x(y)\| < \varepsilon r$  for all sufficient large  $n$ .

Commutativity of the diagram implies

$$F(\mathcal{P}_n(x)) \cap P \neq \emptyset \Leftrightarrow F_n(\varphi_n^{-1}(\mathcal{P}_n(x))) \cap \varphi_n^{-1}(P) \neq \emptyset.$$

But  $\varphi_n^{-1}(P)$  is an element of  $\mathcal{P}_1$  translated by some vector  $y_0 \in \mathbb{R}^N$ . Moreover  $\varphi_n^{-1}(\mathcal{P}_n(x)) \subset Q$  and so  $v_n(x) \leq \#\{P \in \mathcal{P}_1 : F_n(Q) \cap (P + y_0) \neq \emptyset\}$ . Because  $\alpha_n$  depends on  $x$  only

$$\begin{aligned} v_n(x) &\leq \#\left\{P \in \mathcal{P}_1 : \left(n \cdot DF(x)\left(\frac{1}{n}Q\right) + q_n^x(Q) + \alpha_n(x) - y_0\right) \cap P \neq \emptyset\right\} \\ &\leq \sup_{z \in \mathbb{R}^N} \#\{P \in \mathcal{P}_1 : (DF(x)Q + q_n^x(Q) + z) \cap P \neq \emptyset\} \end{aligned} \quad (30)$$

Since  $q_n^x \rightarrow 0$  on compact subsets we get

$$\limsup_{n \rightarrow \infty} v_n(x) \leq \sup_{z \in \mathbb{R}^N} \#\{P \in \mathcal{P}_1 : (DF(x)Q + z) \cap P \neq \emptyset\}$$

concluding the proof of the lemma.  $\square$

For the arguments which use the convergence properties of the sequence  $\log v_n$  we need the following result.

**Lemma 5.3.** *There exists a  $\mu$ -integrable function  $g$  such that  $0 \leq \log v_n \leq g$  for  $\mu$ -almost every point in  $M$  and for all  $n \geq 1$ .*

*Proof.* Fix  $n \geq 1$  and consider  $x \in M \setminus \partial\mathcal{P}$ . On the one hand since  $\mathcal{P}_n$  is a partition we must have  $v_n(x) \geq 1$ . On the other hand, by the bound (30) since the size of the edge of the cubes of  $\mathcal{P}_1$  is  $1/2$  in  $\mathbb{R}^N$  we get

$$v_n(x) \leq \left(2(\text{diam } DF(x)(Q) + \text{diam } q_n^x(Q))\right)^N \quad (31)$$

$$\begin{aligned} \text{diam } DF(x)(Q) &\leq 2\sqrt{N} \cdot \|DF(x)\| \\ &\leq 2\sqrt{N} \max\{\|Df(x)\|, \|DF \mid (T_x M)^\perp\|\}. \end{aligned} \quad (32)$$

Note that for  $x$  far away from  $\mathcal{S}$  we always get bounded expressions above since  $F$  is a local diffeomorphism outside of  $\Phi(T_{\mathcal{S}}M)$ . To bound  $\text{diam } q_n^x(Q)$  we use (29) and consider two cases.

First assume that  $d(x, \mathcal{S}) \geq 2/n$  and take  $y \in Q$ . Then for some  $\theta \in [0, 1]$

$$\begin{aligned} q_n^x(y) &= n \cdot p_x(y/n + x) = n \cdot \left(F(x + y/n) - F(x) - DF(x)(y/n)\right) \\ &= DF(x + \theta \cdot y/n)(y) - DF(x)(y) \end{aligned}$$

so we get by condition (S1) on  $\mathcal{S}$

$$\begin{aligned} \|q_n^x(y)\| &\leq \sqrt{N} \cdot (\|DF(x)\| + \|DF(x + \theta \cdot y/n)\|) \\ &\leq B\sqrt{N} \left(d(x, \mathcal{S})^{-\beta} + (d(x, \mathcal{S}) - 1/n)^{-\beta}\right) \\ &\leq B\sqrt{N} \cdot d(x, \mathcal{S})^{-\beta} \cdot (1 + 2^\beta) \end{aligned} \quad (33)$$

since  $1 - 1/(nd(x, \mathcal{S})) \geq 1/2$  and  $\|DF \mid (T_x M)^\perp\| \leq \rho < 1 \ll d(x, \mathcal{S})^{-\beta}$  for  $x$  close to  $\mathcal{S}$ , because  $\beta > 0$ .

Now assume that  $d(x, \mathcal{S}) < 2/n$ . Then we bound as follows

$$\begin{aligned} \|q_n^x(y)\| &\leq n \cdot \|F(x + y/n) - F(x)\| + \|DF(x)\| \cdot \|y\| \\ &\leq n \cdot \text{diam } W + B\sqrt{N} \cdot d(x, \mathcal{S})^{-\beta} \end{aligned} \quad (34)$$

Hence putting (31), (32), (33) and (34) together we see that there exists a constant  $\tilde{C} > 0$  such that

$$\log v_n(x) \leq \begin{cases} N \log (\tilde{C} d(x, \mathcal{S})^{-\beta}) & \text{if } d(x, \mathcal{S}) \geq 2/n, \\ N \log (\tilde{C} d(x, \mathcal{S})^{-\beta} + 2n \cdot \text{diam } W) & \text{if } d(x, \mathcal{S}) < 2/n. \end{cases}$$

But  $d(x, \mathcal{S})^{-\beta} > 0$  and we may assume without loss that  $2n \cdot \text{diam } W \geq 2$ , so

$$\log (\tilde{C} d(x, \mathcal{S})^{-\beta} + 2n \cdot \text{diam } W) \leq \log (\tilde{C} d(x, \mathcal{S})^{-\beta}) + \log (2n \cdot \text{diam } W)$$

and if  $d(x, \mathcal{S}) < 2/n$  we also get

$$\begin{aligned} \log d(x, \mathcal{S})^{-\beta} &= -\beta \log d(x, \mathcal{S}) \geq -\beta \log (2/n) = \beta \log (n/2) \\ &= \beta \log (2n \cdot \text{diam } W) - \beta \log (4 \text{diam } W) \quad \text{or} \\ \log (2n \cdot \text{diam } W) &\leq \log (4 \text{diam } W) - \log d(x, \mathcal{S}) \end{aligned}$$

Hence in all cases we arrive at

$$\log v_n(x) \leq N \log (C d(x, \mathcal{S})^{-\beta} + D)$$

for some positive constants  $C$  and  $D$ . This concludes the proof.  $\square$

**Lemma 5.4.** *The following bound on the entropy holds*

$$h_\mu(f, \mathcal{P}_n \cap M) = h_\mu(F \mid M, \mathcal{P}_n \cap M) \leq \int_M \log v_n^F d\mu.$$

*Proof.* This is [29, Lemma 12.2] without change.  $\square$

**Corollary 5.5.**  $h_\mu(f) = h_\mu(F \mid M) \leq \int_M \log v^F d\mu.$

*Proof.* Since  $\bigvee_{n \geq 1} (\mathcal{P}_n \cap M)$  is the Borel  $\sigma$ -algebra  $\mu \bmod 0$  we get

$$h_\mu(F \mid M) = \lim_{n \rightarrow \infty} h_\mu(F \mid M, \mathcal{P}_n \cap M) \leq \limsup_{n \rightarrow \infty} \int_M \log v_n^F d\mu.$$

By Lemma 5.3 we can use the Dominated Convergence Theorem to obtain

$$\limsup_{n \rightarrow \infty} \int_M \log v_n^F d\mu \leq \int_M \limsup_{n \rightarrow \infty} \log v_n^F d\mu = \int_M \log v^F d\mu$$

since  $\log$  is monotonous increasing. This concludes the proof.  $\square$

In what follows write  $v^n(x) = v^{F^n}(x)$  for the analogous to  $v^F(x)$  with  $F^n$  in the place of  $F$ .

**Lemma 5.6.** *We have*

$$h_\mu(f) = h_\mu(F \mid M) \leq \int \limsup_{n \rightarrow \infty} \frac{1}{n} \log v^n(x) d\mu(x).$$

*Proof.* Using [45, Thm. 4.13] and Corollary 5.5 we get for all  $n \geq 1$

$$h_\mu(F \mid M) = \frac{1}{n} h_\mu(F^n \mid M) \leq \int \frac{1}{n} \log v^n(x) d\mu(x). \quad (35)$$

Consider the sequence  $g_n(x) = n^{-1} \log v^n(x)$  and observe that by Lemma 5.2 and by (32)

$$\begin{aligned} g_n(x) &\leq \frac{1}{n} \log (2 \operatorname{diam}(DF^n(x)Q))^N \\ &\leq \frac{N}{n} \log(2\sqrt{N}) + \frac{N}{n} \log \|DF^n(x)\| = G_n(x). \end{aligned} \quad (36)$$

Again by (32) and by definition of  $F$  since  $x \in M$  we get  $\log \|DF(x)\| \leq \log^+ \|Df(x)\|$ . Hence by the  $f$ -invariance of  $\mu$  and the Sub-additive Ergodic Theorem [45, Thm. 10.1], the sequence  $G_n(x)$  tends to a finite limit  $G(x)$  for  $\mu$ -a.e.  $x$  when  $n \rightarrow \infty$ .

Now by (36) and by Fatou's Lemma [45, Thm. 0.9]

$$\int \liminf_{n \rightarrow \infty} (G_n - g_n) d\mu \leq \liminf_{n \rightarrow \infty} \int (G_n - g_n) d\mu. \quad (37)$$

On the one hand since  $\lim_{n \rightarrow \infty} G_n(x)$  exists  $\mu$ -a.e.

$$\int \liminf_{n \rightarrow \infty} (G_n - g_n) d\mu = \int (G - \limsup_{n \rightarrow \infty} g_n) d\mu \quad (38)$$

and, on the other hand, since  $\lim_{n \rightarrow \infty} \int G_n(x) d\mu$  exists  $\mu$ -a.e. we also get

$$\liminf_{n \rightarrow \infty} \int (G_n - g_n) d\mu = \int G d\mu - \limsup_{n \rightarrow \infty} \int g_n d\mu. \quad (39)$$

Altogether (37), (38) and (39) imply

$$\limsup_{n \rightarrow \infty} \int \frac{1}{n} \log v^n(x) d\mu(x) \leq \int \limsup_{n \rightarrow \infty} \frac{1}{n} \log v^n(x) d\mu(x)$$

which together with (35) conclude the proof of the Lemma.  $\square$

To finish we need to relate  $\limsup_{n \rightarrow \infty} \frac{1}{n} \log v^n(x)$  with the sum of the positive Lyapunov exponents at  $x$ . This is done just as in [29, Chap. IV, Sec. 12] where it is proved that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log v^n(x) \leq \Sigma^+(x)$$

for  $\mu$ -almost all  $x \in M$ . This together with Lemma 5.6 implies Ruelle's Inequality. The proof of Theorem 5.1 is complete.

## REFERENCES

- [1] J. Alves. *Statistical analysis of non-uniformly expanding dynamical systems*. Publicações Matemáticas do IMPA. [IMPA Mathematical Publications]. Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2003. 24<sup>o</sup> Colóquio Brasileiro de Matemática. [24th Brazilian Mathematics Colloquium].
- [2] J. F. Alves. SRB measures for non-hyperbolic systems with multidimensional expansion. *Ann. Sci. École Norm. Sup.*, 33:1–32, 2000.
- [3] J. F. Alves and V. Araujo. Random perturbations of nonuniformly expanding maps. *Astérisque*, 286:25–62, 2003.

- [4] J. F. Alves, V. Araújo, and B. Saussol. On the uniform hyperbolicity of nonuniformly hyperbolic systems. *Proceedings of the American Mathematical Society*, 131(4):1303–1309, 2003.
- [5] J. F. Alves and V. Araújo. Hyperbolic times: frequency versus integrability. *Ergodic Theory and Dynamical Systems*, 24:1–18, 2004.
- [6] J. F. Alves, C. Bonatti, and M. Viana. SRB measures for partially hyperbolic systems whose central direction is mostly expanding. *Invent. Math.*, 140(2):351–398, 2000.
- [7] J. F. Alves, S. Luzzatto, and V. Pinheiro. Markov structures and decay of correlations for non-uniformly expanding dynamical systems. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 22(6):817–839, 2005.
- [8] V. Araujo and M. J. Pacifico. Physical measures for infinite-modal maps. *Preprint IMPA Serie A*, 328/2004.
- [9] V. Araújo and A. Tahzibi. Stochastic stability at the boundary of expanding maps. *Nonlinearity*, 18:939–959, 2005.
- [10] M. Benedicks and L. Carleson. On iterations of  $1 - ax^2$  on  $(-1, 1)$ . *Annals of Math.*, 122:1–25, 1985.
- [11] M. Benedicks and L. Carleson. The dynamics of the Hénon map. *Annals of Math.*, 133:73–169, 1991.
- [12] M. Benedicks and L.-S. Young. Absolutely continuous invariant measures and random perturbations for certain one-dimensional maps. *Ergod. Th. & Dynam. Sys.*, 12:13–37, 1992.
- [13] M. Benedicks and L.-S. Young. SBR-measures for certain Hénon maps. *Invent. Math.*, 112:541–576, 1993.
- [14] M. Benedicks and L.-S. Young. Markov extensions and decay of correlations for certain Hénon maps. *Astérisque*, 261:13–56, 2000.
- [15] C. Bonatti, L. J. Díaz, and M. Viana. *Dynamics beyond uniform hyperbolicity*, volume 102 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2005. A global geometric and probabilistic perspective, Mathematical Physics, III.
- [16] C. Bonatti and M. Viana. SRB measures for partially hyperbolic systems whose central direction is mostly contracting. *Israel J. Math.*, 115:157–193, 2000.
- [17] R. Bowen and D. Ruelle. The ergodic theory of Axiom A flows. *Invent. Math.*, 29:181–202, 1975.
- [18] W. de Melo and S. van Strien. *One-dimensional dynamics*. Springer Verlag, 1993.
- [19] M. Denker. The central limit theorem for dynamical systems. In *Dynamical Systems and Ergodic Theory*, volume 23. Banach Center Publ., Warsaw, 1989.
- [20] M. Denker and M. Kesseböhmer. Thermodynamic formalism, large deviation, and multifractals. In *Stochastic climate models (Chorin, 1999)*, volume 49 of *Progr. Probab.*, pages 159–169. Birkhäuser, Basel, 2001.
- [21] S. Gouezel. Decay of correlations for nonuniformly expanding systems. *Preprint: <http://arxiv.org/math.DS/0401184>*, 2004.
- [22] V. Guillemin and A. Pollack. *Differential Topology*. Prentice Hall, New Jersey, 1974.
- [23] M. Hirsch. *Differential Topology*. Springer-Verlag, New-York, 1976.
- [24] M. Jakobson. Absolutely continuous invariant measures for one-parameter families of one-dimensional maps. *Comm. Math. Phys.*, 81:39–88, 1981.
- [25] A. Katok and J. M. Strelcyn. *Invariant manifolds, entropy and billiards. Smooth maps with singularities*, volume 1222 of *Lect. Notes in Math*. Springer Verlag, 1986.
- [26] A. Lasota and J. Yorke. On the existence of invariant measures for piecewise monotonic transformations. *Trans. Amer. Math. Soc.*, 186:481–488, 1973.
- [27] P.-D. Liu. Pesin’s Entropy Formula for endomorphisms. *Nagoya Math. J.*, 150:197–209, 1998.
- [28] S. Luzzatto and M. Viana. Positive Lyapunov exponents for Lorenz-like families with criticalities. *Astérisque*, 261:201–237, 2000. Géométrie complexe et systèmes dynamiques (Orsay, 1995).
- [29] R. Mañé. *Ergodic theory and differentiable dynamics*. Springer Verlag, New York, 1987.
- [30] R. J. Metzger. Sinai-Ruelle-Bowen measures for contracting Lorenz maps and flows. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 17(2):247–276, 2000.

- [31] L. Mora and M. Viana. Abundance of strange attractors. *Acta Math.*, 171(1):1–71, 1993.
- [32] J. Munkres. *Topology*. Prentice-Hall, New Deli, 2nd edition, 2004.
- [33] K. Oliveira and M. Viana. Existence and uniqueness of maximizing measures for robust classes of local diffeomorphisms. *Preprint IMPA serie A 383/2005*, 2005.
- [34] V. I. Oseledets. A multiplicative ergodic theorem: Lyapunov characteristic numbers for dynamical systems. *Trans. Moscow Math. Soc.*, 19:197–231, 1968.
- [35] Y. Pesin and Y. Sinai. Gibbs measures for partially hyperbolic attractors. *Ergod. Th. & Dynam. Sys.*, 2:417–438, 1982.
- [36] V. Pinheiro. SRB measures for weakly expanding maps. [arxiv.org/pdf/math.DS/0503433](http://arxiv.org/pdf/math.DS/0503433), 2005.
- [37] V. Pliss. On a conjecture due to Smale. *Diff. Uravneniya*, 8:262–268, 1972.
- [38] A. Rovella. The dynamics of perturbations of the contracting Lorenz attractor. *Bull. Braz. Math. Soc.*, 24(2):233–259, 1993.
- [39] D. Ruelle. A measure associated with Axiom A attractors. *Amer. J. Math.*, 98:619–654, 1976.
- [40] D. Ruelle. An inequality for the entropy of differentiable maps. *Bol. Soc. Bras. Mat.*, 9:83–87, 1978.
- [41] M. Rychlik. Bounded variation and invariant measures. *Studia Math.*, 76:69–80, 1983.
- [42] Y. Sinai. Gibbs measures in ergodic theory. *Russian Math. Surveys*, 27:21–69, 1972.
- [43] M. Viana. Multidimensional nonhyperbolic attractors. *Inst. Hautes Études Sci. Publ. Math.*, 85:63–96, 1997.
- [44] M. Viana. *Stochastic dynamics of deterministic systems*. Publicações Matemáticas do IMPA. [IMPA Mathematical Publications]. Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 1997. 21º Colóquio Brasileiro de Matemática. [21th Brazilian Mathematics Colloquium].
- [45] P. Walters. *An introduction to ergodic theory*. Springer Verlag, 1982.
- [46] L.-S. Young. Some large deviation results for dynamical systems. *Trans. Amer. Math. Soc.*, 318(2):525–543, 1990.
- [47] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Annals of Math.*, 147:585–650, 1998.

VÍTOR ARAÚJO, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL. *Presently at* INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, C. P. 68.530, 21.945-970 RIO DE JANEIRO, RJ-BRAZIL

*E-mail address:* [vdaraujo@fc.up.pt](mailto:vdaraujo@fc.up.pt)

MARIA JOSÉ PACIFICO, INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, C. P. 68.530, 21.945-970 RIO DE JANEIRO, BRAZIL

*E-mail address:* [pacifico@im.ufrj.br](mailto:pacifico@im.ufrj.br)