

# Eigenfunctions and fundamental solutions of the fractional two-parameter Laplacian

Semyon Yakubovich\*

November 2, 2009

Department of Pure Mathematics, Faculty of Science,  
University of Porto, Campo Alegre str., 687,  
4169-007 Porto, Portugal

## Abstract

We deal with the following fractional generalization of the Laplace equation for rectangular domains  $(x, y) \in (x_0, X_0) \times (y_0, Y_0) \subset \mathbb{R}_+ \times \mathbb{R}_+$ , which is associated with the Riemann-Liouville fractional derivatives

$$\Delta^{\alpha, \beta} u(x, y) = \lambda u(x, y), \quad \Delta^{\alpha, \beta} := D_{x_0+}^{1+\alpha} + D_{y_0+}^{1+\beta},$$

where  $\lambda \in \mathbb{C}$ ,  $(\alpha, \beta) \in [0, 1] \times [0, 1]$ . Reducing the left-hand side of this equation to the sum of fractional integrals by  $x$  and  $y$ , we then use the operational technique for the conventional right-sided Laplace transformation and its extension to generalized functions to describe a complete family of eigenfunctions and fundamental solutions of the operator  $\Delta^{\alpha, \beta}$  in classes of functions represented by the left-sided fractional integral of a summable function or just admitting a summable fractional derivative. A symbolic operational form of the solutions in terms of the Mittag-Leffler functions is exhibited. The case of the separation of variables is also considered. An analog of the fractional logarithmic solution is presented. Classical particular cases of solutions are demonstrated.

---

\*E-mail: syakubov@fc.up.pt. Work supported by *Fundação para a Ciência e a Tecnologia* (FCT, the programmes POCTI and POSI) through the *Centro de Matemática da Universidade do Porto* (CMUP). Available as a PDF file from <http://www.fc.up.pt/cmup>. The author thanks Vladimir Mityushev for the statement of the problem and fruitful discussions of possible applications during the 7th ISAAC Congress at Imperial College London in July of 2009.

**Key words:** Laplace equation, Riemann-Liouville fractional integrals and derivatives, generalized functions, Laplace transform, operational calculus, Mittag-Leffler function, logarithmic solution, strong and weak solutions, Hypergeometric functions

*2000 Mathematics Subject Classification:* 35J05, 26A33, 35A22, 44A10, 35C05, 33E12, 34B27, 33C20, 35D05

## 1 Introduction

Let  $D_{a+}^{\gamma}f$  and  $I_{a+}^{\gamma}f$  be the Riemann-Liouville fractional derivative and integral of order  $\gamma > 0$  defined by [3, 4]

$$(D_{a+}^{\gamma}f)(x) = \left(\frac{d}{dx}\right)^n \frac{1}{\Gamma(n-\gamma)} \int_a^x \frac{f(t)}{(x-t)^{\gamma-n+1}} dt, \quad a, x > 0, \quad n = [\gamma] + 1, \quad (1.1)$$

$$(I_{a+}^{\gamma}f)(x) = \frac{1}{\Gamma(\gamma)} \int_a^x \frac{f(t)}{(x-t)^{1-\gamma}} dt, \quad a, x > 0, \quad (1.2)$$

where  $[ \ ]$  means the integer part of  $\gamma$ . Consider a class of the linear non-homogeneous differential equations

$$(D_{x_0+}^{1+\alpha}u)(x, y) + (D_{y_0+}^{1+\beta}u)(x, y) - \lambda u(x, y) = f(x, y), \quad (1.3)$$

where  $\lambda \in \mathbb{C}$ ,  $(\alpha, \beta) \in [0, 1] \times [0, 1]$ ,  $f(x, y)$ ,  $0 \leq x_0 \leq x \leq X_0 < \infty$ ,  $0 \leq y_0 \leq y \leq Y_0 < \infty$  is a prescribed function and  $u(x, y)$  is to be determined. Denoting by

$$\Delta^{\alpha, \beta} := D_{x_0+}^{1+\alpha} + D_{y_0+}^{1+\beta}, \quad (1.4)$$

equation (1.3) can be written in the form  $(\Delta^{\alpha, \beta} - \lambda E)u = f$ , where  $E$  is the identity operator. When  $\alpha = \beta = 1$  we come out with the classical Poisson equation. Therefore we call fractional partial differential equation (1.3) *the fractional two-parameter Poisson equation (FPE)*. Its homogeneous analog it is naturally to call *the fractional Laplace equation (FLE)* or fractional two-parameter Laplacian.

In this paper we present a general operational approach [1] to describe eigenfunctions and fundamental solutions of the fractional two-parameter Laplacian basing on the conventional right-sided Laplace transform [5]

$$F(s) = \int_{T_f}^{\infty} f(t)e^{-st}dt, \quad T_f \geq 0, \quad \operatorname{Re} s > a_0 \quad (1.5)$$

of absolutely integrable functions  $f \in L_1((T_f, \infty); e^{-a_0 t} dt)$  with respect to the measure  $e^{-a_0 t} dt$  and its distributional analog

$$F(s) = \langle f(t), e^{-st} \rangle \quad (1.6)$$

in Zemanian's space  $\mathcal{L}'(a_0)$  defined below. Operational solutions will be written in terms of the generalized Mittag-Leffler function  $E_{\mu, \nu}(z)$ , [2], [3], [4] which is defined in terms of the power series

$$E_{\mu, \nu}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\mu n + \nu)}, \quad \mu > 0, \nu \in \mathbb{R}, z \in \mathbb{C}. \quad (1.7)$$

In particular, the function  $E_{\mu, \nu}(z)$  is entire of the order  $\rho = \frac{1}{\mu}$  and type  $\sigma = 1$ . The exponential function, trigonometric and hyperbolic functions are expressed through (1.7) as follows

$$E_{1,1}(z) = e^z, \quad E_{2,1}(-z^2) = \cos z, \quad E_{2,1}(z) = \cosh z, \quad (1.8)$$

$$zE_{2,2}(-z^2) = \sin z, \quad zE_{2,2}(z^2) = \sinh z. \quad (1.9)$$

We will consider in the sequel the existence and uniqueness of general solutions of the fractional Laplacian and its particular cases. Possible applications and an investigation of the fractional two-parameter Poisson equation (1.3) are still out of the framework of this paper and will be done in forthcoming articles of the author.

## 2 Eigenfunctions and fundamental solutions of the fractional Laplace equation

We begin with

**Definition 1** [3]. By  $AC^n([a, b])$ ,  $n \in \mathbb{N}$  we denote the class of functions  $f(x)$ , which are continuously differentiable on the segment  $[a, b]$  up to the order  $n - 1$  and  $f^{(n-1)}(x)$  is absolutely continuous on  $[a, b]$ .

It is known [3], that the class  $AC^n([a, b])$  contains only functions represented in the form

$$f(x) = \frac{1}{(n-1)!} \int_a^x (x-t)^{n-1} \varphi(t) dt + \sum_{k=0}^{n-1} c_k (x-a)^k, \quad (2.1)$$

where  $\varphi(t) \in L_1(a, b)$  and  $c_k$  are arbitrary constants. It is not difficult to find that  $\varphi(t) = f^{(n)}(t)$ ,  $c_k = f^{(k)}(a)/k!$ . Moreover, if  $f(x) \in AC^n([a, b])$  then fractional derivative (1.1) exists almost everywhere and can be represented by the formula

$$(D_{a+}^{\gamma} f)(x) = \sum_{k=0}^{n-1} \frac{f^{(k)}(a)}{\Gamma(1+k-\gamma)} (x-a)^{k-\gamma}$$

$$+\frac{1}{\Gamma(n-\gamma)} \int_a^x \frac{f^{(n)}(t)}{(x-t)^{\gamma-n+1}} dt, \quad n = [\gamma] + 1. \quad (2.2)$$

**Definition 2 [3].** By  $I_{a+}^\gamma(L_1)$  we denote the class of functions  $f$  represented by the left-sided fractional integral (1.2) of a summable function, i.e.  $f = I_{a+}^\gamma \varphi$ ,  $\varphi \in L_1(a, b)$ .

A description of this class is given by the following

**Theorem 1 [3].** A function  $f(x) \in I_{a+}^\gamma(L_1)$ ,  $\gamma > 0$  if and only if  $(I_{a+}^{n-\gamma} f)(x) \in AC^n([a, b])$ ,  $n = [\gamma] + 1$  and  $(I_{a+}^{n-\gamma} f)^{(k)}(a) = 0, k = 0, 1, \dots, n-1$ .

**Definition 3 [3].** We will say that a function  $f \in L_1(a, b)$  has a summable fractional derivative  $(D_{a+}^\gamma f)(x)$  if  $(I_{a+}^{n-\gamma} f)(x) \in AC^n([a, b])$ ,  $n = [\gamma] + 1$ .

If  $(D_{a+}^\gamma f)(x) = \left(\frac{d}{dx}\right)^n (I_{a+}^{n-\gamma} f)(x)$  exists in the ordinary sense, i.e.  $(I_{a+}^{n-\gamma} f)(x)$  is differentiable in each point up to the order  $n$ , then  $f(x)$  evidently admits the derivative  $(D_{a+}^\gamma f)(x)$  in the sense of Definition 3.

So, if  $f(x) \in I_{a+}^\gamma(L_1)$ , then  $(I_{a+}^\gamma D_{a+}^\gamma f)(x) = f(x)$ . Otherwise if  $f$  just admits a summable fractional derivative, then the composition of fractional operators (1.1), (1.2) can be written in the form (see [3])

$$\begin{aligned} (I_{a+}^\gamma D_{a+}^\gamma f)(x) &= f(x) - \sum_{k=0}^{n-1} \frac{(x-a)^{\gamma-k-1}}{\Gamma(\gamma-k)} \\ &\quad \times (I_{a+}^{n-\gamma} f)^{(n-k-1)}(a), \quad n = [\gamma] + 1. \end{aligned} \quad (2.3)$$

Nevertheless we note, that  $(D_{a+}^\gamma I_{a+}^\gamma f)(x) = f(x)$  for any summable function  $f$ .

Consider now the eigenfunction problem for the fractional Laplace equation in the rectangular domain  $(x, y) \in (x_0, X_0) \times (y_0, Y_0)$

$$(D_{x_0+}^{1+\alpha} u)(x, y) + (D_{y_0+}^{1+\beta} u)(x, y) = \lambda u(x, y), \quad (2.4)$$

where  $\lambda \in \mathbb{C}$ ,  $(\alpha, \beta) \in (0, 1] \times (0, 1]$  in the following three cases:

- i)  $u(x, y)$  belongs to classes  $I_{x_0+}^{1+\alpha}(L_1)$ ,  $I_{y_0+}^{1+\beta}(L_1)$  by  $x \in [x_0, X_0]$  and  $y \in [y_0, Y_0]$ , respectively;
- ii)  $u(x, y)$  admits a summable fractional derivative  $(D_{x_0+}^{1+\alpha} u)(x, y)$  by  $x \in [x_0, X_0]$  and belongs to  $I_{y_0+}^{1+\beta}(L_1)$  by  $y \in [y_0, Y_0]$  or vice versa;
- iii)  $u(x, y)$  admits summable fractional derivative  $(D_{x_0+}^{1+\alpha} u)(x, y)$ ,  $(D_{y_0+}^{1+\beta} u)(x, y)$  by  $x \in [x_0, X_0]$  and  $y \in [y_0, Y_0]$ , respectively.

**Theorem 2.** In the case i) trivial solution of the equation (2.4) is the only solution.

**Proof.** Indeed, taking the operator  $I_{x_0+}^{1+\alpha}$  from both sides of the equation (2.4) and using the identity  $(I_{x_0+}^{1+\alpha} D_{x_0+}^{1+\alpha} u)(x, y) = u(x, y)$  it becomes

$$u(x, y) + (I_{x_0+}^{1+\alpha} D_{y_0+}^{1+\beta} u)(x, y) - \lambda (I_{x_0+}^{1+\alpha} u)(x, y) = 0, \quad (2.5)$$

Hence, applying the operator  $I_{y_0+}^{1+\beta}$  to both sides of the equation (2.5), we use the fact that due Fubini's theorem this operator commutes with  $I_{x_0+}^{1+\alpha}$ . Then we obtain

$$(I_{x_0+}^{1+\alpha} u)(x, y) + (I_{y_0+}^{1+\beta} u)(x, y) - \lambda(I_{x_0+}^{1+\alpha} I_{y_0+}^{1+\beta} u)(x, y) = 0. \quad (2.6)$$

Hence from conditions of the theorem we observe that fractional integrals of the equation (2.6) are Laplace-transformable functions. Therefore we may act on (2.8) by the conventional right-sided Laplace transform (1.5), let say, by  $y$  with  $T_f = y_0$ . Taking into account its convolution and operational properties [1] after straightforward calculations we arrive at the following second kind homogeneous integral equation of the Volterra type

$$U(x, s) + \frac{(s^{1+\beta} - \lambda)}{\Gamma(1 + \alpha)} \int_{x_0}^x (x - t)^\alpha U(t, s) dt = 0, \quad (2.7)$$

where  $\lambda \in \mathbb{C}, s \in \mathbb{C}, \operatorname{Res} \geq a_0 > 0, s^{1+\beta} = |s|^{1+\beta} e^{i\theta}, \theta = \arg s \in (-\frac{\pi}{2}, \frac{\pi}{2})$  and

$$U(x, s) = \int_{y_0}^{Y_0} e^{-st} u(x, t) dt. \quad (2.8)$$

Appealing to [2, Chapter 3] we find that (2.7) has the only trivial solution in the space of summable functions and  $U(x, s) \in L_1(x_0, X_0)$  because  $U(x, s) \in I_{x_0+}^{1+\alpha}(L_1)$  for each  $s$ . Cancelling the Laplace transform and using its uniqueness property for summable functions we get  $u(x, y) = 0$ . Theorem 2 is proved.

In the case ii) equation (2.6) should be substituted by the following equalities (see (2.3))

$$\begin{aligned} & (I_{x_0+}^{1+\alpha} u)(x, y) + (I_{y_0+}^{1+\beta} u)(x, y) - \lambda(I_{x_0+}^{1+\alpha} I_{y_0+}^{1+\beta} u)(x, y) \\ &= \frac{(x - x_0)^{\alpha-1}}{\Gamma(\alpha)} (I_{y_0+}^{1+\beta} f_0)(y) + \frac{(x - x_0)^\alpha}{\Gamma(1 + \alpha)} (I_{y_0+}^{1+\beta} f_1)(y), \end{aligned} \quad (2.9)$$

or

$$\begin{aligned} & (I_{x_0+}^{1+\alpha} u)(x, y) + (I_{y_0+}^{1+\beta} u)(x, y) - \lambda(I_{x_0+}^{1+\alpha} I_{y_0+}^{1+\beta} u)(x, y) \\ &= + \frac{(y - y_0)^{\beta-1}}{\Gamma(\beta)} (I_{x_0+}^{1+\alpha} h_0)(x) + \frac{(y - y_0)^\beta}{\Gamma(1 + \beta)} (I_{x_0+}^{1+\alpha} h_1)(x), \end{aligned} \quad (2.10)$$

where we denoted by

$$f_0(y) = (I_{x_0+}^{1-\alpha} u)(x_0, y), \quad (2.11)$$

$$f_1(y) = (D_{x_0+}^\alpha u)(x_0, y), \quad (2.12)$$

$$h_0(x) = (I_{y_0+}^{1-\beta} u)(x, y_0), \quad (2.13)$$

$$h_1(x) = (D_{y_0+}^\beta u)(x, y_0) \quad (2.14)$$

Cauchy's fractional initial conditions. Treating, for instance, equation (2.9) we take the Laplace transform from its both sides and arrive at the following integral equation

$$U(x, s) + \frac{(s^{1+\beta} - \lambda)}{\Gamma(1 + \alpha)} \int_{x_0}^x (x - t)^\alpha U(t, s) dt = F(x, s), \quad x \in (x_0, X_0), \quad \lambda \in \mathbb{C}, \quad (2.15)$$

where

$$F(x, s) = \frac{(x - x_0)^{\alpha-1}}{\Gamma(\alpha)} F_0(s) + \frac{(x - x_0)^\alpha}{\Gamma(1 + \alpha)} F_1(s) \quad (2.16)$$

and

$$F_i(s) = \int_{y_0}^{Y_0} e^{-st} f_i(t) dt, \quad i = 0, 1. \quad (2.17)$$

It is known [2], that a unique solution of the equation (2.15) in the class of summable functions is

$$U(x, s) = F(x, s) - (s^{1+\beta} - \lambda) \int_{x_0}^x (x - t)^\alpha \times E_{1+\alpha, 1+\alpha} \left( -(s^{1+\beta} - \lambda)(x - t)^{1+\alpha} \right) F(t, s) dt, \quad (2.18)$$

which involves as the kernel the generalized Mittag-Leffler function (1.7). Next, substituting (2.16) into (2.15), using (1.7), (2.17), index laws for fractional operators [3] and the estimates

$$\begin{aligned} & \left| (s^{1+\beta} - \lambda) \int_{x_0}^x (x - t)^\alpha E_{1+\alpha, 1+\alpha} \left( -(s^{1+\beta} - \lambda)(x - t)^{1+\alpha} \right) F(t, s) dt \right| \\ & \leq |F_0(s)| \sum_{n=0}^{\infty} \frac{(|s|^{1+\beta} + |\lambda|)^{n+1}}{\Gamma((n+1)(1+\alpha))\Gamma(\alpha)} \int_{x_0}^x (x - t)^{n(1+\alpha)+\alpha} (t - x_0)^{\alpha-1} dt \\ & + |F_1(s)| \sum_{n=0}^{\infty} \frac{(|s|^{1+\beta} + |\lambda|)^{n+1}}{\Gamma((n+1)(1+\alpha))\Gamma(1+\alpha)} \int_{x_0}^x (x - t)^{n(1+\alpha)+\alpha} (t - x_0)^\alpha dt \\ & \leq (|s|^{1+\beta} + |\lambda|)(X_0 - x_0)^{2\alpha} \left[ E_{1+\alpha, 2\alpha+1} \left( (|s|^{1+\beta} + |\lambda|)(X_0 - x_0)^{1+\alpha} \right) \int_{y_0}^{Y_0} e^{-a_0 t} |f_0(t)| dt \right. \\ & \quad \left. + (X_0 - x_0) E_{1+\alpha, 2(\alpha+1)} \left( (|s|^{1+\beta} + |\lambda|)(X_0 - x_0)^{1+\alpha} \right) \left[ \int_{y_0}^{Y_0} e^{-a_0 t} |f_1(t)| dt \right] \right] \end{aligned}$$

we write solution (2.18) of the Volterra type equation (2.15) in terms of the Mittag-Leffler functions

$$\begin{aligned} U(x, s) &= (x - x_0)^{\alpha-1} E_{1+\alpha, \alpha} \left( -(s^{1+\beta} - \lambda)(x - x_0)^{1+\alpha} \right) F_0(s) \\ &+ (x - x_0)^\alpha E_{1+\alpha, 1+\alpha} \left( -(s^{1+\beta} - \lambda)(x - x_0)^{1+\alpha} \right) F_1(s). \end{aligned} \quad (2.19)$$

In order to cancel the Laplace transformation by  $s$  in (2.19) we will appeal to its distributional form (1.6) in Zemanian's space  $\mathcal{L}'(0)$  (see [5]), which is dual of the countable-union space of test functions  $\mathcal{L}(0)$  defined by

$$\mathcal{L}(0) = \cup_{\nu=1}^{\infty} \mathcal{L}_{a_\nu},$$

where  $\{a_\nu\}_{\nu=1}^{\infty}$  is a sequence of real numbers  $a_\nu > 0$ , which converges monotonically to  $0+$  as  $\nu \rightarrow \infty$  and each  $\mathcal{L}_{a_\nu}$  is a testing-function space of smooth functions  $\varphi(y)$ ,  $y \in (y_0, \infty)$  and for each nonnegative integer  $k$  it satisfies

$$\rho_{k,\nu}(\varphi) = \sup_{y \in (y_0, \infty)} e^{a_\nu y} |\varphi^{(k)}(y)| < \infty, \quad k \in \mathbb{N}_0. \quad (2.20)$$

According to [5, Chapter III] we assign  $\mathcal{L}_{a_\nu}$  a topology generated by the multinorm (2.20). Consequently,  $\mathcal{L}_{a_\nu}$  is a countably multinormed space and the kernel of the Laplace transform  $e^{-st}$  is a member of  $\mathcal{L}_{a_\nu}$  if and only if  $\text{Re } s \geq a_\nu$ . Taking the space  $\mathcal{L}(0)$  we have an advantage that the space of smooth functions with compact support  $\mathcal{D}$  is dense in  $\mathcal{L}(0)$  and the members of the dual  $\mathcal{L}'(0)$  are distributions. Moreover, any  $f \in \mathcal{L}'(0)$  is a right-sided Laplace-transformable generalized function via the formula (1.6) with the right half-plane  $\text{Re } s > 0$  as a region of definition. Meanwhile, any analytic function  $F(s)$  on the half-plane  $\text{Re } s \geq a > 0$ , which satisfies the estimate

$$|F(s)| \leq e^{-T \text{Re } s} P(|s|), \quad \text{Re } s \geq a \quad (2.21)$$

where  $P(z)$  is a polynomial, may be identified as the Laplace transform (1.6) of a right-sided Laplace -transformable generalized function which is concentrated on  $T \leq t < \infty$ . Finally, the uniqueness and inversion properties are true and the inversion formula has the form

$$f(t) = \lim_{r \rightarrow \infty} \int_{\sigma - ir}^{\sigma + ir} F(s) e^{st} ds, \quad (2.22)$$

in the sense of convergence in  $\mathcal{D}'$  for any  $\sigma > 0$ .

So in order to find eigenfunctions and general fundamental solutions of the fractional Laplace equation (2.4) we will invert the Laplace transform in (2.19) by using formula (2.22). Of course, we understand that the conventional right-sided Laplace transform (1.5) is a particular case of (1.6) being applied to a regular generalized function  $f \in L_1((T_f, \infty); e^{-a_0 t} dt)$ ,  $a_0 > 0$ .

Further, we have

$$\begin{aligned} \lim_{r \rightarrow \infty} \int_{\sigma - ir}^{\sigma + ir} s^{n(1+\beta)} F_i(s) e^{sy} ds &= \lim_{r \rightarrow \infty} \left( \frac{d}{dy} \right)^{[n(1+\beta)]+1} \int_{\sigma - ir}^{\sigma + ir} s^{\{n(1+\beta)\}-1} F_i(s) e^{sy} ds \\ &= (D_{y_0}^{n(1+\beta)} f_i)(y), \quad i = 0, 1, \quad n \in \mathbb{N}_0, \end{aligned}$$

where  $\{ \}$  is a fractional part of the number, the convergence is in  $\mathcal{D}'$  and we assume that  $(I_{y_0}^{1-\{n(1+\beta)\}} f_i)(y) \in AC^{[n(1+\beta)]+1}[y_0, Y_0]$  for any  $n \in \mathbb{N}_0$ . Therefore, canceling the Laplace transformation in (2.19) and taking into account (2.16) after straightforward calculations we get the expression for a family of eigenfunctions of the equation (2.4)

$$\begin{aligned} u_\lambda(x, y) = & \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (x - x_0)^{n(1+\alpha)+\alpha-1} \\ & \times {}_1\Psi_1 \left( (1, n+1); (1+\alpha, n(1+\alpha)+\alpha); \lambda(x-x_0)^{1+\alpha} \right) (D_{y_0}^{n(1+\beta)} f_0)(y) \\ & + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (x - x_0)^{n(1+\alpha)+\alpha} {}_1\Psi_1 \left( (1, n+1); (1+\alpha, (n+2)(1+\alpha)); \lambda(x-x_0)^{1+\alpha} \right) \\ & \times (D_{y_0}^{n(1+\beta)} f_1)(y), \end{aligned} \quad (2.23)$$

where  ${}_1\Psi_1((a_1, b_1); (c_1, d_1); z)$  is the generalized Wright function [4]

$${}_1\Psi_1((a_1, b_1); (c_1, d_1); z) = \sum_{m=0}^{\infty} \frac{\Gamma(a_1 m + b_1)}{\Gamma(b_1 m + d_1)} \frac{z^m}{m!}$$

and the convergence of series in (2.23) is in  $\mathcal{D}'$ . Letting in (2.23)  $\lambda = 0$  we immediately come out with a classical fundamental solution of the equation (2.4)

$$\begin{aligned} u_f(x, y) = & \sum_{n=0}^{\infty} (-1)^n \frac{(x - x_0)^{n(1+\alpha)+\alpha-1}}{\Gamma(n(1+\alpha)+\alpha)} (D_{y_0}^{n(1+\beta)} f_0)(y) \\ & + \sum_{n=0}^{\infty} (-1)^n \frac{(x - x_0)^{n(1+\alpha)+\alpha}}{\Gamma((n+1)(1+\alpha))} (D_{y_0}^{n(1+\beta)} f_1)(y). \end{aligned} \quad (2.24)$$

Taking into account definition (1.7) of the Mittag-Leffler function, solution (2.24) may be written in the operational form

$$\begin{aligned} u_f(x, y) = & (x - x_0)^{\alpha-1} E_{1+\alpha, \alpha} \left( -(x - x_0)^{1+\alpha} D_{y_0}^{1+\beta} \right) f_0(y) \\ & + (x - x_0)^{\alpha} E_{1+\alpha, 1+\alpha} \left( -(x - x_0)^{1+\alpha} D_{y_0}^{1+\beta} \right) f_1(y). \end{aligned} \quad (2.25)$$

Analogously, in the case of equation (2.10) we show that

$$\begin{aligned} u_\lambda(x, y) = & \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (y - y_0)^{n(1+\beta)+\beta-1} \\ & \times {}_1\Psi_1 \left( (1, n+1); (1+\beta, n(1+\beta)+\beta); \lambda(y-y_0)^{1+\beta} \right) (D_{x_0}^{n(1+\alpha)} h_0)(x) \end{aligned}$$



$$\begin{aligned}
 & + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (y - y_0)^{n(1+\beta)+\beta} {}_1\Psi_1 \left( (1, n+1); (1+\beta, (n+2)(1+\beta)); \lambda(y - y_0)^{1+\beta} \right) \\
 & \quad \times (D_{h_0}^{n(1+\alpha)} h_1)(x),
 \end{aligned} \tag{2.26}$$

and

$$\begin{aligned}
 u_h(x, y) &= (y - y_0)^{\beta-1} E_{1+\beta, \beta} \left( -(y - y_0)^{1+\beta} D_{x_0}^{1+\alpha} \right) h_0(x) \\
 & \quad + (y - y_0)^{\beta} E_{1+\beta, 1+\beta} \left( -(y - y_0)^{1+\beta} D_{x_0}^{1+\alpha} \right) h_1(x)
 \end{aligned} \tag{2.27}$$

are also correspondingly eigenfunctions and a fundamental solution of the equation (2.4).

On the other hand we may write solutions (2.25), (2.27) in the form of the generalized Neumann series. Namely, we find

$$\begin{aligned}
 u_f(x, y) &= \frac{1}{\Gamma(\alpha)} \sum_{n=0}^{\infty} (-1)^n I_{x_0}^{n(1+\alpha)} (x - x_0)^{\alpha-1} (D_{y_0}^{n(1+\beta)} f_0)(y) \\
 & \quad + \frac{1}{\Gamma(1+\alpha)} \sum_{n=0}^{\infty} (-1)^n I_{x_0}^{n(1+\alpha)} (x - x_0)^{\alpha} (D_{y_0}^{n(1+\beta)} f_1)(y),
 \end{aligned} \tag{2.28}$$

taking in mind the analyticity of series in (2.28) by  $x$  in the interval  $0 < \varepsilon \leq |x - x_0| < R$  and by  $y \in (y_0, Y_0)$ . In the same manner, we represent (2.27) by the expression

$$\begin{aligned}
 u_h(x, y) &= \frac{1}{\Gamma(\beta)} \sum_{n=0}^{\infty} (-1)^n I_{y_0}^{n(1+\beta)} (y - y_0)^{\beta-1} (D_{x_0}^{n(1+\alpha)} h_0)(x) \\
 & \quad + \frac{1}{\Gamma(1+\beta)} \sum_{n=0}^{\infty} (-1)^n I_{y_0}^{n(1+\beta)} (y - y_0)^{\beta} (D_{x_0}^{n(1+\alpha)} h_1)(x)
 \end{aligned} \tag{2.29}$$

with arbitrary  $h_0, h_1$  assuming analyticity of the corresponding series by  $y$  in the interval  $0 < \varepsilon \leq |y - y_0| < R$  and by  $x \in (x_0, X_0)$ . Now taking into account zero values  $D_{x_0+}^{1+\alpha} (x - x_0)^{\alpha-1} = 0$ ,  $D_{x_0+}^{1+\alpha} (x - x_0)^{\alpha} = 0$ ,  $D_{y_0+}^{1+\beta} (y - y_0)^{\beta-1} = 0$ ,  $D_{y_0+}^{1+\beta} (y - y_0)^{\beta} = 0$  it is not difficult to verify that (2.28), (2.29) are classical fundamental solutions of the fractional Laplacian  $\Delta^{\alpha, \beta} u = 0$  subject to conditions (2.11), (2.12) and (2.13), (2.14), respectively. Thus we have proved

**Theorem 3.** *In the case ii) functions (2.23), (2.26) represent eigenfunctions of the fractional Laplacian (2.4) and expressions (2.24), (2.27) are unique classical fundamental solutions subject to conditions (2.11), (2.12) and (2.13), (2.14), respectively. These solutions can be written in the corresponding form of generalized Neumann series (2.28), (2.29) under additional conditions of analyticity.*

Finally, in the case iii) an analog of equations (2.9), (2.10) is

$$(I_{x_0+}^{1+\alpha} u)(x, y) + (I_{y_0+}^{1+\beta} u)(x, y) - \lambda (I_{x_0+}^{1+\alpha} I_{y_0+}^{1+\beta} u)(x, y)$$

$$\begin{aligned}
&= \frac{(x - x_0)^{\alpha-1}}{\Gamma(\alpha)} (I_{y_0+}^{1+\beta} f_0)(y) + \frac{(x - x_0)^\alpha}{\Gamma(1 + \alpha)} (I_{y_0+}^{1+\beta} f_1)(y) \\
&+ \frac{(y - y_0)^{\beta-1}}{\Gamma(\beta)} (I_{x_0+}^{1+\alpha} h_0)(x) + \frac{(y - y_0)^\beta}{\Gamma(1 + \beta)} (I_{x_0+}^{1+\alpha} h_1)(x).
\end{aligned} \tag{2.30}$$

Consequently, in the right-hand side of Volterra's equation (2.15) we get an additional term

$$\begin{aligned}
U(x, s) + \frac{(s^{1+\beta} - \lambda)}{\Gamma(1 + \alpha)} \int_{x_0}^x (x - t)^\alpha U(t, s) dt &= F(x, s) \\
&+ e^{-y_0 s} (I_{x_0+}^{1+\alpha} (s h_0 + h_1))(x),
\end{aligned} \tag{2.31}$$

which will give a source for generalized eigenfunctions and fundamental solutions of the fractional Laplacian (2.4). In fact, owing to the estimates

$$\begin{aligned}
&\left| (s^{1+\beta} - \lambda) \int_{x_0}^x (x - t)^\alpha E_{1+\alpha, 1+\alpha} \left( -(s^{1+\beta} - \lambda)(x - t)^{1+\alpha} \right) F(t, s) dt \right| \\
&\leq |F_0(s)| \sum_{n=0}^{\infty} \frac{(|s|^{1+\beta} + |\lambda|)^{n+1}}{\Gamma((n+1)(1+\alpha))\Gamma(\alpha)} \int_{x_0}^x (x - t)^{n(1+\alpha)+\alpha} (t - x_0)^{\alpha-1} dt \\
&+ |F_1(s)| \sum_{n=0}^{\infty} \frac{(|s|^{1+\beta} + |\lambda|)^{n+1}}{\Gamma((n+1)(1+\alpha))\Gamma(1+\alpha)} \int_{x_0}^x (x - t)^{n(1+\alpha)+\alpha} (t - x_0)^\alpha dt \\
&+ \sum_{n=0}^{\infty} \frac{|s|(|s|^{1+\beta} + |\lambda|)^{n+1}}{\Gamma((n+1)(1+\alpha))\Gamma(1+\alpha)} \int_{x_0}^x (x - t)^{n(1+\alpha)+\alpha} \int_{x_0}^t (t - v)^\alpha |h_0(v)| dv dt \\
&+ \sum_{n=0}^{\infty} \frac{(|s|^{1+\beta} + |\lambda|)^{n+1}}{\Gamma((n+1)(1+\alpha))\Gamma(1+\alpha)} \int_{x_0}^x (x - t)^{n(1+\alpha)+\alpha} \int_{x_0}^t (t - v)^\alpha |h_1(v)| dv dt \\
&\leq (|s|^{1+\beta} + |\lambda|)(X_0 - x_0)^{2\alpha} \left[ E_{1+\alpha, 2\alpha+1} \left( (|s|^{1+\beta} + |\lambda|)(X_0 - x_0)^{1+\alpha} \right) \int_{y_0}^{Y_0} e^{-a_0 t} |f_0(t)| dt \right. \\
&+ (X_0 - x_0) E_{1+\alpha, 2(\alpha+1)} \left( (|s|^{1+\beta} + |\lambda|)(X_0 - x_0)^{1+\alpha} \right) \left[ \int_{y_0}^{Y_0} e^{-a_0 t} |f_1(t)| dt \right. \\
&\quad \left. \left. + |s| \int_{x_0}^{X_0} |h_0(t)| dt + \int_{x_0}^{X_0} |h_1(t)| dt \right] \right] < \infty,
\end{aligned}$$

we write solution of the Volterra type equation (2.31) in terms of the Mittag-Leffler functions and generalized Neumann series

$$U(x, s) = (x - x_0)^{\alpha-1} E_{1+\alpha, \alpha} \left( -(s^{1+\beta} - \lambda)(x - x_0)^{1+\alpha} \right) F_0(s)$$

$$\begin{aligned}
 & + (x - x_0)^\alpha E_{1+\alpha, 1+\alpha} \left( -(s^{1+\beta} - \lambda)(x - x_0)^{1+\alpha} \right) F_1(s) \\
 & + e^{-y_0 s} \sum_{n=0}^{\infty} (-1)^n (s^{1+\beta} - \lambda)^n \left( I_{x_0}^{(n+1)(1+\alpha)} (s h_0 + h_1) \right) (x).
 \end{aligned}$$

Cancelling the Laplace transformation we take in mind the relations

$$\begin{aligned}
 & \lim_{r \rightarrow \infty} \int_{\sigma - ir}^{\sigma + ir} s^{n(1+\beta)} e^{s(y-y_0)} ds \\
 & = \lim_{r \rightarrow \infty} \left( \frac{d}{dy} \right)^{[n(1+\beta)]+1} \int_{\sigma - ir}^{\sigma + ir} \left\langle \frac{t_+^{-\{n(1+\beta)\}}}{\Gamma(1 - \{n(1+\beta)\})} * \delta(t - y_0), e^{-st} \right\rangle e^{sy} ds \\
 & = \left( \frac{d}{dy} \right)^{[n(1+\beta)]+1} \left( \frac{t_+^{-\{n(1+\beta)\}}}{\Gamma(1 - \{n(1+\beta)\})} * \delta(t - y_0) \right) (y) = (D_{y_0}^{n(1+\beta)} \delta)(y),
 \end{aligned}$$

where  $\delta$  is Dirac's delta-function and  $*$  denotes the convolution product. Therefore, after straightforward calculations we get the expression for a family of eigenfunctions of the equation (2.4)

$$\begin{aligned}
 u_\lambda(x, y) & = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (x - x_0)^{n(1+\alpha)+\alpha-1} \\
 & \times {}_1\Psi_1 \left( (1, n+1); (1+\alpha, n(1+\alpha)+\alpha); \lambda(x - x_0)^{1+\alpha} \right) (D_{y_0}^{n(1+\beta)} f_0)(y) \\
 & + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (x - x_0)^{n(1+\alpha)+\alpha} {}_1\Psi_1 \left( (1, n+1); (1+\alpha, (n+2)(1+\alpha)); \lambda(x - x_0)^{1+\alpha} \right) \\
 & \times (D_{y_0}^{n(1+\beta)} f_1)(y) + \int_{x_0}^x h_1(t) \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \\
 & \times (x - t)^{n(1+\alpha)+\alpha} {}_1\Psi_1 \left( (1, n+1); (1+\alpha, (n+2)(1+\alpha)); \lambda(x - t)^{1+\alpha} \right) \\
 & \times (D_{y_0}^{n(1+\beta)} \delta)(y) dt + \int_{x_0}^x h_0(t) \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (x - t)^{n(1+\alpha)+\alpha} \\
 & \times {}_1\Psi_1 \left( (1, n+1); (1+\alpha, (n+2)(1+\alpha)); \lambda(x - t)^{1+\alpha} \right) \\
 & \times (D_{y_0}^{n(1+\beta)} \delta')(y) dt, \tag{2.32}
 \end{aligned}$$

where the convergence of series in (2.32) is in  $\mathcal{D}'$ . Letting in (2.32)  $\lambda = 0$  we derive a generalized fundamental solution of the equation (2.4)

$$u(x, y) = \sum_{n=0}^{\infty} (-1)^n \frac{(x - x_0)^{n(1+\alpha)+\alpha-1}}{\Gamma(n(1+\alpha)+\alpha)} (D_{y_0}^{n(1+\beta)} f_0)(y)$$

$$\begin{aligned}
& + \sum_{n=0}^{\infty} (-1)^n \frac{(x-x_0)^{n(1+\alpha)+\alpha}}{\Gamma((n+1)(1+\alpha))} (D_{y_0}^{n(1+\beta)} f_1)(y) \\
& + \int_{x_0}^x h_1(t) \sum_{n=0}^{\infty} (-1)^n \frac{(x-t)^{(n+1)(1+\alpha)-1}}{\Gamma((n+1)(1+\alpha))} (D_{y_0}^{n(1+\beta)} \delta)(y) dt \\
& + \int_{x_0}^x h_0(t) \sum_{n=0}^{\infty} (-1)^n \frac{(x-t)^{(n+1)(1+\alpha)-1}}{\Gamma((n+1)(1+\alpha))} (D_{y_0}^{n(1+\beta)} \delta')(y) dt,
\end{aligned} \tag{2.33}$$

which may be written in the operational form

$$\begin{aligned}
u(x, y) &= (x-x_0)^{\alpha-1} E_{1+\alpha, \alpha} \left( -(x-x_0)^{1+\alpha} D_{y_0}^{1+\beta} \right) f_0(y) \\
&+ (x-x_0)^{\alpha} E_{1+\alpha, 1+\alpha} \left( -(x-x_0)^{1+\alpha} D_{y_0}^{1+\beta} \right) f_1(y) \\
&+ \int_{x_0}^x (x-t)^{\alpha} E_{1+\alpha, 1+\alpha} \left( -(x-t)^{1+\alpha} D_{y_0}^{1+\beta} \right) \delta(y) h_1(t) dt \\
&+ \int_{x_0}^x (x-t)^{\alpha} E_{1+\alpha, 1+\alpha} \left( -(x-t)^{1+\alpha} D_{y_0}^{1+\beta} \right) \delta'(y) h_0(t) dt.
\end{aligned} \tag{2.34}$$

Analogously, functions

$$\begin{aligned}
u_{\lambda}(x, y) &= \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (y-y_0)^{n(1+\beta)+\beta-1} \\
&\times {}_1\Psi_1 \left( (1, n+1); (1+\beta, n(1+\beta)+\beta); \lambda(y-y_0)^{1+\beta} \right) (D_{x_0}^{n(1+\alpha)} h_0)(x) \\
&+ \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (y-y_0)^{n(1+\beta)+\beta} {}_1\Psi_1 \left( (1, n+1); (1+\beta, (n+2)(1+\beta)); \lambda(y-y_0)^{1+\beta} \right) \\
&\times (D_{h_0}^{n(1+\alpha)} h_1)(x) + \int_{y_0}^y h_1(t) \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \\
&\times (y-t)^{n(1+\beta)+\beta} {}_1\Psi_1 \left( (1, n+1); (1+\beta, (n+2)(1+\beta)); \lambda(y-t)^{1+\beta} \right) \\
&\times (D_{x_0}^{n(1+\alpha)} \delta)(x) dt + \int_{y_0}^y h_0(t) \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (y-t)^{n(1+\beta)+\beta} \\
&\times {}_1\Psi_1 \left( (1, n+1); (1+\beta, (n+2)(1+\beta)); \lambda(y-t)^{1+\beta} \right) \\
&\times (D_{x_0}^{n(1+\alpha)} \delta')(x) dt,
\end{aligned} \tag{2.35}$$

and

$$u(x, y) = (y-y_0)^{\beta-1} E_{1+\beta, \beta} \left( -(y-y_0)^{1+\beta} D_{x_0}^{1+\alpha} \right) h_0(x)$$

$$\begin{aligned}
 & + (y - y_0)^\beta E_{1+\beta, 1+\beta} \left( -(y - y_0)^{1+\beta} D_{x_0}^{1+\alpha} \right) h_1(x) \\
 & + \int_{y_0}^y (y - t)^\beta E_{1+\beta, 1+\beta} \left( -(y - t)^{1+\beta} D_{x_0}^{1+\alpha} \right) \delta(x) f_1(t) dt \\
 & + \int_{y_0}^y (y - t)^\beta E_{1+\beta, 1+\beta} \left( -(y - t)^{1+\beta} D_{x_0}^{1+\alpha} \right) \delta'(x) f_0(t) dt
 \end{aligned} \tag{2.36}$$

are also correspondingly eigenfunctions and generalized fundamental solutions of the equation (2.4).

**Theorem 4.** *In the case iii) functions (2.32), (2.35) represent eigenfunctions and expressions (2.33), (2.34), (2.36) are generalized fundamental solutions of the fractional Laplacian (2.4).*

**Example 1.** As a particular case, it is not difficult to obtain from (2.24), (2.27) the classical fundamental solution  $u(x, y) = \frac{1}{2} \log((x - x_0)^2 + (y - y_0)^2)$  of the Laplace equation  $\Delta u = 0$  ( $\alpha = \beta = 1$ ). Indeed, putting  $f_1(y) = h_1(x) = 0$ ,  $f_0(y) = \log(y - y_0)$ ,  $h_0(x) = \log(x - x_0)$  we assume, correspondingly,  $\frac{x - x_0}{y - y_0} < 1$ ,  $\frac{x - x_0}{y - y_0} > 1$  in (2.24), (2.27) and, for instance, solution  $u_f(x, y)$  becomes

$$\begin{aligned}
 u_f(x, y) &= \sum_{n=0}^{\infty} \frac{(-1)^n (x - x_0)^{2n}}{(2n)!} (\log(y - y_0))^{(2n)} = \log(y - y_0) \\
 &+ \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n} \left( \frac{x - x_0}{y - y_0} \right)^{2n} \\
 &= \log(y - y_0) + \sum_{n=0}^{\infty} \frac{(-1)^n}{2(n+1)} \left( \frac{x - x_0}{y - y_0} \right)^{2(n+1)} = \log(y - y_0) \\
 &+ \frac{1}{2} \log \left( 1 + \left( \frac{x - x_0}{y - y_0} \right)^2 \right) = \frac{1}{2} \log((x - x_0)^2 + (y - y_0)^2).
 \end{aligned}$$

Analogously we treat solution (2.27).

### 3 Separation of variables. Analytic solutions

The method of separation of variables allows us to simplify eigenfunctions and fundamental solutions of the fractional Laplacian. Indeed, putting  $u(x, y) = u_1(x)u_2(y)$ , substituting in (2.30) and taking into account initial conditions (2.11), (2.12), (2.13), (2.14) it becomes

$$u_2(y)(I_{x_0+}^{1+\alpha} u_1)(x) + u_1(x)(I_{y_0+}^{1+\beta} u_2)(y) - \lambda(I_{x_0+}^{1+\alpha} u_1)(x)(I_{y_0+}^{1+\beta} u_2)(y)$$

$$\begin{aligned}
&= a_1 \frac{(x - x_0)^{\alpha-1}}{\Gamma(\alpha)} (I_{y_0+}^{1+\beta} u_2)(y) + a_2 \frac{(x - x_0)^\alpha}{\Gamma(1 + \alpha)} (I_{y_0+}^{1+\beta} u_2)(y) \\
&+ b_1 \frac{(y - y_0)^{\beta-1}}{\Gamma(\beta)} (I_{x_0+}^{1+\alpha} u_1)(x) + b_2 \frac{(y - y_0)^\beta}{\Gamma(1 + \beta)} (I_{x_0+}^{1+\alpha} u_1)(x),
\end{aligned} \tag{3.1}$$

where  $a_i, b_i \in \mathbb{C}$ ,  $i = 1, 2$  are arbitrary constants. If  $(I_{x_0+}^{1+\alpha} u_1)(x)(I_{y_0+}^{1+\beta} u_2)(y) \neq 0$ ,  $(x, y) \in (x_0, X_0) \times (y_0, Y_0)$  we divide (3.1) by this product and separate variables, getting two Abel's type second kind integral equations to define  $u_i$ ,  $i = 1, 2$ , namely

$$u_1(x) + \mu(I_{x_0+}^{1+\alpha} u_1)(x) = a_1 \frac{(x - x_0)^{\alpha-1}}{\Gamma(\alpha)} + a_2 \frac{(x - x_0)^\alpha}{\Gamma(1 + \alpha)}, \tag{3.2}$$

$$u_2(y) - (\lambda + \mu)(I_{y_0+}^{1+\beta} u_2)(y) = b_1 \frac{(y - y_0)^{\beta-1}}{\Gamma(\beta)} + b_2 \frac{(y - y_0)^\beta}{\Gamma(1 + \beta)}, \tag{3.3}$$

where  $\lambda, \mu \in \mathbb{C}$  are constants. We note that the equality  $(I_{x_0+}^{1+\alpha} u_1)(x)(I_{y_0+}^{1+\beta} u_2)(y) = 0$  for at least one point  $(\xi, \eta)$  agrees with (3.1) and (3.2), (3.3). So we solve the latter equations similarly to (2.18), arriving at the following family of eigenfunctions  $u_{\lambda, \mu}(x, y) = u_1(x)u_2(y)$ , where

$$\begin{aligned}
u_1(x) &= a_1(x - x_0)^{\alpha-1} E_{1+\alpha, \alpha}(\mu(x - x_0)^{1+\alpha}) \\
&+ a_2(x - x_0)^\alpha E_{1+\alpha, 1+\alpha}(\mu(x - x_0)^{1+\alpha}),
\end{aligned} \tag{3.4}$$

$$\begin{aligned}
u_2(y) &= b_1(y - y_0)^{\beta-1} E_{1+\beta, \beta}((\lambda - \mu)(y - y_0)^{1+\beta}) \\
&+ b_2(y - y_0)^\beta E_{1+\beta, 1+\beta}((\lambda - \mu)(y - y_0)^{1+\beta}).
\end{aligned} \tag{3.5}$$

On the other hand, we may write these solutions in terms of the generalized Neumann series. Precisely, denoting by

$$v_\alpha(x) = a_1 \frac{(x - x_0)^{\alpha-1}}{\Gamma(\alpha)} + a_2 \frac{(x - x_0)^\alpha}{\Gamma(1 + \alpha)}, \tag{3.6}$$

$$v_\beta(y) = b_1 \frac{(y - y_0)^{\beta-1}}{\Gamma(\beta)} + b_2 \frac{(y - y_0)^\beta}{\Gamma(1 + \beta)} \tag{3.7}$$

and recalling index properties for the fractional integral (1.2) we get representations of (3.4), (3.5) in the respective resolvent form for fractional integral operators  $I_{x_0}^{1+\alpha} : L_1(x_0, X_0) \rightarrow L_1(x_0, X_0)$ ,  $I_{y_0}^{1+\beta} : L_1(y_0, Y_0) \rightarrow L_1(y_0, Y_0)$

$$u_1(x) = \sum_{n=0}^{\infty} I_{x_0}^{n(1+\alpha)} v_\alpha \mu^n = (E - \mu I_{x_0}^{1+\alpha})^{-1} v_\alpha = \mathcal{R}(\mu; I_{x_0}^{1+\alpha}) v_\alpha, \tag{3.8}$$

$$\begin{aligned}
 u_2(y) &= \sum_{n=0}^{\infty} I_{y_0}^{n(1+\beta)} v_{\beta} (\lambda - \mu)^n = (E - (\lambda - \mu) I_{y_0}^{1+\beta})^{-1} v_{\beta} \\
 &= \mathcal{R}(\lambda - \mu; I_{y_0}^{1+\beta}) v_{\beta},
 \end{aligned} \tag{3.9}$$

where  $E$  as usual is the identity operator. It's easily seen that series (3.8), (3.9) are analytic with respect to  $x \in (x_0, X_0)$  and  $y \in (y_0, Y_0)$ . Further, since (see (1.2))

$$\|I_{x_0}^{1+\alpha} f\|_{L_1(x_0, X_0)} \leq \frac{(X_0 - x_0)^{\alpha+1}}{\Gamma(1+\alpha)} \|f\|_{L_1(x_0, X_0)}$$

we have

$$\|\mathcal{R}(z; I_{x_0}^{1+\alpha})\| \leq \sum_{n=0}^{\infty} \|I_{x_0}^{1+\alpha}\|^n |z|^n \leq \sum_{n=0}^{\infty} \left( \frac{(X_0 - x_0)^{\alpha+1}}{\Gamma(1+\alpha)} \right)^n |z|^n.$$

Therefore the resolvent functions  $\mathcal{R}(z; I_{x_0}^{1+\alpha})$ ,  $\mathcal{R}(z; I_{y_0}^{1+\beta})$  are analytic in the open discs

$$|z| < \frac{\Gamma(1+\alpha)}{(X_0 - x_0)^{\alpha+1}}, \quad |z| < \frac{\Gamma(1+\beta)}{(Y_0 - y_0)^{\beta+1}},$$

respectively. Thus we write a family of eigenfunctions for the equation (2.4) in the resolvent form

$$u_{\lambda}(x, y) = \mathcal{R}(\mu; I_{x_0}^{1+\alpha}) v_{\alpha} \mathcal{R}(\lambda - \mu; I_{y_0}^{1+\beta}) v_{\beta}. \tag{3.10}$$

Indeed, substituting (3.10) into (2.4) taking into account the values  $D_{x_0+}^{1+\alpha} v_{\alpha} = D_{y_0+}^{1+\beta} v_{\beta} = 0$  after a simple change of the summation index into the series we easily satisfy equation (2.4). But we will extend our family of eigenfunctions considering

$$u_{\lambda}(x, y) = \mathcal{R}(\mu; I_{x_0}^{1+\alpha}) f \mathcal{R}(\lambda - \mu; I_{y_0}^{1+\beta}) g, \quad \mu \in \mathbb{C}, \tag{3.11}$$

with arbitrary  $f(x), g(y)$  such that  $I_{x_0}^{n(1+\alpha)+1-\alpha} f \in AC^2([x_0, X_0])$ ,  $I_{y_0}^{n(1+\beta)+1-\beta} g \in AC^2([y_0, Y_0])$  and the corresponding resolvent functions (3.8), (3.9) are analytic by  $x$  and  $y$ . So substituting (3.11) into (2.4) and ignoring trivial cases  $D_{x_0+}^{1+\alpha} f = D_{y_0+}^{1+\beta} g = 0$  which drive immediately to (3.11) with  $f = v_{\alpha}$ ,  $g = v_{\beta}$  (see (2.3)), after separation of variables we obtain fractional differential equations to define  $f, g$

$$D_{x_0+}^{1+\alpha} f = c \mathcal{R}(\mu; I_{x_0}^{1+\alpha}) f, \tag{3.12}$$

$$D_{y_0+}^{1+\beta} g = -c \mathcal{R}(\lambda - \mu; I_{y_0}^{1+\beta}) g, \tag{3.13}$$

where  $c$  is an arbitrary constant. Hence acting by inverse operators  $E - \mu I_{x_0}^{1+\alpha}$  and  $E - (\lambda - \mu) I_{y_0}^{1+\beta}$  on (3.12), (3.13) with the use of (2.3) we get, correspondingly,

$$D_{x_0+}^{1+\alpha} f = (\mu + c) f(x) + \mu v_{\alpha},$$

$$D_{y_0+}^{1+\beta} g = (\lambda - \mu - c)g(y) + (\lambda - \mu)v_\beta.$$

The latter equations are solved, for instance, in [3], [4] and we obtain the following solutions ( $c_i, d_i, i = 1, 2$  are constants)

$$\begin{aligned} f(x) = & c_1(x - x_0)^{\alpha-1} E_{1+\alpha, \alpha}((\mu + c)(x - x_0)^{1+\alpha}) \\ & + c_2(x - x_0)^\alpha E_{1+\alpha, 1+\alpha}((\mu + c)(x - x_0)^{1+\alpha}) \\ & + \mu a_1(x - x_0)^{2\alpha} E_{1+\alpha, 2\alpha+1}((\mu + c)(x - x_0)^{1+\alpha}) \\ & + \mu a_2(x - x_0)^{2\alpha+1} E_{1+\alpha, 2(\alpha+1)}((\mu + c)(x - x_0)^{1+\alpha}), \end{aligned} \quad (3.14)$$

$$\begin{aligned} g(y) = & d_1(y - y_0)^{\beta-1} E_{1+\beta, \beta}((\lambda - \mu - c)(y - y_0)^{1+\beta}) \\ & + d_2(y - y_0)^\beta E_{1+\beta, 1+\beta}((\lambda - \mu - c)(y - y_0)^{1+\beta}) \\ & + (\lambda - \mu)b_1(y - y_0)^{2\beta} E_{1+\beta, 2\beta+1}((\lambda - \mu - c)(y - y_0)^{1+\beta}) \\ & + (\lambda - \mu)b_2(y - y_0)^{2\beta+1} E_{1+\beta, 2(\beta+1)}((\lambda - \mu - c)(y - y_0)^{1+\beta}). \end{aligned} \quad (3.15)$$

Consequently, equation (2.4) has families of eigenfunctions (3.10) and (3.11) with  $f$  and  $g$  given by (3.14), (3.15). The case  $\lambda = 0$  naturally gives classical fundamental solutions  $u_{0,\mu}(x, y) = u_1(x)u_2(y)$  with  $u_1, u_2$ , for instance, in the form (3.4), (3.5).

**Remark 1.** Letting  $\alpha = \beta = 1$  in (3.4), (3.5) and using (1.8), (1.9) we obtain familiar trigonometric eigenfunctions of the Laplace equation  $\Delta u = \lambda u$ .

Returning again to functions  $f_i(y), h_i(x), i = 0, 1$  from Section 2 we suppose the following power-logarithmic analytic expansions in the neighbourhood of points  $y_0, x_0, 0 < r_1 \leq |y - y_0| \leq r_2 = |Y_0 - y_0|, 0 < \rho_1 \leq |x - x_0| \leq \rho_2 = |X_0 - x_0|$ , namely

$$f_i(y) = a_i \log(y - y_0) + (y - y_0)^{\mu_i} \sum_{k=0}^{\infty} a_{ik} (y - y_0)^k, \quad i = 1, 2, \quad \mu_i > -1, \quad (3.16)$$

$$h_i(x) = b_i \log(x - x_0) + (x - x_0)^{\nu_i} \sum_{k=0}^{\infty} b_{ik} (x - x_0)^k, \quad i = 1, 2, \quad \nu_i > -1. \quad (3.17)$$

Hence owing to [3] and straightforward calculations we get for each  $n \in \mathbb{N}_0$

$$\begin{aligned} (D_{y_0}^{n(1+\beta)} f_i)(y) &= a_i D_{y_0}^{n(1+\beta)} \log(y - y_0) + (y - y_0)^{\mu_i - n(1+\beta)} \\ &\times \frac{(-1)^n}{\pi} \sin(\pi(\{n\beta\} - \mu_i)) \sum_{k=0}^{\infty} (-1)^k a_{ik} \Gamma(k + \mu_i + 1) \Gamma(n(1 + \beta) - \mu_i - k) (y - y_0)^k, \\ (D_{x_0}^{n(1+\alpha)} h_i)(x) &= b_i D_{x_0}^{n(1+\alpha)} \log(x - x_0) + (x - x_0)^{\nu_i - n(1+\alpha)} \end{aligned}$$



$$\times \frac{(-1)^n}{\pi} \sin(\pi(\{n\alpha\} - \nu_i)) \sum_{k=0}^{\infty} (-1)^k b_{ik} \Gamma(k + \nu_i + 1) \Gamma(n(1 + \alpha) - \nu_i - k) (x - x_0)^k,$$

where  $i = 0, 1$ . Substituting the latter expressions into (2.28), (2.29) we get

$$\begin{aligned} u_f(x, y) = & \sum_{n=0}^{\infty} (-1)^n \frac{(x - x_0)^{n(1+\alpha)+\alpha-1}}{\Gamma(n(1 + \alpha) + \alpha)} \left[ a_0 + \frac{(x - x_0)a_1}{n(1 + \alpha) + \alpha} \right] D_{y_0}^{n(1+\beta)} \log(y - y_0) \\ & + (x - x_0)^{\alpha-1} (y - y_0)^{\mu_0} \frac{1}{\pi} \sum_{k,n=0}^{\infty} (-1)^k a_{0k} \sin(\pi(\{n\beta\} - \mu_0)) \\ & \times \frac{\Gamma(n(1 + \beta) - \mu_0 - k) \Gamma(k + \mu_0 + 1)}{\Gamma(n(1 + \alpha) + \alpha)} \left( \frac{(x - x_0)^{1+\alpha}}{(y - y_0)^{1+\beta}} \right)^n (y - y_0)^k \\ & + (x - x_0)^{\alpha} (y - y_0)^{\mu_1} \frac{1}{\pi} \sum_{k,n=0}^{\infty} (-1)^k a_{1k} \sin(\pi(\{n\beta\} - \mu_1)) \\ & \times \frac{\Gamma(n(1 + \beta) - \mu_1 - k) \Gamma(k + \mu_1 + 1)}{\Gamma((n + 1)(1 + \alpha))} \left( \frac{(x - x_0)^{1+\alpha}}{(y - y_0)^{1+\beta}} \right)^n (y - y_0)^k, \end{aligned} \quad (3.18)$$

$$\begin{aligned} u_h(x, y) = & \sum_{n=0}^{\infty} (-1)^n \frac{(y - y_0)^{n(1+\beta)+\beta-1}}{\Gamma(n(1 + \beta) + \beta)} \left[ b_0 + \frac{(y - y_0)b_1}{n(1 + \beta) + \beta} \right] D_{x_0}^{n(1+\alpha)} \log(x - x_0) \\ & + (y - y_0)^{\beta-1} (x - x_0)^{\nu_0} \frac{1}{\pi} \sum_{k,n=0}^{\infty} (-1)^k b_{0k} \sin(\pi(\{n\alpha\} - \nu_0)) \\ & \times \frac{\Gamma(n(1 + \alpha) - \nu_0 - k) \Gamma(k + \nu_0 + 1)}{\Gamma(n(1 + \beta) + \beta)} \left( \frac{(y - y_0)^{1+\beta}}{(x - x_0)^{1+\alpha}} \right)^n (x - x_0)^k \\ & + (y - y_0)^{\beta} (x - x_0)^{\nu_1} \frac{1}{\pi} \sum_{k,n=0}^{\infty} (-1)^k b_{1k} \sin(\pi(\{n\alpha\} - \nu_1)) \\ & \times \frac{\Gamma(n(1 + \alpha) - \nu_1 - k) \Gamma(k + \nu_1 + 1)}{\Gamma((n + 1)(1 + \beta))} \left( \frac{(y - y_0)^{1+\beta}}{(x - x_0)^{1+\alpha}} \right)^n (x - x_0)^k, \end{aligned} \quad (3.19)$$

where double series in (3.18), (3.19) are absolutely and uniformly convergent on the compact  $0 < r_1 \leq |y - y_0| \leq r_2 = |Y_0 - y_0|$ ,  $0 < \rho_1 \leq |x - x_0| \leq \rho_2 = |X_0 - x_0|$  owing to conditions

$$\sum_{k,n=0}^{\infty} \frac{|a_{1k} \Gamma(n(1 + \beta) - \mu_1 - k) \Gamma(k + \mu_1 + 1)|}{\Gamma((n + 1)(1 + \alpha))} \left( \frac{\rho_2^{1+\alpha}}{r_1^{1+\beta}} \right)^n r_2^k < \infty, \quad (3.20)$$

$$\sum_{k,n=0}^{\infty} \frac{|b_{1k}\Gamma(n(1+\alpha) - \nu_1 - k)|\Gamma(k + \nu_1 + 1)}{\Gamma((n+1)(1+\beta))} \left( \frac{r_2^{1+\beta}}{\rho_1^{1+\alpha}} \right)^n \rho_2^k < \infty. \quad (3.21)$$

The ordinary series in (3.18), (3.19) we will call fractional logarithmic solutions. Taking into account, for instance, the representation (see (1.1))

$$\begin{aligned} D_{y_0}^{n(1+\beta)} \log(y - y_0) &= \left( \frac{d}{dy} \right)^{[n(1+\beta)]+1} [(y - y_0)^{1-\{n\beta\}} \\ &\quad \times \int_0^1 (1-t)^{-\{n\beta\}} \log((y - y_0)t) dt] \\ &= (-1)^{[n(1+\beta)]+1} (y - y_0)^{-n(1+\beta)} \left[ \frac{\Gamma(n(1+\beta))}{\Gamma(\{n\beta\} - 1)} \left[ d_n - \frac{\log(y - y_0)}{\{n\beta\} - 1} \right] \right. \\ &\quad \left. + \frac{([n(1+\beta)] + 1)!}{\Gamma(\{n\beta\})} \sum_{m=1}^{[n(1+\beta)]+1} \frac{\Gamma(n(1+\beta) - m)}{m([n(1+\beta)] + 1 - m)!} \right], \end{aligned}$$

where  $d_n = \int_0^1 (1-t)^{-\{n\beta\}} \log t \, dt$ , we may substitute it in (3.18) to write the ordinary series in a different form and to guarantee its absolute convergence in the region  $(x - x_0)^{1+\alpha}/(y - y_0)^{1+\beta} < 1$ . Finally we note that in the similar manner we treat the ordinary series in (3.19).

## References

- [1] V. A. Ditkin and A. P. Prudnikov, *Integral Transforms and Operational Calculus*, Pergamon Press, Oxford, 1965.
- [2] M. M. Dzhrbashyan, *Integral Transforms and Representations of Functions in a Complex Domain*, Nauka, Moscow, 1966 (in Russian).
- [3] S. G. Samko, A. A. Kilbas and O. I. Marichev, *Fractional Integrals and Derivatives. Theory and Applications*, Gordon and Breach, Yverdon *et alibi*, 1993.
- [4] S.B. Yakubovich and Yu.F. Luchko, *The Hypergeometric Approach to Integral Transforms and Convolutions*, (Kluwers Ser. Math. and Appl.: vol. 287). Dordrecht, Boston, London, 1994.
- [5] A.H. Zemanian, *Generalized Integral Transformations*, Dover Publ., New York, 1987.